



UNIVERSITY OF  
ILLINOIS LIBRARY  
AT URBANA-CHAMPAIGN  
BIOLOGY

1111 . . .





# FIELDIANA

6

## Zoology

NEW SERIES, NO. 35

### Catalogue of Type Specimens of Recent Fishes in Field Museum of Natural History

Myriam Ibarra  
Donald J. Stewart

THE LIBRARY OF THE  
NOV 9 1987  
UNIVERSITY OF ILLINOIS  
URBANA-CHAMPAIGN

BIOLOGY LIBRARY  
101 DOWELL HALL

September 30, 1987  
Publication 1377

PUBLISHED BY FIELD MUSEUM OF NATURAL HISTORY



Field Museum of Natural History  
Roosevelt Road at Lake Shore Drive  
Chicago, Illinois 60605-2496  
Telephone: (312) 922-9410

---

---

# FIELDIANA

---

---

## Zoology

NEW SERIES, NO. 35

### **Catalogue of Type Specimens of Recent Fishes in Field Museum of Natural History**

**Myriam Ibarra and Donald J. Stewart**

*Division of Fishes  
Department of Zoology  
Field Museum of Natural History  
Chicago, Illinois 60605-2496*

*Present address:  
Center for Limnology  
University of Wisconsin  
Madison, Wisconsin 53706*

**Accepted for publication August 7, 1985**

**September 30, 1987**

**Publication 1377**

---

---

**PUBLISHED BY FIELD MUSEUM OF NATURAL HISTORY**

---

---

© 1987 Field Museum of Natural History  
*Library of Congress Catalog Card Number: 87-81351*  
ISSN 0015-0754  
PRINTED IN THE UNITED STATES OF AMERICA



Table of Contents

ABSTRACT ..... 1

INTRODUCTION ..... 1

FORMAT OF THE LIST ..... 3

LIST OF TYPES

    Arranged Alphabetically by Genus ..... 4

LIST OF TAXA REPRESENTED BY TYPES

    Arranged Alphabetically by Family ..... 86

ACKNOWLEDGMENTS ..... 104

LITERATURE CITED ..... 104

SPECIES INDEX ..... 105

List of Illustrations

1. Frequency histogram for number of species of Recent fishes described in each year through 1984 that are represented by type material in Field Museum of Natural History ..... 2



# Catalogue of Type Specimens of Recent Fishes in Field Museum of Natural History

---

## Abstract

We list 1,406 nominal taxa of Recent fishes for which there are type specimens present in Field Museum of Natural History, including 837 holotypes, eight lectotypes, three neotypes, 48 taxa represented by syntypes, and 510 taxa represented only by secondary types. A large proportion of the species was described between 1905 and 1930 by Seth E. Meek, Samuel F. Hildebrand, Carl H. Eigenmann, David S. Jordan, and their associates. About 500 primary types came to Field Museum in 1952 with the purchase of the Carnegie Museum of Natural History fish collection. The main list of types presented here is arranged alphabetically by genus; each entry includes genus, species, and subspecies (if any) as it was originally proposed; author(s) and year of publication; family where the taxon is presently classified; abbreviated literature citation to original description; current type status; catalogue number(s) for primary types; total number of syntypes and secondary types; and brief locality information. In a few cases, comments are included to indicate discrepancies in previously published spellings or catalogue numbers, citations to subsequent designations of neotypes and lectotypes, and status of the material if uncertain. Also included are a list of the same taxa arranged by family and an alphabetical index of species.

## Introduction

Field Museum of Natural History is one of seven recognized international depositories of Recent fishes in North America and is ranked among the

top five in the United States and Canada with respect to its type holdings (Collette & Lachner, 1976; Nitecki, 1980). The collection includes type material for 1,406 nominal taxa, of which 896 are represented by primary types (including 837 holotypes, eight lectotypes, three neotypes, and 48 taxa represented by syntypes). This catalogue was prepared in response to Recommendation 17D(4) of the International Code of Zoological Nomenclature (ICZN) to publish information on holdings of type materials. It also stems from our sense of obligation to the international community of systematic ichthyologists to provide the best possible curation for the types that have been entrusted to this institution since it was founded in 1893.

The type specimens of Recent fishes in Field Museum's collection stem from (1) exploratory fieldwork and systematic research of curators and associates, (2) purchase of the fish collection of the Carnegie Museum, Pittsburgh, in 1952, and (3) gifts and exchanges received from researchers throughout the world. Acquisition of the Carnegie Museum collection, with about 500 taxa represented by primary types, more than doubled the type collection at Field Museum. Most of the taxa in the collection were described after 1900, and a large proportion of them between 1905 and 1930 (fig. 1). Most of the specimens in our collection from that time period were described by Seth E. Meek (Field Museum), Samuel F. Hildebrand (United States National Museum), Carl H. Eigenmann (Indiana University), David S. Jordan (Stanford University), and their associates.

This catalogue represents a synthesis of information presented in two previously published type lists (Henn, 1928; Grey, 1947) with that in our catalogues and associated recent literature. Henn (1928) reported only the primary types in the Car-




FIG. 1. Frequency histogram for number of species of Recent fishes described in each year through 1984 that are represented by type material in Field Museum of Natural History.

negie Museum collection, while Grey (1947) listed all types in Field Museum. These two lists cover about 1,000 of our 1,406 taxa. In preparing this catalogue, the published lists and subsequent catalogue entries were checked against all jars on the shelf and outstanding loan invoices. A first draft of this list was then prepared, and nearly every entry was checked against the original publication. Those few taxa for which we were not able to get a copy of the original publication were included only if the names were clearly established in the literature; references for all such taxa (i.e., five species) are indicated as "not seen."

This list includes the types of all taxa present in Field Museum's fish collection that, to our knowledge, were described by mid-1985 when the final revision of this catalogue was submitted. It seems likely that a few other species based in part on FMNH materials were published in the last few months, but that we have not received a copy of the paper(s). The fish collections of Field Museum include specimens for about 50 taxa which have been labeled "manuscript types", none of which were included in this list. We were able to trace most of these hopeful taxa to determine their present status, but a few remain unresolved. We urge colleagues who find omissions or other errors in this list to notify Division of Fishes staff at Field Museum so that records can be updated.

## Format of the List

The list of types is arranged alphabetically by genus and, within each genus, by species or subspecies as it was originally proposed without subsequent relocation. Information included for each species is as follows:

**GENUS**—This is spelled as it appears in the original description of the species, except where an emendation is clearly called for according to the rules of the ICZN, or where the author apparently used an inappropriate emendation of a previously published generic name. In almost every such case, the spelling used here has appeared in the recent literature. When the spelling used here differs from that used in the species description, the presumed incorrect spelling is given in a comment at the end of the entry. Subgeneric assignments were included in some species descriptions, but are omitted here.

**SPECIES AND SUBSPECIES**—These are spelled as they appear in the original description, except that

diacritical marks and hyphens have been deleted as recommended by the ICZN (i.e., hyphens are retained only in species names such as *x-punctata*). In a few cases, the original publication included a typographic error; such names are corrected to agree with current usage, and a comment to that effect is added at the end of the entry.

**AUTHOR(S) AND DATE**—The surname of the author(s) of the species is given followed by the year in which the publication was issued.

**LITERATURE REFERENCE**—Title of the journal where the description was published is abbreviated (using BIOSIS, 1981, where possible) and followed by volume number, issue number in parentheses, and page number on which the description begins. Some book titles are abbreviated. When authorship of the publication differed from that of the species described within, author(s) of the publication precedes the title of the journal or book.

**PRIMARY TYPES**—Categories used here for primary types are holotype, syntype(s), lectotype, and neotype. Field Museum of Natural History (FMNH) catalogue numbers are given for all primary types, and for former Carnegie Museum (CM) types, catalogue numbers are given in parentheses after FMNH number. For syntypes, the total number of specimens is given just ahead of the catalogue number(s). Missing types are indicated as such, and omission of a catalogue number means simply that no catalogue entry was ever made for the missing lot. A question mark indicates some uncertainty about type status of our material, and the problem is discussed at the end of the entry. If a lectotype or neotype has been designated, the relevant literature reference is given. Finally, the type locality is presented in general terms. Geographic and geopolitical place names have been changed in some cases to agree with modern usage (e.g., Guyana for British Guiana). We follow Bartholomew et al. (1980) for spellings of many of the place names.

**SECONDARY TYPES**—Categories of secondary types used here are paratype, lectoparatype, and allotype. Allotypes are considered equivalent to paratypes, but they are catalogued separately in the collection. The total number of secondary types is given followed by brief locality information. We consider secondary types to be all materials other than primary types that were used by the author to describe the taxon. Many early workers and occasionally recent workers gave incomplete data on the secondary types, and deciding which should be considered as such is, in some cases, subjective.

We excluded any material for which there was not at least a vague reference to the collection locality in the original description. In this regard, we differ from Grey (1947) who included materials from localities not given in the original description. We also excluded specimens which were not explicitly called types in the original description *and* indicated by the author as differing in some way from the other types.

**FAMILY**—Assignment of each taxon to a family and spelling of family names follows Nelson (1984) for most entries, because that work is the most recent and comprehensive synthesis dealing with all of the world's fish families. We differ from Nelson (1984) and follow Robins et al. (1980) for the more widely used spellings of Cobitidae, Eleotridae, Engraulidae, and Paralepididae.

## List of Types

### Arranged Alphabetically by Genus

**Abudefduf lorenzi** Hensley and Allen, 1977

Rec. W. Aust. Mus., 6(1): 109.

Holotype: FMNH 23564. Solomon Islands: Isabel Island, Tenibuli Bay.

Paratypes: 2. Solomon and Philippine Islands: 2 localities.

POMACENTRIDAE

**Acahara jusanensis** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 180.

Holotype: FMNH 58722 (CM 7828). Japan: Lake Jusan near Aomori.

CYPRINIDAE

**Acanthemblemaria greenfieldi** Smith-Vaniz and Palacio, 1974

Proc. Acad. Nat. Sci. Phila., 125(11): 210.

Paratypes: 7. Belize: Glover's Reef.

CHAENOPSIDAE

**Acanthicus canensis** Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 80.

Holotype: FMNH 7581. Panama: Río Cana, Cana.

Paratypes: 30. Panama: 2 localities.

LORICARIIDAE

**Acanthocharax microlepis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 405.

Holotype: FMNH 54475 (CM 2138). Guyana: Tumatumari.

Paratypes: 31. Guyana: 3 localities.

CHARACIDAE

**Acanthophaelus bifurcus** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 52.

Holotype: FMNH 53539 (CM 1088). Guyana: Christianburg.

Paratypes: 36. Guyana: 2 localities.

POECILIIDAE

**Acanthophaelus melanzonus** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 51.

Holotype: FMNH 52717 (CM 1086). Guyana: Georgetown Trenches.

Paratypes: 6. Same locality.

POECILIIDAE

**Acanthopthalmus mariae** Inger and Chin, 1962

Fieldiana Zool., 45: 118.

Holotype: FMNH 68161. Malaysia: Sabah, tributary of Kinabatangan at Deramakot.

Paratypes: 112. Malaysia: Sabah, 2 localities.

COBITIDAE

**Acanthopthalmus sandakanensis** Inger and Chin, 1962

Fieldiana Zool., 45: 120.

Holotype: FMNH 68158. Malaysia: Sabah, Sepilok Forest Reserve.

Paratypes: 36. Malaysia: Sabah, 4 localities.

COBITIDAE

**Acanthoplesiops hiatti** Schultz, 1953

Bull. U.S. Natl. Mus., 202(1): 407.

Paratype. Marshall Islands: Bikini Atoll, Boro Island.

ACANTHOCLINIDAE

**Acanthurus randalli** Briggs and Caldwell, 1957

Bull. Fla. State Mus. Biol. Sci., 2(4): 43.

Paratype. Florida: Bay County near Panama City.

ACANTHURIDAE

**Acanthurus triostegus marquesensis** Schultz and Woods, 1948

J. Wash. Acad. Sci., 38(7): 250.

Paratypes: 5. Marquesas Islands: 2 localities.

ACANTHURIDAE

**Acestridium discus** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 319.

Holotype: FMNH 54339 (CM 2985). Brazil: Igarapé de Cachoeira Grande, near Manaus.

Paratype. Same locality.

LORICARIIDAE

- Acestrorhynchus altus** Menezes, 1969  
Arq. Zool. (São Paulo), 18(1–2): 52.  
Paratypes: 10. Brazil: Mato Grosso, Descalvados.  
CHARACIDAE
- Acestrorhynchus guianensis** Menezes, 1969  
Arq. Zool. (São Paulo), 18(1–2): 70.  
Holotype: FMNH 74359. Guyana: Botanic Garden.  
Paratypes: 7. Guyana: 4 localities.  
CHARACIDAE
- Acestrorhynchus nasutus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 411.  
Holotype: FMNH 53475 (CM 1959). Guyana: Rockstone.  
CHARACIDAE
- Acheilognathus barbatus** Nichols, 1926  
Am. Mus. Nov., 214: 6.  
Paratype. China: Anhwei, Ningkwo.  
CYPRINIDAE
- Acheilognathus gracilis** Nichols, 1926  
Am. Mus. Nov., 214: 5.  
Paratype. China: Hunan, Tungting Lake.  
CYPRINIDAE
- Acheilognathus tabira** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 220.  
Holotype: FMNH 57071 (CM 6007). Japan: Lake Biwa at Matsubara.  
Paratypes: 41. Same locality.  
CYPRINIDAE
- Achirus fluviatilis** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 1002.  
Paratypes: 40. Panama: 4 localities.  
SOLEIDAE
- Acipenser oxyrhynchus desotoi** Vladykov, 1955  
J. Fish. Res. Board Can., 12(5): 755.  
Paratype. Mississippi: off Gautier at mouth of Singing River.  
ACIPENSERIDAE
- Adioryx poco** Woods, 1965  
Not. Nat. (Phila.), 377: 1.  
Paratype. Grand Cayman Island: north side of Georgetown.  
HOLOCENTRIDAE
- Adontosternarchus clarkae** Mago-Leccia, Lundberg, and Baskin, 1985  
Nat. Hist. Mus. Los Ang. Cty. Contrib. Sci., 358: 14.  
Paratypes: 6. Colombia and Ecuador: 2 localities.  
APTERONOTIDAE
- Adontosternarchus devenanzii** Mago-Leccia, Lundberg, and Baskin, 1985  
Nat. Hist. Mus. Los Ang. Cty. Contrib. Sci., 358: 11.  
Paratypes: 12. Venezuela: Río Orinoco, north of Isla Tres Caños.  
APTERONOTIDAE
- Aequidens awani** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 335.  
Holotype: FMNH 54036 (CM 2576). Brazil: São Antonio de Guaporé.  
Paratype. Same locality.  
CICHLIDAE
- Aequidens azurifer** Fowler, 1911  
Proc. Acad. Nat. Sci. Phila., 63: 515.  
Paratype. Ecuador: near Bucay.  
CICHLIDAE
- Aequidens chimantanus** Inger, 1956  
Fieldiana Zool., 34(37): 437.  
Holotype: FMNH 45702. Venezuela: Río Abácapa, west side of Chimantá-tepui.  
Paratypes: 2. Same locality.  
CICHLIDAE
- Aequidens duopunctata** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 338.  
Holotype: FMNH 54033 (CM 2573). Brazil: Manaus.  
Paratype. Same locality.  
CICHLIDAE
- Aequidens guaporensis** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 335.  
Holotype: FMNH 54035 (CM 2575). Brazil: São Antonio de Guaporé.  
CICHLIDAE
- Aequidens mariaae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 240.  
Paratypes: 3. Colombia: Río Negro at Villaviciencio.  
CICHLIDAE

**Aequidens plagiozonatus** Kullander, 1984

Zool. Scr., 13(2): 155.

Paratype. Brazil: Descalvados.

CICHLIDAE

**Aequidens potaroensis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 490.

Holotype: FMNH 53892 (CM 2407). Guyana: Amatuk.

Paratypes: 79. Guyana: 14 localities.

CICHLIDAE

**Ageneiosus guianensis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 204.

Holotype: FMNH 53247 (CM 1712). Guyana: Wismar.

AGENEIOSIDAE

**Ageneiosus madeirensis** Fisher, 1917

Ann. Carnegie Mus., 11(3-4): 426.

Holotype: FMNH 58143 (CM 7143). Bolivia: San Joaquín.

Paratypes: 14. Same locality.

AGENEIOSIDAE

**Ageneiosus marmoratus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 206.

Holotype: FMNH 53245 (CM 1710). Guyana: creek below Potaro Landing.

AGENEIOSIDAE

**Agmus lyriformis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 128.

Holotype: FMNH 53120 (CM 1554). Guyana: Gluck Island.

ASPREDINIDAE

**Agoniates anchovia** Eigenmann, 1914

Indiana Univ. Stud., 19: 46.

Holotype: FMNH 56401 (CM 5216). Bolivia: Villa Bella.

Paratypes: 8. Same locality.

CHARACIDAE

**Agosia adobe** Jordan and Evermann, 1891

Bull. U.S. Fish Comm. 1889, 9: 36.

Paratypes: 3. Utah: Sevier River near Juab.

CYPRINIDAE

**Agosia yarrowi** Jordan and Evermann, 1891

Bull. U.S. Fish Comm. 1889, 9: 28.

Syntypes: 21, FMNH 1878, 6688. Colorado: Tomichi Creek, Gunnison.

CYPRINIDAE

**Albula virgata** Jordan and Jordan, 1922

Mem. Carnegie Mus., 10(1): 6.

Holotype: FMNH 55190 (CM 3896). Hawaiian Islands: Honolulu.

ALBULIDAE

**Alburnops illecebrosus** Girard, 1857

Proc. Acad. Nat. Sci. Phila., 1856: 194.

Lectoparatype. Arkansas: Arkansas River, Fort Smith.

Lectotype designated by Meek (In: Jordan, 1885, Proc. U.S. Natl. Mus., 8(500): 123).

CYPRINIDAE

**Alectis breviventralis** Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 216.

Holotype: FMNH 59411 (CM 7753). Japan: Kii. Paratype. Same locality.

CARANGIDAE

**Alectis temmincki** Wakiya, 1924

Ann. Carnegie Mus., 15(2-3): 215.

Holotype: FMNH 59410 (CM 7752). Japan: Bonin Islands.

CARANGIDAE

**Alepideleotris tigris** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 413.

Holotype: FMNH 17348. Ecuador: Islas Galápagos, Isla South Seymour.

ELEOTRIDAE

**Alepocephalus owstoni** Tanaka, 1908

J. Coll. Sci. Imp. Univ. Tokyo, 23(7): 27.

Holotype: FMNH 89529. Japan: Sagami Sea outside Okinose.

ALEPOCEPHALIDAE

**Alepocephalus umbriceps** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 209.

Holotype: (CM 6030) MISSING. Japan: Aomori.

ALEPOCEPHALIDAE

**Alestes langi** Fowler, 1935

Ann. Transvaal Mus., 16(2): 258.

Paratypes: 3. Botswana: Thamalakane River at Maun.

CHARACIDAE

**Alestes thamalakanensis** Fowler, 1935

Ann. Transvaal Mus., 16(2): 257.

Paratypes: 5. Botswana: Thamalakane River at Maun.

CHARACIDAE



**Alfaro acutiventralis** Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 72.  
Holotype: FMNH 7679. Costa Rica: Guapilis.  
Allotype and paratypes: 31. Same locality.  
POECILIIDAE

**Algansea aphanea** Barbour and Miller, 1978

Misc. Publ. Mus. Zool. Univ. Mich., 155: 21.  
Paratypes: 10. Mexico: Jalisco, Río Ayutla.  
CYPRINIDAE

**Algansea monticola** Barbour and Contreras, 1968

Proc. Biol. Soc. Wash., 81: 101.  
Paratypes: 2. Mexico: Zacatecas, Río Juchipila  
near Jalpa.  
CYPRINIDAE

**Algansea rubescens** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 78.  
Holotype: FMNH 3653. Mexico: Río Santiago  
at Ocotlán.  
Paratypes: 8. Same locality.  
CYPRINIDAE

**Allanetta crenolepis** Schultz, 1953

Bull. U.S. Natl. Mus., 202(1): 302.  
Paratypes: 3. Philippine Islands: Tara Island.  
ATHERINIDAE

**Allolepis hollandi** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 323.  
Holotype: FMNH 58842 (CM 7952). Sea of Ja-  
pan: near Fukui.  
ZOARCIDAE

**Alosa ohiensis** Evermann, 1902

Rept. U.S. Fish Comm. 1901, 27: 277.  
Paratype. Kentucky: Ohio River at Louisville.  
CLUPEIDAE

**Alticus novemmaculosus** Snyder, 1908

Proc. U.S. Natl. Mus., 35(1635): 107.  
Paratypes: 129. Japan: Ryukyu Islands, Oki-  
nawa, Naha.  
BLENNIIDAE

**Ambassis lafa** Jordan and Seale, 1906

Bull. Bur. Fish. 1905, 25: 253.  
Paratypes: 16. Samoan Islands: Apia.  
CENTROPOMIDAE

**Ambassis vaivasensis** Jordan and Seale, 1906

Bull. Bur. Fish. 1905, 25: 254.  
Paratypes: 3. Samoan Islands: Vaivase River,  
Apia.  
CENTROPOMIDAE

**Amblygobius myersi** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 426.  
Holotype: FMNH 17352. Solomon Islands: New  
Georgia.  
GOBIIDAE

**Amblypharyngodon grandisquamis** Jordan and  
Starks, 1917

Ann. Carnegie Mus., 11: 438.  
Holotype: FMNH 58964 (CM 8050). Sri Lanka:  
Ceylon River at Colombo.  
Paratypes: 17. Same locality.  
CYPRINIDAE

**Ameiurus australis** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 13.  
Holotype: FMNH 4474. Mexico: Tamaulipas,  
Forlón.  
Spelled *Amiurus* in Meek (1904).  
ICTALURIDAE

**Ameiurus mexicanus** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 15.  
Holotype: FMNH 4507. Mexico: San Luis Po-  
tosí, Rascón.  
Paratypes: 3. Mexico: San Luis Potosí, 2 local-  
ities.  
Spelled *Amiurus* in Meek (1904).  
ICTALURIDAE

**Amia sialis** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6: 246.  
Holotype: FMNH 57084 (CM 6021). Japan: Su-  
ruga Bay.  
APOGONIDAE

**Ammocryptocharax elegans** Weitzman and Ka-  
nazawa, 1976

Proc. Biol. Soc. Wash., 89(26): 331.  
Paratypes: 7. Colombia: Río Vichada basin,  
Caño Muco west of Puerto Gaitán.  
CHARACIDAE

**Anarchias yoshiae** Kanazawa, 1952

Fieldiana Zool., 34(7): 75.  
Holotype: FMNH 48729. Bermuda: ship's  
channel.  
MURAENIDAE

**Anchoa hepsetus colonensis** Hildebrand, 1943

Bull. Bingham Oceanogr. Collect. Yale Univ.,  
8(2): 60.  
Paratypes: 12. Jamaica.  
ENGRAULIDAE

**Anchoa lamprotaenia** Hildebrand, 1943  
Bull. Bingham Oceanogr. Collect. Yale Univ.,  
8(2): 62.  
Paratypes: 86. Gulf of Mexico and Caribbean  
Sea: 7 localities.  
ENGRAULIDAE

**Anchoa mitchilli diaphana** Hildebrand, 1943  
Bull. Bingham Oceanogr. Collect. Yale Univ.,  
8(2): 91.  
Paratypes: 10. Louisiana: Grande Isle.  
ENGRAULIDAE

**Anchovia arenicola** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 201.  
Paratypes: 122. Panama: 4 localities.  
ENGRAULIDAE

**Anchovia brevirostra** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 198.  
Paratypes: 10. Panama: Canal Zone, Balboa Bay  
at Balboa.  
ENGRAULIDAE

**Anchovia eigenmannia** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 205.  
Paratypes: 7. Panama: Taboga Island, Balboa.  
ENGRAULIDAE

**Anchovia elongata** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 198.  
Paratypes: 3. Panama: Canal Zone, Mindi Cut.  
ENGRAULIDAE

**Anchovia parva** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 202.  
Paratypes: 24. Panama: 2 localities.  
ENGRAULIDAE

**Anchoviella belizensis** Thomerson and Greenfield,  
1975  
Copeia, 1975(1): 50.  
Holotype: FMNH 78363. Belize: Sibun River.  
Paratypes: 35. Same locality.  
ENGRAULIDAE

**Anchoviella helleri** Hubbs, 1921  
Proc. Biol. Soc. Wash., 34: 47.  
Holotype: FMNH 3332. Mexico: Baja California,  
Bahía San Felipe.  
Paratypes: 2. Same locality.  
ENGRAULIDAE

**Ancistrus lithurgicus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 241.  
Holotype: FMNH 53091 (CM 1524). Guyana:  
Crab Falls.  
Paratypes: 5. Same locality.  
LORICARIIDAE

**Ancistrus melas** Eigenmann, 1916  
Ann. Carnegie Mus., 10(1-2): 83.  
Holotype: FMNH 58339 (CM 7335). Colombia:  
Condoto.  
Paratype. Colombia: Raspadura.  
LORICARIIDAE

**Ancistrus planiceps** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 79.  
Holotype: FMNH 7580. Panama: Río Tuyra,  
Boca de Cupe.  
Paratypes: 16. Panama: 6 localities.  
LORICARIIDAE

**Ancistrus spinosus** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 252.  
Holotype: FMNH 8942. Panama: Río Calobre.  
LORICARIIDAE

**Ancylosetta antillarum** Gutherz, 1966  
Bull. Mar. Sci., 16(3): 449.  
Paratypes: 2. Caribbean Sea: 2 localities.  
BOTHIDAE

**Ancylosetta cycloidea** Tyler, 1959  
Copeia, 1959(2): 140.  
Paratypes: 9. Western Atlantic Ocean: 2 localities.  
BOTHIDAE

**Ancylosetta kumperae** Tyler, 1959  
Copeia, 1959(2): 143.  
Paratypes: 2. Western Atlantic Ocean: 2 localities.  
BOTHIDAE

**Ancylosetta microctenus** Gutherz, 1966  
Bull. Mar. Sci., 16(3): 455.  
Paratypes: 2. Caribbean Sea.  
BOTHIDAE

**Anduzedoras arleo** Fernández-Yépez, 1968  
Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 29.  
Holotype: FMNH 84068. Venezuela: Río  
Autana, east of Puerto Ayacucho.  
DORADIDAE

**Anduzedoras copei** Fernández-Yépez, 1968  
Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 31.  
Holotype: FMNH 84069. Venezuela: Lagoon at  
side of Río Capanaparo.  
DORADIDAE

**Anisotremus williamsi** Meek, 1905  
Proc. Biol. Soc. Wash., 18: 241.  
Holotype: FMNH 3350. Brazil: Santos.  
Paratypes: 3. Same locality.  
Misspelled *Anisotrema* in Meek (1905); *wil-*  
*liamsoni* in Grey (1947, p. 116).  
HAEMULIDAE

**Anoplarchus purpureus archolepis** Hubbs, 1926  
Pap. Mich. Acad. Sci. Arts and Lett., 7: 375.  
Paratypes: 34. California: 6 localities.  
Misspelled *purpurascens* in Grey (1947, p.  
118).  
STICHAEIDAE

**Anostomus plicatus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 296.  
Holotype: FMNH 53393 (CM 1861). Guyana:  
Crab Falls.  
Paratypes: 3. Guyana: 3 localities.  
ANOSTOMIDAE

**Antennarius bermudensis** Schultz, 1957  
Proc. U.S. Natl. Mus., 107(3383): 98.  
Paratype: Bermuda: Harrington Sound.  
ANTENNARIIDAE

**Antennarius pauciradiatus** Schultz, 1957  
Proc. U.S. Natl. Mus., 107(3383): 100.  
Paratypes: 2. Florida: south end of Biscayne Bay.  
ANTENNARIIDAE

**Antennarius verrucosus** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 31.  
Holotype: FMNH 4853. Bermuda: Saint  
George's Island.  
ANTENNARIIDAE

**Anthias conspicuus** Heemstra, 1973  
Copeia, 1973(2): 200.  
Paratypes: 12. India: Arabian Sea, off Bombay.  
SERRANIDAE

**Anthias mooreanus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 405.  
Holotype: FMNH 17325. Society Islands: Moo-  
rea Island.  
SERRANIDAE

**Anthias woodsi** Anderson and Heemstra, 1980  
Copeia, 1980(1): 74.  
Holotype: FMNH 70709. Straits of Florida: off  
Dry Tortugas.  
Paratypes: 2. Straits of Florida off Dry Tortugas:  
2 localities.  
SERRANIDAE

**Antigonía combatia** Berry and Rathjen, 1958  
Q. J. Fla. Acad. Sci., 21(3): 255.  
Paratypes: 52. Gulf of Mexico and off Cuba: 9  
localities.  
CAPROIDAE

**Antigonía malayana** Weber, 1913  
Die Fische der Siboga-Expedition, Leiden, p.  
299.  
Syntype: FMNH 52432 (CM 689). Arafura Sea.  
CAPROIDAE

**Apareiodon hasemani** Eigenmann, 1916  
Ann. Carnegie Mus., 10(1–2): 75.  
Holotype: FMNH 57615 (CM 6587). Brazil:  
Pirapora.  
Paratypes: 59. Brazil: 5 localities.  
PARODONTIDAE

**Apareiodon itapicuruensis** Eigenmann and Henn,  
1916  
Ann. Carnegie Mus., 10(1–2): 72.  
Holotype: FMNH 56988 (CM 5804). Brazil: Rio  
Itapicuru basin, Rio Paiaia.  
Paratypes: 7. Brazil: 2 localities.  
PARODONTIDAE

**Aparrius aurocingulus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 425.  
Holotype: FMNH 17353. Fiji Islands: Ovalau  
Island.  
Paratypes: 2. Same locality.  
GOBIIDAE

**Aphyocharax anisitsi** Eigenmann and Kennedy,  
1903  
Proc. Acad. Nat. Sci. Phila., 59: 517.  
Paratypes: 2. Paraguay: Campo Grande.  
CHARACIDAE

**Aphyocharax erythrurus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 313.  
Holotype: FMNH 53579 (CM 1879). Guyana:  
Rockstone sand-bank.  
Paratypes: 14. Guyana: 3 localities.  
CHARACIDAE

- Aphyocharax melanotus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 312.  
Holotype: FMNH 53578 (CM 1877). Guyana:  
Rockstone sand-bank.  
CHARACIDAE
- Aphyocharax paraguayensis** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 33.  
Holotype: FMNH 57922 (CM 6906). Brazil:  
Cáceres.  
Paratype. Same locality.  
CHARACIDAE
- Aphyocheiroidon hemigrammus** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 59.  
Holotype: FMNH 57819 (CM 6802). Brazil: Ja-  
quara.  
Paratypes: 25. Brazil: 3 localities.  
CHARACIDAE
- Aphyodite grammica** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 314.  
Holotype: FMNH 53408 (CM 1882). Guyana:  
Konawaruk.  
CHARACIDAE
- Apistogramma caetei** Kullander, 1980  
Bonn. Zool. Monogr., 14: 76.  
Holotype: FMNH 54164 (CM 2732). Brazil:  
Pará, Igarapé in Bragança.  
Paratypes: 2. Same locality.  
CICHLIDAE
- Apistogramma inconspicua** Kullander, 1982  
Zool. Scr., 11(4): 307.  
Paratypes: 3. Brazil: São Luiz de Cáceres.  
CICHLIDAE
- Apistogramma macmasteri** Kullander, 1979  
Zool. Scr., 8: 70.  
Paratype. Colombia: Villavicencio.  
CICHLIDAE
- Aplatophis chauliodus** Böhlke, 1956  
Not. Nat. (Phila.), 289: 3.  
Holotype: FMNH 61772. Puerto Rico: Maya-  
güez pier.  
OPHICHTHIDAE
- Aplocheilus chobensis** Fowler, 1935  
Ann. Transvaal Mus., 16(2): 276.  
Paratypes: 4. Botswana: Chobe River at Ka-  
bulabula.  
APLOCHEILIDAE
- Aplocheilus luzonensis** Herre and Ablan, 1934  
Philipp. J. Sci., 54(2): 275.  
Paratypes: 4. Philippine Islands: Luzon, Ilocos  
Norte, Solsona.  
APLOCHEILIDAE
- Apogon abo** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 402.  
Holotype: FMNH 17307. New Guinea: Sepik  
River at Koragu.  
Paratypes: 3. New Guinea: 2 localities.  
APOGONIDAE
- Apogon beauforti** Weber, 1908  
Nova Guinea, 5(2): 246.  
Syntypes: 3, FMNH 52390 (CM 646). New  
Guinea: Lake Sentani.  
APOGONIDAE
- Apogon leptocaulus** Gilbert, 1972  
Proc. Biol. Soc. Wash., 85(36): 421.  
Paratype. Belize: Glover's Reef.  
APOGONIDAE
- Apogon nigrofasciatus** Lachner, 1953  
Bull. U.S. Natl. Mus., 202(1): 466.  
Paratypes: 3. Marshall Islands: Bikini Atoll, 2  
localities.  
APOGONIDAE
- Apogon snyderi** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 180.  
Paratypes: 2. Hawaiian Islands: Hilo.  
APOGONIDAE
- Apogon wichmanni** Weber, 1908  
Nova Guinea, 5(2): 248.  
Syntype: FMNH 52389 (CM 645). New Guinea:  
Lake Sentani.  
APOGONIDAE
- Apogonichthys melampodus** Blosser, 1909  
Ann. Carnegie Mus., 6(1): 296.  
Holotype: FMNH 53049 (CM 1474). Lesser An-  
tilles: Saint Croix.  
APOGONIDAE
- Apuredoras rivasi** Fernández-Yépez, 1950  
Mem. Soc. Cienc. Nat. La Salle, 10(27): 195.  
Paratypes: 7. Venezuela: Río Apure near San  
Fernando.  
DORADIDAE

- Araiophos gracilis** Grey, 1961  
Pac. Sci., 15(3): 465.  
Holotype: FMNH 73901. Hawaiian Islands: off Mauna Loa lava flow.  
Paratypes: 8. Same locality.  
STERNOPTYCHIDAE
- Archicheir minutus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 46.  
Holotype: FMNH 52786 (CM 1186). Guyana: Christianburg.  
LEBIASINIDAE
- Archosargus oviceps** Ginsburg, 1952  
J. Wash. Acad. Sci., 42(3): 94.  
Paratypes: 2. Texas: Harbor Island.  
SPARIDAE
- Areliscus hollandi** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 62.  
Holotype: FMNH 55627 (CM 4369). Korea: Fusan.  
CYNOGLOSSIDAE
- Argentina alicae** Cohen and Atsides, 1969  
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 19.  
Paratypes: 5. Pacific Ocean: off Peru.  
ARGENTINIDAE
- Argentina brucei** Cohen and Atsides, 1969  
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 31.  
Paratypes: 20. Caribbean Sea: 7 localities.  
ARGENTINIDAE
- Argentina georgei** Cohen and Atsides, 1969  
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 27.  
Paratypes: 3. Bahama Banks and Antilles: 3 localities.  
ARGENTINIDAE
- Argentina stewarti** Cohen and Atsides, 1969  
U.S. Fish Wildl. Serv. Fish. Bull., 68(1): 29.  
Paratypes: 4. Antilles: 3 localities.  
ARGENTINIDAE
- Arius kanganamanensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 387.  
Holotype: FMNH 17194. New Guinea: Sepik River at Kanganaman.  
ARIIDAE
- Arius nox** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 388.  
Holotype: FMNH 17195. New Guinea: Sepik River at Nyaurangai.  
Paratypes: 5. New Guinea: 2 localities.  
ARIIDAE
- Arius solidus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 385.  
Holotype: FMNH 17201. New Guinea: Sepik River at Timbunke.  
Paratypes: 9. New Guinea: Sepik River, 3 localities.  
ARIIDAE
- Arius tuyra** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 128.  
Paratypes: 5. Panama: 2 localities.  
ARIIDAE
- Asiphonichthys hemigrammus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 403.  
Holotype: FMNH 53665 (CM 2137). Guyana: Gluck Island.  
CHARACIDAE
- Aspidoras maculosus** Nijssen and Isbrucker, 1976  
Bijdr. Dierkd., 46(1): 119.  
Holotype: FMNH 54810 (CM 3457). Brazil: Rio Paiaia.  
Paratypes: 6. Brazil: 3 localities.  
CALLICHTHYIDAE
- Astroblepus cyclopus santanderensis** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 675.  
Lectotype: FMNH 58433 (CM 7430). Colombia: Quebrada de Guapota.  
Lectoparatypes: 30. Colombia: 8 localities.  
Lectotype designated by Henn (1928, p. 82).  
ASTROBLEPIDAE
- Astroblepus frenatus** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 676.  
Holotype: FMNH 58384 (CM 7380). Colombia: Quebrada de San Joaquín.  
ASTROBLEPIDAE
- Astroblepus grixalvii micrescens** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 677.  
Holotype: FMNH 58376 (CM 7372). Colombia: Quebrada de Agua Larga.  
Paratypes: 41. Colombia: 9 localities.  
ASTROBLEPIDAE
- Astroblepus latidens** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 674.  
Holotype: FMNH 58366 (CM 7362). Colombia: Piperel.  
Paratypes: 40. Colombia: 5 localities.  
ASTROBLEPIDAE

- Astroblepus pholeter** Collette, 1962  
Proc. Biol. Soc. Wash., 75: 311.  
Paratype. Ecuador: Napo Province, cave in Latastas.  
ASTROBLEPIDAE
- Astroblepus rosei** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 55.  
Paratype. Peru: Río Jequetepeque at Llállán.  
ASTROBLEPIDAE
- Astronesthes ijimai** Tanaka, 1908  
J. Coll. Sci. Imp. Univ. Tokyo, 23(13): 9.  
Syntypes (?): 10, FMNH 89100 (CM 4610). Japan: Sagami Sea.  
These differ from material listed in original publication in being a series of 10 collected on 1 June 1907 (versus 5 from May).  
ASTRONESTHIDAE
- Astronotus orbiculatus** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 331.  
Holotype: FMNH 53992 (CM 2515). Brazil: Santarém.  
Paratype. Same locality.  
CICHLIDAE
- Astyanax abramoides** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 21.  
Holotype: FMNH 52863 (CM 1028). Guyana: Potaro River at Tumatumari.  
Paratypes: 78. Guyana: 8 localities.  
CHARACIDAE
- Astyanax aeneus costaricensis** Meek, 1914  
Field Mus. Nat. Hist. Zool. Ser., 10(10): 105.  
Syntypes: 203, FMNH 6849 and 7866-7873.  
Costa Rica: 8 localities.  
CHARACIDAE
- Astyanax albeolus** Eigenmann, 1908  
Bull. Mus. Comp. Zool. Harv. Coll., 52(6): 97.  
Holotype: FMNH 6241. Costa Rica: Río Machuca.  
CHARACIDAE
- Astyanax aurocaudatus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 26.  
Holotype: FMNH 56882 (CM 5162). Colombia: Boquia.  
Paratypes: 3. Same locality.  
CHARACIDAE
- Astyanax bimaculatus novae** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 175.  
Holotype (?): FMNH 54641 (CM 3278). Brazil: Rio Sapon at Prazeres.  
Paratypes: 23. Brazil: 2 localities.  
One of the 11 specimens in the paratype lot FMNH 54641 could be the illustrated holotype.  
CHARACIDAE
- Astyanax bimaculatus paraguayensis** Eigenmann, 1921  
Mem. Mus. Comp. Zool. Harv. Coll., 43(3): 256.  
Syntypes: 36, FMNH 54643-54648 (CM 3280-3286). Bolivia, Brazil, and Paraguay: 7 localities.  
Two lots CM 3274 and 3287 with 6 specimens are MISSING.  
CHARACIDAE
- Astyanax daguae** Eigenmann, 1913  
Indiana Univ. Stud., 18: 23.  
Holotype: FMNH 56251 (CM 5052). Colombia: Córdova.  
CHARACIDAE
- Astyanax essequibensis** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 17.  
Holotype: FMNH 53519 (CM 1018). Guyana: Tumatumari.  
Paratypes: 80. Guyana: 5 localities.  
CHARACIDAE
- Astyanax grandis** Meek and Hildebrand, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(6): 67.  
Holotype: FMNH 7571. Panama: Río Juan Díaz.  
Paratypes: 196. Panama: 7 localities.  
CHARACIDAE
- Astyanax guaporensis** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 176.  
Holotype: FMNH 54709 (CM 3351). Brazil: Rio Guaporé at Maciel.  
Paratype. Same locality.  
CHARACIDAE
- Astyanax guianensis** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 16.  
Holotype: FMNH 53518 (CM 1013). Guyana: Warraputa.  
Paratypes: 45. Guyana: 3 localities.  
CHARACIDAE

***Astyanax gymnogynys*** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 179.  
Holotype: FMNH 54707 (CM 3350). Brazil: Rio Iguaçu at Porto União.  
Paratype. Same locality.  
CHARACIDAE

***Astyanax hasemani*** Eigenmann, 1914  
Indiana Univ. Stud., 19: 10.  
Holotype: FMNH 56659 (CM 5476). Brazil: Porto Alegre.  
Paratypes: 6. Same locality.  
CHARACIDAE

***Astyanax heterurus*** Eigenmann and Wilson, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 11.  
Holotype: FMNH 56577 (CM 5392). Colombia: Río Truando.  
CHARACIDAE

***Astyanax kompi*** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 260.  
Paratypes: 6. Panama: Volcán, Laguna Gulnar or Grande.  
CHARACIDAE

***Astyanax magdalenae*** Eigenmann and Henn, 1916  
Ann. Carnegie Mus., 10(1–2): 89.  
Holotype: FMNH 57006 (CM 5822). Colombia: Girardot.  
CHARACIDAE

***Astyanax marionae*** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 175.  
Holotype: FMNH 54711 (CM 3353). Brazil: near São Luiz de Cáceres.  
Paratypes: 8. Brazil: 2 localities.  
CHARACIDAE

***Astyanax metae*** Eigenmann, 1914  
Indiana Univ. Stud., 19: 11.  
Holotype: FMNH 56640 (CM 5457). Colombia: Río Negro at Villavicencio.  
Paratypes: 2. Same locality.  
CHARACIDAE

***Astyanax microlepis*** Eigenmann, 1913  
Indiana Univ. Stud., 18: 24.  
Holotype: FMNH 56209 (CM 5001). Colombia: Piedra de Moler.  
Paratypes: 843. Colombia: 5 localities.  
CHARACIDAE

***Astyanax mucronatus*** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 19.  
Holotype: FMNH 53520 (CM 1025). Guyana: Potaro River at Tumatumari.  
Paratypes: 7. Guyana: 2 localities.  
CHARACIDAE

***Astyanax mutator*** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 18.  
Holotype: FMNH 52688 (CM 1023). Guyana: Potaro River at Savannah Landing.  
Paratypes: 61. Guyana: 2 localities.  
CHARACIDAE

***Astyanax nasutus*** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 108.  
Holotype: FMNH 5909. Nicaragua: Lago Managua at Managua.  
Paratypes: 15. Nicaragua: 2 localities.  
CHARACIDAE

***Astyanax notemigonoides*** Fowler, 1911  
Proc. Acad. Nat. Sci. Phila., 63: 506.  
Paratype. Ecuador: near Bucay.  
CHARACIDAE

***Astyanax paranahybae*** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 177.  
Holotype: FMNH 54714 (CM 3356). Brazil: Rio Paranahyba.  
CHARACIDAE

***Astyanax potaroensis*** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 22.  
Holotype: FMNH 52695 (CM 1037). Guyana: Potaro River at Amatuk Cataract.  
Paratypes: 7. Guyana: 3 localities.  
CHARACIDAE

***Astyanax regani*** Meek, 1909  
Field Mus. Nat. Hist. Zool. Ser., 7(7): 207.  
Holotype: FMNH 6257. Costa Rica: Las Cañas.  
CHARACIDAE

***Astyanax ribeirae*** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 177.  
Holotype: FMNH 54725 (CM 3368). Brazil: Xiririca.  
Paratypes: 103. Brazil: 4 localities.  
CHARACIDAE

- Astyanax robustus** Meek, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(7): 69.  
Holotype: FMNH 7682. Costa Rica: Virginia River.  
Paratypes (?): 116. Costa Rica.  
See comment in Grey (1947, pp. 122–123); status of various paratypes uncertain.  
CHARACIDAE
- Astyanax ruberrimus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 25.  
Holotype: FMNH 56122 (CM 4912). Colombia: Istmina.  
Paratypes: 315. Colombia: 4 localities.  
CHARACIDAE
- Astyanax scabripinnis paranae** Eigenmann, 1914  
Indiana Univ. Stud., 20: 47.  
Paratypes: 307. Brazil: 10 localities.  
CHARACIDAE
- Astyanax scierus** Fowler, 1911  
Proc. Acad. Nat. Sci. Phila., 63: 509.  
Paratype: Ecuador: near Bucay.  
CHARACIDAE
- Atherina evermanni** Eigenmann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 228.  
Paratypes: 5. Cuba: 2 localities.  
ATHERINIDAE
- Atherina insularum** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 170.  
Paratype. Hawaiian Islands: Honolulu.  
ATHERINIDAE
- Atherina ovalaua** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 401.  
Holotype: FMNH 17272. Fiji Islands: Ovalau Island.  
Paratypes: 33. Fiji Islands: 2 localities.  
ATHERINIDAE
- Atherinichthys nouhuysi** Weber, 1910  
Notes Leiden Mus., 32: 229.  
Syntypes: 2, FMNH 52500 (CM 758). New Guinea: Lorentz River, Alkmaar.  
ATHERINIDAE
- Atherinops affinis littoralis** Hubbs, 1918  
Bull. Am. Mus. Nat. Hist., 38(13): 426.  
Holotype: FMNH 2064. California: San Diego Bay, North Island.  
Paratypes: 5. California: 3 localities.  
ATHERINIDAE
- Atherinops insularum cedroscensis** Hubbs, 1918  
Bull. Am. Mus. Nat. Hist., 38(13): 434.  
Paratypes: 2. Mexico: Baja California, Islas Cedros.  
ATHERINIDAE
- Atherinops insularum guadalupae** Hubbs, 1918  
Bull. Am. Mus. Nat. Hist., 38(13): 435.  
Paratypes: 2. Mexico: Baja California, Isla Guadalupe.  
ATHERINIDAE
- Atherion elymus aphrozoicus** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 295.  
Paratype. Philippine Islands: Tara Island.  
ATHERINIDAE
- Atherion elymus freyi** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 292.  
Paratypes: 2. Guam: 2 localities.  
ATHERINIDAE
- Atuona tricuspidata** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 429.  
Holotype: FMNH 17356. Marquesas Islands: Hiva Oa Island, Autona.  
Paratypes: 4. Same locality.  
GOBIIDAE
- Auchenipterus brevior** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 202.  
Holotype: FMNH 53249 (CM 1715). Guyana: Tumatumari.  
Paratypes: 6. Same locality.  
AUCHENIPTERIDAE
- Auchenipterus demerarae** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 202.  
Holotype: FMNH 53248 (CM 1714). Guyana: Wismar.  
Paratype. Guyana: Malali.  
AUCHENIPTERIDAE
- Awaous decemlineatus** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 686.  
Holotype: FMNH 58476 (CM 7478). Colombia: Quibdo.  
Paratypes: 9. Colombia: 2 localities.  
GOBIIDAE
- Aztecula mexicana** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 81.  
Holotype: FMNH 3606. Mexico: Río Moctezuma at San Juan del Río.  
Paratypes: 220. Same locality.  
CYPRINIDAE



**Barbatula toni posteroventralis** Nichols, 1925

Am. Mus. Nov., 171: 4.

Paratypes: 3. China: Shansi, Kwei-hwan.

COBITIDAE

**Barbatula yarkandensis sellaefer** Nichols, 1925

Am. Mus. Nov., 171: 4.

Paratypes: 3. China: Shansi, Chin-ssu.

COBITIDAE

**Barbuligobius boehlkei** Lachner and McKinney, 1974

Copeia, 1974(4): 871.

Paratype. Sri Lanka: Galle.

GOBIIDAE

**Barbus akeleyi** Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 13.

Holotype: FMNH 6106. Kenya: Athi River.

Paratypes: 12. Same locality.

CYPRINIDAE

**Barbus anoplus** Weber, 1897

Zool. Jahrb., 10: 151.

Syntypes: 3, FMNH 52397 (CM 653). South

Africa: Buffels River near Laingsburg.

CYPRINIDAE

**Barbus athi** Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 10.

Holotype: FMNH 6108. Kenya: Athi River.

Paratypes: 63. Same locality.

CYPRINIDAE

**Barbus bantolanensis** Day, 1914

Philipp. J. Sci., 9(2): 188.

Paratypes: 12. Philippine Islands: Palawan Island, Lake Manguao.

CYPRINIDAE

**Barbus bifrenatus** Fowler, 1935

Ann. Transvaal Mus., 16(2): 266.

Paratypes: 10. Botswana: Tsotsoroga Pan.

CYPRINIDAE

**Barbus carens** Boulenger, 1912

Ann. Mus. Congo Belge Ser. Quarto Zool., Ser. 1, 2(3): 15.

Syntype: FMNH 56148 (CM 4939). Zaïre: Le-buzi River at Kuka Muno.

CYPRINIDAE

**Barbus fitzsimonsi** Fowler, 1935

Ann. Transvaal Mus., 16(2): 267.

Paratypes: 15. Botswana: 2 localities.

CYPRINIDAE

**Barbus helleri** Hubbs, 1918

Field Mus. Nat. Hist. Zool. Ser., 12(2): 12.

Holotype: FMNH 6105. Kenya: Athi River.

Paratypes: 103. Same locality.

CYPRINIDAE

**Barbus oatesii** Boulenger, 1893

Ann. Mag. Nat. Hist., Ser. 6, 12: 201.

Syntypes: 2, FMNH 5708. Burma: Nampandet.

CYPRINIDAE

**Barbus tsotsorogensi** Fowler, 1935

Ann. Transvaal Mus., 16(2): 265.

Paratypes: 23. Botswana: 2 localities.

CYPRINIDAE

**Bathylagus tetranema** Bailey and Stewart, 1984

Misc. Publ. Mus. Zool. Univ. Mich., 168:11.

Paratypes: 2. Zambia: Lake Tanganyika near Mpulungu.

BAGRIDAE

**Bathylupea schroederi** Dick, 1962

Breviora, 167: 1.

Paratypes: 4. Western Caribbean Sea: 3 localities.

BATHYCLUPEIDAE

**Bathylagus greyae** Cohen, 1958

Breviora, 98: 5.

Holotype: FMNH 49730. Western North Atlantic: near Bermuda.

BATHYLAGIDAE

**Bathypterois bigelowi** Mead, 1958

J. Wash. Acad. Sci., 48(11): 366.

Holotype: FMNH 64435. Caribbean Sea: off Nicaragua.

Paratypes: 42. Caribbean Sea: 12 localities.

CHLOROPHTHALMIDAE

**Bathypterois dubius** Vaillant, 1888

Exp. Sci. Trav. et Talis. . . Poissons, Paris, p. 124 (not seen).

Paratype. Off Mauritania: Banc d'Arguin.

CHLOROPHTHALMIDAE

**Bathypterois pectoralis** Garman, 1899

Mem. Mus. Comp. Zool. Harv. Coll., 24: 257.

Syntype: FMNH 33730. Pacific Ocean: off Colombia.

CHLOROPHTHALMIDAE

**Bathytyphlops marionae** Mead, 1958  
J. Wash. Acad. Sci., 48(11): 370.  
Holotype: FMNH 64439. Caribbean Sea.  
CHLOROPHTHALMIDAE

**Batrachoides waltersi** Collette and Russo, 1981  
Bull. Mar. Sci., 31(2): 208.  
Paratypes: 3. El Salvador: Bahía Jiquilisco, La Venadona.  
BATRACHOIDIDAE

**Beirabarus palustris** Herre, 1936  
Proc. Biol. Soc. Wash., 49: 100.  
Paratypes: 12. Mozambique: near Beira.  
CYPRINIDAE

**Bellator ribeiroi** Miller, 1965  
Q. J. Fla. Acad. Sci., 28(3): 259.  
Paratypes: 2. Western Atlantic Ocean: off Colombia.  
TRIGLIDAE

**Belligobio eristigma** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 173.  
Holotype: FMNH 58714 (CM 7820). Japan: near Okayama.  
CYPRINIDAE

**Belone svetovidovi** Knudsen and Wolff, 1970  
Atl. Rep., 11: 21.  
Paratypes: 7. Tunisia: Gabes.  
BELONIDAE

**Bembrops greyi** Poll, 1959  
Résult. Sci. Exp. Oceanogr. Belge . . . (1948–1949), 4(3B): 40.  
Paratype. South Atlantic: off Angola, Ambri-zete.  
PERCOPHIDAE

**Bivibranchia protractila** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 259.  
Holotype: FMNH 53576 (CM 1873). Guyana: Bartica sand-bank.  
Paratypes: 8. Guyana: 3 localities.  
HEMIODONTIDAE

**Blenniulus proteus** Krejsa, 1960  
Copeia, 1960(4): 326.  
Paratypes: 2. Mexico: Islas Revilla Gigedo southeast of Isla San Benedictino.  
BLENNIIDAE

**Blennius nicholsi** Tavalga, 1954  
Copeia, 1954(2): 135.  
Paratypes: 2. Florida: 2 localities.  
BLENNIIDAE

**Bleptonema amazonae** Eigenmann, 1914  
Indiana Univ. Stud., 20: 44.  
Holotype: FMNH 56679 (CM 5497). Brazil: Santarém.  
Paratypes: 2. Same locality.  
CHARACIDAE

**Bleptonema paraguayensis** Eigenmann, 1914  
Indiana Univ. Stud., 20: 44.  
Holotype: FMNH 56681 (CM 5499). Paraguay: Corumbá.  
Paratypes: 4. Same locality.  
CHARACIDAE

**Bodianus stellatus** Blosser, 1909  
Ann. Carnegie Mus., 6(1): 297.  
Holotype: FMNH 53048 (CM 1473). Lesser Antilles: Saint Croix Island.  
SERRANIDAE

**Bombonia uxorius** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 395.  
Holotype: FMNH 17493. Indonesia: Waigiu Island.  
SYNGNATHIDAE

**Boroda malua** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 414.  
Holotype: FMNH 17349. New Guinea: Sepik River at Malu.  
Paratypes: 2. Same locality.  
ELEOTRIDAE

**Bostrychus zonatus** Weber, 1908  
Nova Guinea, 5(2): 259.  
Syntype: FMNH 52457 (CM 715). New Guinea: Merauke River.  
ELEOTRIDAE

**Botia purpurea** Nichols, 1925  
Am. Mus. Nov., 177: 4.  
Paratype. China: Hunan, Tungting Lake.  
COBITIDAE

**Brachirus bellus** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 274.  
Holotype: FMNH 58783 (CM 7894). Japan: Misaki.  
SCORPAENIDAE

- Brachychalcinus guianensis** Boeseman, 1952  
Zool. Meded. (Leiden), 31(26): 303.  
Syntypes: 6, FMNH 51535 and 59525. Guyana:  
New River drainage, Itabu Creek.  
CHARACIDAE
- Brachyglanis frenata** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 156.  
Holotype: (CM 1670) MISSING. Guyana:  
Amatuk.  
PIMELODIDAE
- Brachyglanis melas** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 157.  
Holotype: FMNH 53217 (CM 1672). Guyana:  
Crab Falls.  
Paratypes: 4. Guyana: 3 localities.  
PIMELODIDAE
- Brachyglanis phalacra** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 157.  
Holotype: FMNH 53216 (CM 1671). Guyana:  
Amatuk.  
PIMELODIDAE
- Brachygobius kabiliensis** Inger, 1958  
Fieldiana Zool., 39(14): 110.  
Holotype: FMNH 47991. Malaysia: Sabah, Ka-  
bili River.  
Paratypes: 6. Malaysia: Sabah, 2 localities.  
GOBIIDAE
- Brachygobius sabanus** Inger, 1958  
Fieldiana Zool., 39(14): 113.  
Holotype: FMNH 47990. Malaysia: Sabah, La-  
mag.  
Paratypes: 21. Malaysia: Sabah, 3 localities.  
GOBIIDAE
- Brachygobius xanthomelas** Herre, 1937  
Bull. Raffles Mus., 13: 43.  
Paratype. Malaysia: Johore, Mawai district.  
GOBIIDAE
- Bramocharax elongatus** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 110.  
Holotype: FMNH 5922. Nicaragua: Lago Ma-  
nagua at Managua.  
Paratypes: 26. Nicaragua: 3 localities.  
See comment in Grey (1947, p. 126); status  
of various paratypes uncertain.  
CHARACIDAE
- Branchiostoma bermudae** Hubbs, 1922  
Occas. Pap. Mus. Zool. Univ. Mich., 105: 9.  
Paratypes: 71. Bermuda: The Flatts.  
BRANCHIOSTOMATIDAE
- Branchiostoma floridae** Hubbs, 1922  
Occas. Pap. Mus. Zool. Univ. Mich., 105: 7.  
Paratypes: 7. Florida: Tampa Bay.  
BRANCHIOSTOMATIDAE
- Bryconamericus alfredeae** Eigenmann, 1927  
Mem. Mus. Comp. Zool. Harv. Coll., 43(4):  
394.  
Paratypes: 5. Peru: Río Urubamba at Santa Ana.  
CHARACIDAE
- Bryconamericus alpha** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana  
Univ. Stud., 19: 7.  
Holotype: FMNH 56646 (CM 5463). Colombia:  
Villavicencio.  
Paratypes: 2. Same locality.  
CHARACIDAE
- Bryconamericus beta** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana  
Univ. Stud., 19: 7.  
Holotype: FMNH 56648 (CM 5465). Colombia:  
Villavicencio.  
Paratypes: 4. Same locality.  
CHARACIDAE
- Bryconamericus cascajalensis** Meek and Hilde-  
brand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 284.  
Holotype: FMNH 8946. Panama: Río Cascajal  
at Porto Bello.  
Paratypes: 24. Same locality.  
CHARACIDAE
- Bryconamericus caucanus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 17.  
Holotype: FMNH 56229 (CM 5031). Colombia:  
Piedra de Moler.  
Paratypes: 106. Colombia: 6 localities.  
CHARACIDAE
- Bryconamericus chocoensis** Eigenmann, 1913  
Indiana Univ. Stud., 18: 14.  
Holotype: FMNH 56235 (CM 5036). Colombia:  
Istmina.  
Paratypes: 270. Colombia: 6 localities.  
CHARACIDAE

**Bryconamericus cismontanus** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 4.  
Holotype: FMNH 56642 (CM 5459). Colombia: Villavicencio.  
CHARACIDAE

**Bryconamericus conventus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 13.  
Holotype: FMNH 56261 (CM 5060). Colombia: Soplaviento.  
Paratypes: 57. Same locality.  
CHARACIDAE

**Bryconamericus decurrens** Eigenmann, 1913  
Indiana Univ. Stud., 18: 20.  
Holotype: FMNH 56255 (CM 5055). Colombia: Soplaviento.  
CHARACIDAE

**Bryconamericus dentatus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 19.  
Holotype: FMNH 56253 (CM 5054). Colombia: Piedra de Moler.  
Paratypes: 21. Colombia: 3 localities.  
CHARACIDAE

**Bryconamericus deuterodonoides** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 5.  
Holotype: FMNH 56644 (CM 5461). Colombia: Río Negro at Villavicencio.  
Paratypes: 6. Same locality.  
CHARACIDAE

**Bryconamericus deuterodonoides caudovittatus** Inger, 1956  
Fieldiana Zool., 34(37): 435.  
Holotype: FMNH 45706. Venezuela: Río Abácapa, west side of Chimantá-tepui.  
Paratypes: 12. Same locality.  
CHARACIDAE

**Bryconamericus diquensis** Eigenmann, 1913  
Indiana Univ. Stud., 18: 14.  
Holotype: FMNH 56272 (CM 5072). Colombia: Soplaviento.  
Paratypes: 11. Same locality.  
CHARACIDAE

**Bryconamericus grosvenori** Eigenmann, 1927  
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 365.  
Paratypes: 3. Peru: 2 localities.  
CHARACIDAE

**Bryconamericus henni** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 6.  
Holotype: FMNH 56594 (CM 5410). Ecuador: Vinces.  
Paratypes: 10. Ecuador: 2 localities.  
CHARACIDAE

**Bryconamericus hyphesson** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 32.  
Holotype: FMNH 52708 (CM 1070). Guyana: Tumatumari.  
Paratypes: 3. Same locality.  
CHARACIDAE

**Bryconamericus magdalenensis** Eigenmann, 1913  
Indiana Univ. Stud., 18: 14.  
Holotype: FMNH 56263 (CM 5063). Colombia: Girardot.  
Paratypes: 675. Colombia: 8 localities.  
CHARACIDAE

**Bryconamericus novae** Eigenmann and Henn, 1914  
Indiana Univ. Stud., 24: 234.  
Holotype: FMNH 54905 (CM 3568). Brazil: Rio Novo near Piabana.  
Paratypes: 9. Same locality.  
CHARACIDAE

**Bryconamericus ortholepis** Eigenmann, 1913  
Indiana Univ. Stud., 18: 15.  
Holotype: (CM 5088) MISSING. Colombia: Boca de Raspadura.  
CHARACIDAE

**Bryconamericus osgoodi** Eigenmann and Allen, 1942  
Fishes of Western South America, Lexington, Kentucky, p. 225.  
Paratypes: 3. Peru: Moyobamba.  
CHARACIDAE

**Bryconamericus pachacuti** Eigenmann, 1927  
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 376.  
Paratypes: 12. Peru: 2 localities.  
CHARACIDAE

**Bryconamericus peruanus ricae** Eigenmann, 1908  
Bull. Mus. Comp. Zool. Harv. Coll., 52(6): 106.  
Lectotype: FMNH 59523. Costa Rica: Chitaria.  
Lectoparatypes: 2. Same locality.  
Lectotype designated by Grey (1947, pp. 127–128).  
CHARACIDAE

**Bryconamericus scopiferus** Eigenmann, 1913

Indiana Univ. Stud., 18: 16.

Holotype: FMNH 56224 (CM 5026). Colombia:  
Río San Juan at Itsmina.

Paratypes: 122. Colombia: 3 localities.

CHARACIDAE

**Bryconamericus scopiferus guaytarae** Eigenmann  
and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana  
Univ. Stud., 19: 7.

Holotype: FMNH 56657 (CM 5474). Colombia:  
Río Patia at mouth of Río Guáytara.

Paratypes: 122. Colombia: 3 localities.

CHARACIDAE

**Bryconamericus tolimae** Eigenmann, 1913

Indiana Univ. Stud., 18: 18.

Holotype: FMNH 56257 (CM 5057). Colombia:  
Ibagué.

Paratypes: 7. Same locality.

CHARACIDAE

**Bryconamericus zeteki** Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 265.

Paratypes: 8. Panama: Pacific slope, creek in El  
Valle.

CHARACIDAE

**Brycon argenteus** Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.

Holotype: FMNH 7588. Panama: Río Aruza at  
Aruza.

Paratypes: 168. Panama: 11 localities.

CHARACIDAE

**Brycon behreae** Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 278.

Paratype. Panama: Río Chiriquí del Tire above  
Caldera.

CHARACIDAE

**Brycon henni** Eigenmann, 1913

Indiana Univ. Stud., 18: 26.

Holotype: FMNH 56384 (CM 5152). Colombia:  
Caldas.

Paratypes: 102. Colombia: 6 localities.

CHARACIDAE

**Brycon obscurus** Hildebrand, 1938

Field Mus. Nat. Hist. Zool. Ser., 22(4): 283.

Paratypes: 2. Panama: Pacific slope, creek in El  
Valle.

CHARACIDAE

**Brycon petrosus** Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.

Holotype: FMNH 7589. Panama: Upper  
Chagres.

Paratypes: 89. Panama: 10 localities.

CHARACIDAE

**Brycon siebenthalae** Eigenmann, 1912

Mem. Carnegie Mus., 5: 372.

Holotype: FMNH 53353 (CM 1819). Guyana:  
Aruka River, Mud Creek.

CHARACIDAE

**Bunocephalus amaurus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 126.

Holotype: FMNH 53121 (CM 1555). Guyana:  
Konawaruk.

ASPREDINIDAE

**Bunocephalus chamaizelus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 127.

Holotype: FMNH 53122 (CM 1556). Guyana:  
Erukin.

Paratypes: 7. Guyana: 3 localities.

ASPREDINIDAE

**Bunocephalus colombianus** Eigenmann, 1912

Indiana Univ. Stud., 16: 10.

Holotype: FMNH 56038 (CM 4828). Colombia:  
Raspadura.

ASPREDINIDAE

**Bunocephalus depressus** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 319.

Holotype: FMNH 54338 (CM 2984). Bolivia:  
Río Machupo near San Joaquín.

ASPREDINIDAE

**Calamopteryx jeb** Cohen, 1973

Proc. Biol. Soc. Wash., 86(28): 343.

Holotype: FMNH 71603. Ecuador: Islas Galá-  
pagos, Tagus Cove.

Paratypes: 24. Same locality.

OPHIDIIDAE

**Callechelys bilinearis** Kanazawa, 1952

Fieldiana Zool., 34(7): 72.

Holotype: FMNH 48973. Bermuda: Cooper's  
Island and Saint David's.

OPHICHTHIDAE

**Callionymus distethommatatus** Fowler, 1941

Proc. U.S. Natl. Mus., 90(3106): 18.

Paratypes: 2. Philippine Islands: Port Matalvi  
off western Luzon.

CALLIONYMIDAE

- Callionymus hudsoni** Fowler, 1941  
Proc. U.S. Natl. Mus., 90(3106): 8.  
Paratype. Philippine Islands: Mantacao Island.  
CALLIONYMIDAE
- Callionymus tenuis** Fricke, 1981  
Ann. Mus. Civ. Stor. Nat. "Giacomo Doria"  
Genova, 83: 387.  
Holotype: FMNH 78916. Maldives Archipelago: South Nilandu Atoll.  
CALLIONYMIDAE
- Callionymus wilburi** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 431.  
Holotype: FMNH 17491. Indonesia: Waigiu Island.  
Paratype. Same locality.  
CALLIONYMIDAE
- Calliurichthys astrinius** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 80.  
Holotype: FMNH 55207 (CM 3903). Hawaiian Islands: Honolulu Market.  
CALLIONYMIDAE
- Calliurichthys zanectes** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 81.  
Holotype: FMNH 55208 (CM 3904). Hawaiian Islands: Honolulu Market.  
CALLIONYMIDAE
- Callogobius ocellatus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 422.  
Holotype: FMNH 17363. Fiji Islands: Ovalau Island.  
GOBIIDAE
- Calymmichthys xenicus** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6: 296.  
Holotype: FMNH 57089 (CM 6027). Japan: Sagami Bay.  
CALLIONYMIDAE
- Caranx formosanus** Jordan and Snyder, 1909  
Mem. Carnegie Mus., 4(2): 38.  
Holotype: FMNH 55364 (CM 412). Taiwan: Takao.  
CARANGIDAE
- Caranx ishikawai** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 193.  
Holotype: FMNH 59390 (CM 7734). Japan: Bonin Islands.  
CARANGIDAE
- Caranx miyakamii** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 201.  
Holotype: FMNH 59407 (CM 7742). Taiwan.  
CARANGIDAE
- Caranx oshimai** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 189.  
Holotype: FMNH 59492 (CM 7731). Taiwan.  
CARANGIDAE
- Caranx rastrosus** Jordan and Snyder, 1909  
Mem. Carnegie Mus., 4(2): 37.  
Holotype: FMNH 55363 (CM 411). Taiwan: Takao.  
CARANGIDAE
- Caranx tanakai** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 177.  
Paratype. Japan: Kii.  
CARANGIDAE
- Caranx uii** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 174.  
Holotype: FMNH 59494 (CM 7720). Japan: Kii.  
Paratype. Same locality.  
CARANGIDAE
- Careproctus burkei** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 281.  
Holotype: FMNH 57515 (CM 6457). Japan: Yokohama Market.  
CYCLOPTERIDAE
- Careproctus gilberti** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 282.  
Holotype: FMNH 57514 (CM 6456). Japan: Sagami Bay, Misaki.  
CYCLOPTERIDAE
- Carpiodes elongatus** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 28.  
Holotype: FMNH 4425. Mexico: Nuevo León, Linares.  
Paratypes: 13. Mexico: 3 localities.  
CATOSTOMIDAE
- Carpiodes labiosus** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 29.  
Holotype: FMNH 4492. Mexico: San Luis Potosí, Valles.  
Paratype. Same locality.  
CATOSTOMIDAE

- Carpiodes microstomus** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 27.  
Holotype: FMNH 3542. Mexico: Río Noavaco at Santa Rosalia.  
Paratypes: 18. Mexico: 2 localities.  
CATOSTOMIDAE
- Cascadura maculocephala** Ellis, 1913  
Ann. Carnegie Mus., 8(3–4): 387.  
Holotype: FMNH 54878 (CM 3539). Brazil: Rio Uruguay at Uruguayana.  
CALLICHTHYIDAE
- Catostomus conchos** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 75.  
Holotype: FMNH 3557. Mexico: Río Conchos at Jiménez.  
Paratypes: 8. Same locality.  
CATOSTOMIDAE
- Catostomus sonorensis** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 75.  
Holotype: FMNH 3564. Mexico: Río Paphigochic at Miñaca.  
Paratypes: 21. Same locality.  
CATOSTOMIDAE
- Centriscus capito** Oshima, 1922  
Ann. Carnegie Mus., 13(3–4): 263.  
Holotype: FMNH 59149 (CM 8287). Taiwan: Toko.  
CENTRISCIDAE
- Centropristes melanus** Ginsburg, 1952  
J. Wash. Acad. Sci., 42(3): 84.  
Holotype: FMNH 33719. Florida: Newport near Wakulla.  
Paratypes: 6. Same locality.  
SERRANIDAE
- Centropyge argi** Woods and Kanazawa, 1951  
Fieldiana Zool., 31(53): 636.  
Holotype: FMNH 48494. Bermuda: Argus Bank.  
POMACANTHIDAE
- Centropyge heraldi** Woods and Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 602.  
Paratype. Marshall Islands: Bikini Atoll, Boby Island.  
POMACANTHIDAE
- Centropyge tutuilae** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 62.  
Holotype: FMNH 55206 (CM 3902). Samoan Islands: Tetuila, Pago Pago.  
POMACANTHIDAE
- Cephalopholis boninius** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 248.  
Holotype: FMNH 57099 (CM 6038). Japan: Bonin Islands.  
SERRANIDAE
- Cephalosilurus fowleri** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 317.  
Holotype: FMNH 54254 (CM 2880). Brazil: Bahia, Cidade de Barra.  
PIMELODIDAE
- Ceratobranchia binghami** Eigenmann, 1927  
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 357.  
Paratypes: 5. Peru: 2 localities.  
CHARACIDAE
- Cetopsorhamdia boquillae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 37.  
Holotype: FMNH 55212 (CM 3923). Colombia: Boquilla.  
Paratypes: 5. Same locality.  
PIMELODIDAE
- Cetopsorhamdia nasus** Eigenmann and Fisher, 1916  
In: Eigenmann, Ann. Carnegie Mus., 10(1–2): 83.  
Holotype: FMNH 58126 (CM 7124). Colombia: Honda.  
PIMELODIDAE
- Cetopsorhamdia phantasia** Stewart, 1985  
Copeia 1985(2): 340.  
Holotype: FMNH 94601. Ecuador: Río Jivino, west of Limoncocha.  
Paratype. Same locality.  
PIMELODIDAE
- Chaenopsis alepidota californiensis** Böhlke, 1957  
Proc. Acad. Nat. Sci. Phila., 109: 99.  
Paratypes: 4. California: Santa Catalina Island, Avalon.  
CHAENOPSIDAE
- Chaenopsis roseola** Hastings and Shipp, 1980  
Proc. Biol. Soc. Wash., 93(4): 876.  
Paratype. Florida: southwest of Panama City Beach.  
CHAENOPSIDAE

**Chaenothorax eigenmanni** Ellis, 1913  
Ann. Carnegie Mus., 8(3-4): 393.  
Holotype: FMNH 54880 (CM 3542). Brazil: São  
Luiz de Cáceres.  
Paratype. Same locality.  
CALLICHTHYIDAE

**Chaetostomus leucomelas** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 681.  
Paratype. Colombia: Río Patía.  
LORICARIIDAE

**Characidium blennioides** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 37.  
Holotype: FMNH 71043 (CM 1144). Guyana:  
Erukin.  
Paratypes: 19. Guyana: 6 localities.  
CHARACIDAE

**Characidium catenatum** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 40.  
Holotype: FMNH 52761 (CM 1153). Guyana:  
Warraputa.  
Paratypes: 6. Guyana: 2 localities.  
CHARACIDAE

**Characidium caucanum** Eigenmann, 1912  
Indiana Univ. Stud., 16: 25.  
Holotype: FMNH 56057 (CM 4847). Colombia:  
Cali.  
Paratypes: 20. Colombia: 3 localities.  
CHARACIDAE

**Characidium laterale** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 36.  
Holotype: FMNH 72627 (CM 1141). Guyana:  
Amatuk.  
CHARACIDAE

**Characidium pellucidum** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 39.  
Holotype: FMNH 52764 (CM 1156). Guyana:  
Gluck Island.  
CHARACIDAE

**Characidium phoxocephalum** Eigenmann, 1912  
Indiana Univ. Stud., 16: 26.  
Holotype: FMNH 56061 (CM 4851). Colombia:  
Paila.  
CHARACIDAE

**Characidium pteroides** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 40.  
Holotype: FMNH 53548 (CM 1157). Guyana:  
Konawaruk.  
Paratype. Guyana: Rockstone.  
CHARACIDAE

**Characidium vintoni** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 36.  
Holotype: FMNH 52756 (CM 1142). Guyana:  
Shrimp Creek.  
Paratypes: 22. Same locality.  
CHARACIDAE

**Characidium zebra** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 38.  
Holotype: FMNH 53547 (CM 1151). Guyana:  
Ireng River, Maripicru Creek.  
Paratypes: 7. Same locality.  
CHARACIDAE

**Characodon multiradiatus** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 119.  
Holotype: FMNH 4523. Mexico: Río Lerma at  
Lerma.  
Paratypes: 23. Same locality.  
GOODEIDAE

**Charax metae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 238.  
Paratypes: 10. Colombia: Quebrada Cramalote  
at Villavicencio.  
CHARACIDAE

**Charax rupununi** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 402.  
Holotype: FMNH 53663 (CM 2135). Guyana:  
Rupununi.  
Paratype. Same locality.  
CHARACIDAE

**Chasmias misakius** Jordan and Snyder, 1901  
Proc. U.S. Natl. Mus., 23(1235): 761.  
Paratypes: 35. Japan: Sagami Sea, Misaki.  
GOBIIDAE

**Chasmocranus brevior** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 162.  
Holotype: FMNH 53213 (CM 1662). Guyana:  
Warratuk.  
PIMELODIDAE

**Chasmocranus chimantanus** Inger, 1956  
Fieldiana Zool., 34(37): 425.  
Holotype: FMNH 45704. Venezuela: Río Abá-  
capa, west side of Chimantá-tepui.  
Paratypes: 3. Same locality.  
PIMELODIDAE



- Chasmocranus longior** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 160.  
 Holotype: FMNH 53208 (CM 1655). Guyana: Amatuk.  
 Paratypes: 13. Guyana: 5 localities.  
 PIMELODIDAE
- Chasmocranus rosae** Eigenmann, 1922  
 Mem. Carnegie Mus., 9(1): 220.  
 Holotype: FMNH 55140 (CM 3841). Colombia: Río Negro at Villavicencio.  
 Paratypes: 2. Same locality.  
 PIMELODIDAE
- Chasmodes bosquianus longimaxilla** Williams, 1983  
 Bull. Fla. State Mus. Biol. Sci., 29(2): 85.  
 Paratypes: 14. Gulf of Mexico: 4 localities.  
 BLENNIIDAE
- Cheirocerus eques** Eigenmann, 1917  
 Ann. Carnegie Mus., 11(3–4): 398.  
 Holotype: FMNH 58255 (CM 7254). Bolivia: Villa Bella.  
 Paratypes: 5. Bolivia: 3 localities.  
 PIMELODIDAE
- Cheirodon annae** McAtee, 1903  
 Proc. Acad. Nat. Sci. Phila., 59: 515.  
 Paratypes: 2. South America.  
 CHARACIDAE
- Cheirodon dialepturus** Fink and Weitzman, 1974  
 Smithson. Contrib. Zool., 172: 5.  
 Paratypes: 2. Panama: Río San Pedro basin, west of Santiago.  
 CHARACIDAE
- Cheirodon eigenmanni** Meek, 1912  
 Field Mus. Nat. Hist. Zool. Ser., 10(7): 70.  
 Holotype: FMNH 7683. Costa Rica: La Junta.  
 Paratype. Same locality.  
 CHARACIDAE
- Cheirodon galusdae** Eigenmann, 1927  
 Mem. Natl. Acad. Sci. Wash., 22(2): 44.  
 Paratypes (?): 3. Chile: Río Locomilla at San Javier.  
 These came to Field Museum via University of Illinois, Urbana, where they had been sent by Eigenmann for a study of fish parasites.  
 CHARACIDAE
- Cheirodon interruptus ibicuhiensis** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 74.  
 Syntypes: 8, FMNH 57833 (CM 6816). Brazil: Cacequy.  
 CHARACIDAE
- Cheirodon madeirae** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 76.  
 Holotype: FMNH 57864 (CM 6847). Bolivia: San Joaquín.  
 CHARACIDAE
- Cheirodon microdon** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 80.  
 Holotype: FMNH 57867 (CM 6850). Brazil: Cáceres.  
 Paratype. Same locality.  
 CHARACIDAE
- Cheirodon notomelas** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 74.  
 Holotype: FMNH 57829 (CM 6812). Brazil: Rio Tietê basin at Miguel Calmone.  
 Paratypes: 10. Same locality.  
 CHARACIDAE
- Cheirodon parahybae** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 70.  
 Holotype: FMNH 57857 (CM 6841). Brazil: Campos.  
 Paratypes: 5. Same locality.  
 CHARACIDAE
- Cheirodon stenodon** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 82.  
 Holotype: FMNH 57865 (CM 6848). Brazil: Bebedouro near Rio Grande and Rio Paraná.  
 Paratypes: 28. Same locality.  
 CHARACIDAE
- Chelidonichthys ischyryus** Jordan and Thompson, 1914  
 Mem. Carnegie Mus., 6(4): 282.  
 Holotype: FMNH 57517 (CM 6459). Japan: Sagami Bay.  
 TRIGLIDAE
- Chirostoma attenuatum** Meek, 1902  
 Field Mus. Nat. Hist. Zool. Ser., 3(16): 112.  
 Holotype: FMNH 3631. Mexico: Lago de Patzcuaro at Patzcuaro.  
 Paratypes: 12. Same locality.  
 ATHERINIDAE

- Chirostoma consocium** Jordan and Hubbs, 1919  
Stanford Univ. Publ., Univ. Ser., p. 76.  
Holotype: FMNH 3672. Mexico: Lago Chapala at La Palma.  
Paratypes: 88. Mexico: 2 localities.  
ATHERINIDAE
- Chirostoma jordani** Woolman, 1894  
Bull. U.S. Fish Comm. 1894, 14: 62.  
Lectoparatypes: 10. Mexico: Río de Lerma at Salamanca.  
Lectotype designated by Barbour (1973, Tulane Univ. Stud. Zool. Bot., 18(3): 103).  
ATHERINIDAE
- Chirostoma labarcae** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 113.  
Holotype: FMNH 3640. Mexico: Río Lerma at La Barca.  
Paratypes: 8. Mexico: 2 localities.  
ATHERINIDAE
- Chirostoma mezquital** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 170.  
Holotype: FMNH 4389. Mexico: Río Mezquital at Durango.  
Paratypes: 7. Same locality.  
ATHERINIDAE
- Chirostoma patzcuaro** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 113.  
Holotype: FMNH 3628. Mexico: Lago de Patzcuaro at Patzcuaro.  
Paratypes: 8. Same locality.  
ATHERINIDAE
- Chirostoma regani** Jordan and Hubbs, 1919  
Stanford Univ. Publ., Univ. Ser., p. 74.  
Holotype: FMNH 3687. Mexico: Xochimilco.  
Paratypes: 13. Same locality.  
ATHERINIDAE
- Chirostoma zirahuen** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 114.  
Holotype: FMNH 3609. Mexico: Lago de Zirahuén at Zirahuén.  
Paratypes: 29. Same locality.  
ATHERINIDAE
- Chondrostoma cyri orientalis** Bianco and Banarescu, 1982  
Cybium, Ser. 3, 6(2): 80.  
Paratypes: 2. Iran: Pulwar River near Persepolis.  
CYPRINIDAE
- Chonerhinus amabilis** Roberts, 1982  
Proc. Calif. Acad. Sci., 43(1): 5.  
Paratypes: 2. Indonesia: Kapuas River, southwest of Putussibau.  
TETRAODONTIDAE
- Chonerhinus nefastus** Roberts, 1982  
Proc. Calif. Acad. Sci., 43(1): 10.  
Paratypes: 8. Malaysia: Sarawak, 3 localities.  
TETRAODONTIDAE
- Chonerhinus remotus** Roberts, 1982  
Proc. Calif. Acad. Sci., 43(1): 11.  
Holotype: FMNH 68476. Malaysia: Sabah, mouth of Sungei Deramakot.  
Paratypes: 74. Malaysia: Sabah, 4 localities.  
TETRAODONTIDAE
- Chonerhinus silus** Roberts, 1982  
Proc. Calif. Acad. Sci., 43(1): 12.  
Holotype: FMNH 68477. Malaysia: Sarawak, Rejang basin, Sungei Baleh.  
Paratypes: 56. Malaysia: Sarawak, 5 localities.  
TETRAODONTIDAE
- Chromis alta** Greenfield and Woods, 1980  
Copeia, 1980(4): 630.  
Paratypes: 14. Mexico and Ecuador, Galápagos: 2 localities.  
POMACENTRIDAE
- Chromis atripectoralis** Welander and Schultz, 1951  
J. Wash. Acad. Sci., 41(3): 107.  
Paratypes: 26. Philippine and Marshall Islands: 4 localities.  
POMACENTRIDAE
- Chromis limbaughi** Greenfield and Woods, 1980  
Copeia, 1980(4): 627.  
Paratypes: 12. Mexico: Baja California, 3 localities.  
POMACENTRIDAE
- Chromis meridiana** Greenfield and Woods, 1980  
Copeia, 1980(4): 634.  
Holotype: FMNH 75774. Chile: Isla San Felix.  
Paratypes: 7. Same locality.  
POMACENTRIDAE
- Chromis mossambicus** Peters, 1852  
Monatsber. Akad. Wiss. Berl., 1852: 681.  
Syntypes: 2, FMNH 54267 (CM 2898). Mozambique.  
CICHLIDAE

- Chromis randalli** Greenfield and Hensley, 1970  
Copeia, 1970(4): 689.  
Paratype. Easter Island: Motu Nui.  
POMACENTRIDAE
- Chromis sparsidens** Hilgendorf, 1905  
Zool. Jahrb. Abt. Syst. Geogr. Biol. Tiere, 22(4): 408.  
Syntypes: 2, FMNH 54272 (CM 2902). Tanzania: Manyara Lake.  
CICHLIDAE
- Chromis villadolidi** Jordan and Tanaka, 1927  
Ann. Carnegie Mus., 17(12): 287.  
Holotype: FMNH 59185 (CM 8328). Sea of Japan: Kiusiu.  
POMACENTRIDAE
- Chromis weberi** Fowler and Bean, 1928  
Bull. U.S. Natl. Mus., 100(7): 41.  
Paratype. Philippine Islands: Tataan Island.  
POMACENTRIDAE
- Chromis woodsi** Bruner and Arnam, 1979  
Fieldiana Zool., 73(3): 50.  
Holotype: FMNH 85011. Western Indiana Ocean: Somalia coast.  
Paratypes: 13. Western Indian Ocean: 5 localities.  
POMACENTRIDAE
- Cichlasoma acutum** Miller, 1907  
Bull. Am. Mus. Nat. Hist., 23(2): 117.  
Paratype. Guatemala: Río Kilagua at los Amates.  
CICHLIDAE
- Cichlasoma alfari** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(5): 148.  
Holotype: FMNH 6032. Costa Rica: Turrialba.  
Paratypes: 4. Same locality.  
One specimen in the paratype series, FMNH 6033, is outside the size range given in the original description.  
CICHLIDAE
- Cichlasoma boliviense** Kullander, 1983  
A revision of the South American cichlid genus *Cichlasoma* . . . , Stockholm, p. 165.  
Paratypes: 2. Bolivia: Santa Cruz de la Sierra.  
CICHLIDAE
- Cichlasoma calobrense** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 90.  
Holotype: FMNH 7600. Panama: Río Calobre.  
Paratypes: 38. Panama: 7 localities.  
CICHLIDAE
- Cichlasoma centrale** Meek, 1906  
Field Mus. Nat. Hist. Zool. Ser., 7(3): 94.  
Holotype: FMNH 5510. Guatemala: Caballo Blanco.  
CICHLIDAE
- Cichlasoma dorsatum** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 123.  
Holotype: FMNH 5971. Nicaragua: Lago Managua at Managua.  
Paratypes: 4. Nicaragua: 3 localities.  
CICHLIDAE
- Cichlasoma eigenmanni** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 119.  
Holotype: FMNH 3712. Mexico: Río Tehuacán at Venta Salada.  
Paratype. Same locality.  
CICHLIDAE
- Cichlasoma evermanni** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 214.  
Holotype: FMNH 4727. Mexico: Oaxaca, Tehuantepec.  
Paratypes: 5. Same locality.  
CICHLIDAE
- Cichlasoma frontale** Meek, 1909  
Field Mus. Nat. Hist. Zool. Ser., 7(7): 210.  
Holotype: FMNH 6358. Costa Rica: Turru-bares.  
Paratype. Same locality.  
CICHLIDAE
- Cichlasoma grammodes** Taylor and Miller, 1980  
Occas. Pap. Mus. Zool. Univ. Mich., 693: 2.  
Paratypes: 9. Mexico: Río Chico at Chiapa.  
CICHLIDAE
- Cichlasoma granadense** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 121.  
Holotype: FMNH 5951. Nicaragua: Lago Nicaragua at Granada.  
Paratypes: 9. Nicaragua: 4 localities.  
CICHLIDAE
- Cichlasoma hartwegi** Taylor and Miller, 1980  
Occas. Pap. Mus. Zool. Univ. Mich., 693: 8.  
Paratype. Mexico: Río Salado near Chiapilla.  
CICHLIDAE
- Cichlasoma hedricki** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 208.  
Holotype: FMNH 4673. Mexico: Oaxaca, Obispo.  
Paratypes: 42. Mexico: 3 localities.  
CICHLIDAE

- Cichlasoma meeki** Hildebrand, 1925  
Bull. Bur. Fish., 41: 275.  
Paratypes: 11. El Salvador: 4 localities.  
CICHLIDAE
- Cichlasoma milleri** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(5): 142.  
Holotype: FMNH 5529. Guatemala: El Rancho.  
Paratypes: 11. Same locality.  
CICHLIDAE
- Cichlasoma mojarra** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 217.  
Holotype (?): FMNH 4719. Mexico: Oaxaca, San Gerónimo.  
See comment in Grey (1947, p. 133); FMNH 4719 agrees closely with published size and illustration. Published number FMNH 4718 is a lot identified by Meek as *C. melanurum*, and recently by R. Miller as *C. zonatum*.  
CICHLIDAE
- Cichlasoma nigratum** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 128.  
Holotype: FMNH 5979. Nicaragua: Lago Nicaragua, Granada.  
CICHLIDAE
- Cichlasoma ornatum gephyrum** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 205.  
Holotype: FMNH 58614 (CM 7639). Colombia: Río Dagua at Córdova. Paratypes: 4. Colombia: 3 localities.  
CICHLIDAE
- Cichlasoma paranaense** Kullander, 1983  
A revision of the South American cichlid genus *Cichlasoma* . . . , Stockholm, p. 241.  
Paratypes: 3. Brazil: Rio Tietê at Salto das Cruces.  
CICHLIDAE
- Cichlasoma punctatum** Meek, 1909  
Field Mus. Nat. Hist. Zool. Ser., 7(7): 210.  
Holotype: FMNH 6478. Costa Rica: Buenos Aires de Térraba.  
Paratypes: 10. Same locality.  
CICHLIDAE
- Cichlasoma spilotum** Meek, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(7): 73.  
Holotype: FMNH 7686. Costa Rica: Victoria.  
Paratypes (?): 10. Costa Rica: Zent.  
See comment in Grey (1947, p. 134); status of paratypes uncertain.  
CICHLIDAE
- Cichlasoma tenue** Meek, 1906  
Field Mus. Nat. Hist. Zool. Ser., 7(3): 94.  
Holotype: FMNH 3781. Mexico: Veracruz, Achotal.  
Paratypes: 4. Same locality.  
CICHLIDAE
- Cichlasoma tuba** Meek, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(7): 73.  
Holotype: FMNH 7681 MISSING. Costa Rica: Victoria.  
Paratype (?). Same locality.  
See comment in Grey (1947, pp. 134–135); status of the paratype uncertain.  
CICHLIDAE
- Cichlasoma tuyrense** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 89.  
Holotype: FMNH 7599. Panama: Río Tuyra, Boca de Cupe.  
Paratypes: 65. Panama: 6 localities.  
CICHLIDAE
- Cichlasoma umbriferum** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 88.  
Holotype: FMNH 7598. Panama: Río Cupe at Cituro.  
Paratypes: 20. Panama: 3 localities.  
CICHLIDAE
- Cichlasoma urophthalmus mayorum** Hubbs, 1936  
Carnegie Inst. Wash., Publ. 457: 277.  
Paratypes: 3. Mexico: Yucatán, Chichen Itza, Xtolok Cenote.  
CICHLIDAE
- Cichlasoma zonatum** Meek, 1905  
Proc. Biol. Soc. Wash., 18: 245.  
Holotype: FMNH 3776. Mexico: Oaxaca, Nilotpec.  
Paratype. Same locality.  
CICHLIDAE
- Cirricaecula johnsoni** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 50.  
Paratype. Marshall Islands: Rongerik Atoll, Bock Island.  
OPHICHTHIDAE
- Clarias malaris** Nichols and Griscom, 1917  
Bull. Am. Mus. Nat. Hist., 37(25): 705.  
Paratype. Zaïre: Stanleyville.  
CLARIIDAE

- Clarias mossambicus** Peters, 1852  
 Monatsber. Akad. Wiss. Berl., 1852: 682.  
 Syntype (?): FMNH 54274 (CM 2904). Mozambique: Quelimane.  
 This precise locality was not mentioned in original description.  
 CLARIIDAE
- Clinocottus analis australis** Hubbs, 1926  
 Occas. Pap. Mus. Zool. Univ. Mich., 171: 11.  
 Paratypes: 146. California: 4 localities.  
 COTTIDAE
- Cnesterodon carnegiei** Haseman, 1911  
 Ann. Carnegie Mus., 7(3-4): 385.  
 Holotype: (CM 2868) MISSING. Brazil: Rio Iguaçu at Serrinha Paraná.  
 Paratypes: 120. Brazil and Uruguay: 4 localities.  
 POECILIIDAE
- Cobitis aurata vallahica** Nalbant, 1957  
 Senck. Biol., 38(3/4): 209.  
 Paratypes: 5. Romania: Ialomitza River at Crivina.  
 COBITIDAE
- Cobitis taenia melanoleuca** Nichols, 1925  
 Am. Mus. Nov., 170: 3.  
 Paratype. China: Shansi.  
 COBITIDAE
- Coelorrhynchus gilberti** Jordan and Hubbs, 1925  
 Mem. Carnegie Mus., 10(2): 327.  
 Holotype: FMNH 59401 (CM 7960). Japan: market at Shizuoka.  
 MACROURIDAE
- Compsura heterura** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 61.  
 Holotype: FMNH 57825 (CM 6808). Brazil: Rio Itapicurú at Queimadas.  
 Paratypes: 11. Same locality.  
 CHARACIDAE
- Conger jordani** Kanazawa, 1958  
 Proc. U.S. Natl. Mus., 108(3400): 250.  
 Paratype. Japan: Tokyo Market.  
 CONGRIDAE
- Copeina osgoodi** Eigenmann, 1922  
 Mem. Carnegie Mus., 9(1): 232.  
 Holotype: FMNH 15352. Peru: mouth of Río Yavary at Nazareth.  
 Paratypes: 6. Same locality.  
 LEBIASINIDAE
- Copidoglanis novaeguineae** Weber, 1908  
 Nova Guinea, 5(2): 226.  
 Syntype: FMNH 52388 (CM 644). New Guinea: Lake Sentani.  
 PLOTOSIDAE
- Coregonus stanleyi** Kendall, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 366.  
 Paratypes: 2. Maine: Aroostook County, Eagle Lake.  
 SALMONIDAE
- Coripareius septentrionalis** Nichols, 1925  
 Am. Mus. Nov., 181: 2.  
 Paratype: MISSING. China: Mongolia, Yellow River at Paotow.  
 CYPRINIDAE
- Corydoras australe** Eigenmann and Ward, 1907  
 Ann. Carnegie Mus., 4(2): 123.  
 Paratype. Paraguay: Corumbá.  
 CALLICHTHYIDAE
- Corydoras blochi** Nijssen, 1971  
 Beaufortia, 19(250): 92.  
 Holotype: FMNH 75951. Guyana: Moco Creek near Lehtem.  
 Paratypes: 2. Brazil: Rio Amazonas at Santarém.  
 CALLICHTHYIDAE
- Corydoras bondi coppenamensis** Nijssen, 1970  
 Beaufortia, 18(230): 19.  
 Paratypes: 2. Surinam: Tributary of Coppename River.  
 CALLICHTHYIDAE
- Corydoras ellisae** Gosline, 1940  
 Stanford Ichthyol. Bull., 2(1): 17.  
 Holotype: FMNH 57361 (CM 3522). Paraguay: Arroyo Poná at Sapucay.  
 Paratype. Paraguay: near Sapucay.  
 CALLICHTHYIDAE
- Corydoras heteromorphus** Nijssen, 1970  
 Beaufortia, 18(230): 22.  
 Paratypes: 2. Surinam: tributary of Coppename River.  
 CALLICHTHYIDAE
- Corydoras schwartzi surinamensis** Nijssen, 1970  
 Beaufortia, 18(230): 39.  
 Paratype. Surinam: tributary of Coppename River.  
 CALLICHTHYIDAE

- Corydoras weitzmani** Nijssen, 1971  
Beaufortia, 19(250): 91.  
Holotype: FMNH 69934. Peru: Río Vilcanota system at Cuzco.  
Paratypes: 2. Same locality.  
CALLICHTHYIDAE
- Corymbophanes andersoni** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 5.  
Holotype: FMNH 52675 (CM 1001). Guyana: Potaro River at Aruataima Falls.  
LORICARIIDAE
- Coryphaenoides bonanox** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6: 305.  
Holotype: FMNH 57520 (CM 6462). Japan: Sagami Bay.  
MACROURIDAE
- Coryphopterus punctipectophorus** Springer, 1960  
Bull. Mar. Sci. Gulf Caribb., 10(2): 237.  
Paratypes: 3. Gulf of Mexico: off Florida.  
GOBIIDAE
- Corythoichthys elerae** Evermann and Seale, 1907  
Bull. Bur. Fish. 1906, 26: 57.  
Paratype. Philippine Islands: Sorsogon, Bacon. Misspelled *Corythoichthys* in Evermann and Seale (1907).  
SYNGNATHIDAE
- Corythoichthys polynotatus** Dawson, 1977  
Copeia, 1977(2): 331.  
Paratypes: 4. Philippine Islands: Negros Island.  
SYNGNATHIDAE
- Cotylis nigripinnis woodsi** Schultz, 1944  
Proc. U.S. Natl. Mus., 96(3187): 68.  
Holotype: FMNH 41794. Cocos Island: Wafer Bay.  
GOBIESOCIDAE
- Creagrutus beni** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 172.  
Holotype: FMNH 54585 (CM 3216). Bolivia: Río Beni at Villa Bella.  
CHARACIDAE
- Creagrutus brevipinnis** Eigenmann, 1913  
Indiana Univ. Stud., 18: 10.  
Holotype: FMNH 56095 (CM 4887). Colombia: Piedra de Moler.  
Paratypes: 59. Colombia: 3 localities.  
CHARACIDAE
- Creagrutus caucanus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 9.  
Holotype: FMNH 56104 (CM 4895). Colombia: Paila.  
Paratypes: 19. Colombia: 4 localities.  
CHARACIDAE
- Creagrutus magdalenae** Eigenmann, 1913  
Indiana Univ. Stud., 18: 8.  
Holotype: FMNH 56088 (CM 4880). Colombia: Girardot.  
Paratypes: 108. Colombia: 3 localities.  
CHARACIDAE
- Creagrutus melanzonus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 30.  
Holotype: FMNH 52705 (CM 1067). Guyana: Crab Falls.  
Paratypes: 2. Guyana: 2 localities.  
CHARACIDAE
- Creagrutus notropoides** Meek and Hildebrand, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.  
Holotype: FMNH 7574. Panama: Río Indio.  
Paratypes: 67. Panama: 5 localities.  
CHARACIDAE
- Creagrutus simus** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 85.  
Holotype: FMNH 7591. Panama: Río Cupe, Cituro.  
Paratypes: 232. Panama: 5 localities.  
CHARACIDAE
- Crenicara altispinosa** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 344.  
Holotype: FMNH 54090 (CM 2639). Bolivia: Río Mamoré, below mouth of Río Guaporé.  
Paratypes: 7. Bolivia: 2 localities.  
CICHLIDAE
- Crenicichla alta** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 516.  
Holotype: FMNH 53777 (CM 2274). Guyana: Gluck Island.  
Paratypes: 35. Guyana: 13 localities.  
CICHLIDAE
- Crenicichla dorsocellata** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 355.  
Holotype: FMNH 54155 (CM 2721). Brazil: Rio Parahyba at Campos.  
Paratype. Brazil: Santarem.  
CICHLIDAE

- Crenicichla iguassuensis** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 352.  
Holotype: FMNH 54159 (CM 2725). Brazil: Rio Iguaçu at Porto União da Victoria.  
Paratypes: 4. Brazil: 2 localities.  
CICHLIDAE
- Crenicichla jaguarensis** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 351.  
Holotype: FMNH 54157 (CM 2723). Brazil: Rio Grande of the Paraná at Jaguará.  
Paratype. Same locality.  
CICHLIDAE
- Crenicichla santaremensis** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 354.  
Holotype: FMNH 54153 (CM 2719). Brazil: near Santarém.  
Paratype. Same locality.  
CICHLIDAE
- Crenicichla simoni** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 345.  
Holotype: FMNH 54097 (CM 2646). Brazil: São Luiz de Cáceres.  
Paratypes: 7. Brazil, Bolivia, and Paraguay: 4 localities.  
CICHLIDAE
- Crossochir koelzi** Hubbs, 1933  
Proc. U.S. Natl. Mus., 82(2962): 4.  
Paratypes: 6. California: San Diego.  
EMBIOTOCIDAE
- Crossostoma fascicauda** Nichols, 1926  
Am. Mus. Nov., 224: 2.  
Paratype. China: Fukien, Fuching Hsien.  
COBITIDAE
- Cryptocentrus geniornatus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 428.  
Holotype: FMNH 17364. Indonesia: Waigiu Island.  
GOBIIDAE
- Cryptotomus crassiceps** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 32.  
Lectotype: FMNH 4964. Bermuda: Cooper's Island.  
Lectotype designated by Grey (1947, p. 138).  
SCARIDAE
- Ctenogobius aterrimus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 423.  
Holotype: FMNH 17384. Solomon Islands: Kulambangra Island.  
GOBIIDAE
- Ctenogobius leavelli** Herre, 1935  
Lingnan Sci. J. Canton, 14(3): 396.  
Paratypes: 6. China: Kwangsi, Wuchow.  
GOBIIDAE
- Ctenogobius malekulae** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 423.  
Holotype: FMNH 17385. New Hebrides Islands: Malekula Island, Bushman's Bay.  
GOBIIDAE
- Ctenogobius myxodermus** Herre, 1935  
Lingnan Sci. J. Canton, 14(3): 395.  
Paratypes: 10. China: 2 localities.  
GOBIIDAE
- Ctenogobius opalescens** Herre, 1936  
Bull. Raffles Mus., 12: 15.  
Paratypes: 3. Singapore.  
GOBIIDAE
- Ctenogobius villadolidi** Herre, 1936  
Philipp. J. Sci., 59(3): 361.  
Paratypes: 2. Philippine Islands: Negros Island, Dumaguete.  
GOBIIDAE
- Ctenogobius virgatulus** Jordan and Snyder, 1901  
Proc. U.S. Natl. Mus., 24(1244): 63.  
Paratype: MISSING. Japan: Sagami, Misaki.  
GOBIIDAE
- Ctenogobius waigiensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 424.  
Holotype: FMNH 17413. Indonesia: Waigiu Island.  
Paratypes: 77. Same locality.  
GOBIIDAE
- Ctenopoma multispinis** Peters, 1844  
Monatsber. Akad. Wiss. Berl., 1844: 34 (not seen).  
Syntype: FMNH 54262 (CM 2891). Mozambique: Quelimane.  
ANABANTIDAE
- Culter aokii** Oshima, 1919  
Ann. Carnegie Mus., 12(2–4): 250.  
Holotype: FMNH 59110 (CM 8248). Taiwan: Jitsugetsutan (Lake Candidius).  
CYPRINIDAE

- Curimata cerasina** Vari, 1983  
Acta Biol. Venez., 11(4): 30.  
Paratype. Venezuela: Guárico, pit along side of road from Calabozo to Camaguán.  
CURIMATIDAE
- Curimata incompta** Vari, 1983  
Acta Biol. Venez., 11(4): 34.  
Paratypes: 2. Venezuela: 2 localities.  
CURIMATIDAE
- Curimatus atratoensis** Eigenmann, 1912  
Indiana Univ. Stud., 16: 19.  
Holotype: FMNH 56024 (CM 4814). Colombia: Quibdó.  
Paratypes: 48. Same locality.  
CURIMATIDAE
- Curimatus gillii** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 510.  
Paratype. Paraguay: Chagalalina.  
CURIMATIDAE
- Curimatus issororoensis** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 266.  
Holotype: FMNH 53647 (CM 2119). Guyana: Issorora Rubber Plantation.  
Paratype. Same locality.  
CURIMATIDAE
- Curimatus metae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 230.  
Holotype: FMNH 55143 (CM 3845). Colombia: Quebrada Cramalote at Villavicencio.  
Paratypes: 2. Same locality.  
CURIMATIDAE
- Curimatus morawhannae** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 266.  
Holotype: FMNH 53650 (CM 2122). Guyana: Morawhanna.  
Paratypes: 14. Guyana: 2 localities.  
CURIMATIDAE
- Curimatus patiae** Eigenmann, 1914  
Indiana Univ. Stud., 19: 12.  
Holotype: FMNH 56554 (CM 5368). Colombia: Barbacoas.  
Paratypes: 3. Same locality.  
CURIMATIDAE
- Curimatus peruanus** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 104.  
Paratype. Peru: Río Chira at Sullana.  
CURIMATIDAE
- Cyclopium chapmani** Eigenmann, 1912  
Indiana Univ. Stud., 16: 13.  
Holotype: FMNH 56071 (CM 4863). Colombia: Boquía.  
Paratypes: 2. Same locality.  
ASTROBLEPIDAE
- Cyclopium pirrense** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 83.  
Holotype: FMNH 7586. Panama: Río Cana at Cana.  
Paratypes: 33. Same locality.  
ASTROBLEPIDAE
- Cyclopium trifasciatum** Eigenmann, 1912  
Indiana Univ. Stud., 16: 14.  
Holotype: FMNH 56076 (CM 4868). Colombia: Caldas.  
Paratypes: 40. Colombia: 2 localities.  
ASTROBLEPIDAE
- Cyclopium unifasciatum** Eigenmann, 1912  
Indiana Univ. Stud., 16: 15.  
Holotype: FMNH 56079 (CM 4871). Colombia: Caldas.  
Paratypes: 54. Same locality.  
ASTROBLEPIDAE
- Cyclopium vanceae** Eigenmann, 1913  
Ann. Carnegie Mus., 8(3-4): 421.  
Holotype: FMNH 56066 (CM 4856). Peru: Queta, brook southeast of Tarma.  
ASTROBLEPIDAE
- Cyclopium ventrale** Eigenmann, 1912  
Indiana Univ. Stud., 16: 15.  
Holotype: FMNH 56074 (CM 4866). Colombia: Caldas.  
Paratypes: 15. Colombia: 2 localities.  
ASTROBLEPIDAE
- Cynodonichthys tenuis** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 101.  
Holotype: FMNH 4643. Mexico: Oaxaca, El Hule.  
APLOCHEILIDAE
- Cynoglossus heterolepis** Weber, 1910  
Notes Leiden Mus., 32: 237.  
Syntype: FMNH 52501 (CM 759). New Guinea: Lorentz River.  
CYNOGLOSSIDAE



- Cynopotamus essequibensis** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 403.  
Holotype: FMNH 53673 (CM 2146). Guyana: Potaro Landing.  
Paratypes: 4. Guyana: 2 localities.  
CHARACIDAE
- Cyprinodon bifasciatus** Miller, 1968  
Occas. Pap. Mus. Zool. Univ. Mich., 659: 1.  
Paratypes: 20. Mexico: Río Puente Colorado near Cuatro Ciénegas.  
CYPRINODONTIDAE
- Cyprinodon dearborni** Meek, 1909  
Field Mus. Nat. Hist. Zool. Ser., 7(7): 208.  
Holotype: FMNH 6484. Lesser Antilles: Curaçao, Willemstad.  
Paratypes: 13. Same locality.  
CYPRINODONTIDAE
- Cyprinodon fontinalis** Smith and Miller, 1980  
Proc. Biol. Soc. Wash., 93(2): 405.  
Paratypes: 15. Mexico: Chihuahua, Ojo de Car-bonera.  
CYPRINODONTIDAE
- Cyprinodon meeki** Miller, 1976  
Bull. South. Calif. Acad. Sci., 75(2): 73.  
Paratypes: 77. Mexico: 2 localities.  
CYPRINODONTIDAE
- Cyprinodon orthonotus** Peters, 1844  
Monatsber. Akad. Wiss. Berl., 1844: 35 (not seen).  
Syntype: FMNH 54289 (CM 2920). Mozam-bique: Quellimane.  
CYPRINODONTIDAE
- Cypselurus alienus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 392.  
Holotype: FMNH 17224. Cuba: south coast, near Santiago.  
EXOCOETIDAE
- Cypselurus antoncichi** Woods and Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 181.  
Paratypes: 2. Marshall Islands: 2 localities.  
EXOCOETIDAE
- Dangila sabana** Inger and Chin, 1962  
Fieldiana Zool., 45: 94.  
Holotype: FMNH 44778. Malaysia: Sabah, Danau, Bukit Garam.  
Paratypes: 65. Malaysia: Sabah, 3 localities.  
CYPRINIDAE
- Dasyatis ushieii** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 114.  
Holotype: FMNH 59380 (CM 7778). Japan: Mikawa Bay.  
DASYATIDAE
- Dasyscopelus orientalis** Gilbert, 1913  
Mem. Carnegie Mus., 6(2): 70.  
Holotype: FMNH 55834 (CM 4613). Japan: Sa-gami Bay, Misaki.  
Paratypes: 52. Same locality.  
MYCTOPHIDAE
- Decapterus dayi** Wakiya, 1924  
Ann. Carnegie Mus., 15(2–3): 158.  
Holotype: FMNH 59468 (CM 7711). Taiwan.  
CARANGIDAE
- Decapterus tabl** Berry, 1968  
Contrib. Mar. Sci., 13: 152.  
Paratype. Western Atlantic: off Colombia.  
CARANGIDAE
- Dermatocheir catablepta** Durbin, 1909  
Ann. Carnegie Mus., 6(1): 55.  
Holotype: FMNH 53553 (CM 1198). Guyana: Tumatumari, above the falls.  
CHARACIDAE
- Deuterodon acanthogaster** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 179.  
Holotype: FMNH 54748 (CM 3395). Brazil: Corumbá.  
Paratypes: 14. Brazil: 2 localities.  
CHARACIDAE
- Deuterodon atracaudata** Meek and Hildebrand, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.  
Holotype: FMNH 7573. Panama: Canal Zone, Río Frijoles.  
Paratypes: 497. Panama: 22 localities.  
CHARACIDAE
- Deuterodon pinnatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 25.  
Holotype: FMNH 53525 (CM 1046). Guyana: Potaro River at Amatuk.  
Paratypes: 46. Guyana: 5 localities.  
CHARACIDAE
- Deuterodon potaroensis** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 27.  
Holotype: FMNH 52967 (CM 1053). Guyana: Potaro River at Amatuk.  
Paratypes: 2. Guyana: 2 localities.  
CHARACIDAE

- Diaphus gigas** Gilbert, 1913  
 Mem. Carnegie Mus., 6(2): 93.  
 Holotype: FMNH 55825 (CM 4601). Japan: Sagami Bay, Misaki.  
 Paratype: Same locality.  
 MYCTOPHIDAE
- Diaphus glandulifer** Gilbert, 1913  
 Mem. Carnegie Mus., 6(2): 90.  
 Paratypes: 3. Japan: Sagami Bay.  
 MYCTOPHIDAE
- Diaphus latus** Gilbert, 1913  
 Mem. Carnegie Mus., 6(2): 95.  
 Holotype: FMNH 56208 (CM 4604). Japan: Sagami Bay, Misaki.  
 MYCTOPHIDAE
- Diaphus sagamiensis** Gilbert, 1913  
 Mem. Carnegie Mus., 6(2): 96.  
 Holotype: FMNH 55831 (CM 4608). Japan: Sagami Bay, Misaki.  
 Paratypes: 3. Same locality.  
 MYCTOPHIDAE
- Diapterus evermanni** Meek and Hildebrand, 1925  
 Field Mus. Nat. Hist. Zool. Ser., 15(2): 594.  
 Paratypes: 2. Panama: Fox Bay, Colón.  
 GERREIDAE
- Diceratias pileatus** Uwate, 1979  
 Copeia, 1979(1): 140.  
 Paratypes: 2. Western Atlantic Ocean: off Surinam.  
 DICERATIIDAE
- Dicrolene kanazawai** Grey, 1958  
 Fieldiana Zool., 39(16): 170.  
 Paratypes: 6. Gulf of Mexico.  
 OPHIDIIDAE
- Dionda diaboli** Hubbs and Brown, 1956  
 Southwest Nat., 1(2): 69.  
 Paratypes: 2. Texas: Devil's Lake.  
 CYPRINIDAE
- Dionda erimyzonops** Hubbs and Miller, 1974  
 Occas. Pap. Mus. Zool. Univ. Mich., 671: 4.  
 Paratypes: 3. Mexico: Tamaulipas, Forlón.  
 CYPRINIDAE
- Diphyacantha chocoensis** Henn, 1916  
 Ann. Carnegie Mus., 10(1-2): 114.  
 Paratypes: 10. Colombia: creek near mouth of Río Calima.  
 POECILIIDAE
- Diplectrum labarum** Rosenblatt and Johnson, 1974  
 Calif. Fish Game, 60(4): 181.  
 Paratypes: 57. Mexico: Baja California, Bahía Santa Inez.  
 SERRANIDAE
- Diplectrum pacificum** Meek and Hildebrand, 1925  
 Field Mus. Nat. Hist. Zool. Ser., 15(2): 477.  
 Paratypes: 3. Panama: Naos Island.  
 SERRANIDAE
- Diplodus bermudensis** Caldwell, 1965  
 Fieldiana Zool., 44(23): 217.  
 Holotype: FMNH 72520. Bermuda: Saint George's Island, Whalebone Bay.  
 Paratypes: 179. Bermuda: 11 localities.  
 SPARIDAE
- Diplogonurus maderensis** Noronha, 1926  
 Ann. Carnegie Mus., 16(3-4): 381.  
 Holotype: MISSING. Cama de Lobos.  
 The type, apparently the head (dried?) illustrated in Noronha, was never received by the Carnegie Museum (Henn, 1928, p. 79).  
 GEMPYLIDAE
- Diretmus pauciradiatus** Woods, 1973  
 Mem. Sears Fnd. Mar. Res., 1(6): 296.  
 Holotype: FMNH 66797. Southern Caribbean Sea.  
 Paratypes: 15. Caribbean Sea and Gulf of Mexico: 12 localities.  
 DIRETMIDAE
- Disparichthys fluviatilis** Herre, 1935  
 Field Mus. Nat. Hist. Zool. Ser., 18(12): 384.  
 Holotype: FMNH 17192. New Guinea: Sepik River basin, brook at Marienberg.  
 CARAPIDAE
- Distichodus stigmaturus** Fowler, 1935  
 Ann. Transvaal Mus., 16(2): 260.  
 Paratypes: 3. Botswana: Chobe River near Kasana.  
 CITHARINIDAE
- Dollfusichthys sinuarabici** Chabanaud, 1931  
 Bull. Soc. Zool. Fr., 56: 304.  
 Paratype: Egypt: Bay of Suez.  
 CYNOGLOSSIDAE
- Doraops zuloagai** Schultz, 1944  
 Proc. U.S. Natl. Mus., 94(3172): 271.  
 Paratype: Venezuela: Río Apón near mouth of Río Cogollo.  
 DORADIDAE

- Doras lentiginosus** Eigenmann, 1917  
Ann. Carnegie Mus., 11(3-4): 401.  
Holotype: FMNH 58056 (CM 7048). Brazil: Santarém.  
Paratype. Brazil: Manaus.  
DORADIDAE
- Dormitator gymnocephalus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 523.  
Holotype: FMNH 53922 (CM 2438). Guyana: Konawaruk.  
ELEOTRIDAE
- Dorosoma anale** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 93.  
Holotype: FMNH 4637. Mexico: Oaxaca, El Hule.  
Paratypes: 15. Mexico: 4 localities.  
CLUPEIDAE
- Dorosoma chavesi** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 112.  
Holotype: FMNH 5928. Nicaragua: Laguna Janicero, Granada.  
Paratypes: 73. Nicaragua: 4 localities.  
Grey (1947, p. 140) noted that the type, FMNH 5928, was MISSING. We now have 1 specimen labeled type and a separated lot of 6 labeled paratypes with both lots bearing FMNH 5928, but no indication of who separated the type.  
CLUPEIDAE
- Doryrhamphus melanopleura cooki** Weed and Woods, 1953  
Bull. U.S. Natl. Mus., 202(1): 249.  
Holotype: FMNH 16220. Cook Archipelago: Aitutaki Island.  
Paratype. Same locality.  
SYNGNATHIDAE
- Draconetta oregona** Briggs and Berry, 1959  
Copeia, 1959(2): 129.  
Paratypes: 18. Northeast coast of South America: 4 localities.  
DRACONETTIDAE
- Dysomma abyssale** Kamohara, 1938  
On the offshore bottom-fishes of Prov. Tosa, Shikoku, Japan, p. 12 (not seen).  
Neotype: FMNH 76869. Sagami Sea.  
Neotype designated by Robins and Robins (1976, Proc. Acad. Nat. Sci. Phila., 127(18): 256).  
SYNAPHOBRANCHIDAE
- Echidna unicolor** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 106.  
Paratype. Marshall Islands: Bikini Atoll.  
MURAENIDAE
- Echidna vineta** Jenkins, 1903  
Bull. U.S. Fish Comm. 1902, 22: 429.  
Paratype: MISSING. Hawaiian Islands.  
Published as FMNH 2753 (field no. 263), but the specimen apparently was never received.  
MURAENIDAE
- Echidna zonophaea** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 167.  
Paratype. Hawaiian Islands: Honolulu.  
MURAENIDAE
- Echiodon dawsoni** Williams and Shipp, 1982  
Copeia, 1982(4): 486.  
Paratypes: 2. Gulf of Mexico: off Florida.  
CARAPIDAE
- Ectenias brunneus** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 241.  
Holotype: FMNH 57511 (CM 6453). Japan: Misaki.  
Paratype. Same locality.  
CORYPHAENIDAE
- Ekemblemaria myersi** Stephens, 1963  
Univ. Calif. Publ. Zool., 68: 23.  
Paratypes: 5. Mexico: Golfo de California, Bahía Santa Inez.  
CHAENOPSIDAE
- Eleotris herwerdenii** Weber, 1910  
Notes Leiden Mus., 32: 338.  
Syntype: FMNH 52462 (CM 720). New Guinea: Lorentz River.  
ELEOTRIDAE
- Eleotris heterodon** Weber, 1908  
Nova Guinea, 5(2): 255.  
Syntype: FMNH 52460 (CM 718). New Guinea: Lake Sentani.  
ELEOTRIDAE
- Eleotris isthmensis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 359.  
Holotype: FMNH 8951. Panama: Canal Zone, Mindi Cut.  
Paratypes: 29. Panama: 5 localities.  
ELEOTRIDAE

**Eleotris latifasciatus** Meek and Hildebrand, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(6): 68.  
Holotype: FMNH 7575. Panama: Río Cárdenas  
at Corozal.  
Paratypes: 3. Panama: 2 localities.  
ELEOTRIDAE

**Eleotris macrolepis** Meek, 1914  
Field Mus. Nat. Hist. Zool. Ser., 10(10): 130.  
Holotype: FMNH 7775. Costa Rica: Jesús Ma-  
ría.  
ELEOTRIDAE

**Emblemaria angelica** Böhlke and Mead, 1957  
Not. Nat. (Phila.), 301: 2.  
Paratypes: 24 MISSING. Mexico: Acapulco Bay.  
CHAENOPSIDAE

**Emblemaria hyltoni** Johnson and Greenfield, 1976  
Fieldiana Zool., 70(2): 17.  
Holotype: FMNH 80412. Honduras: Isla Roa-  
tán.  
Paratypes: 2. Same locality.  
CHAENOPSIDAE

**Emblemaria nigra** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 953.  
Paratype. Panama: Porto Bello.  
CHAENOPSIDAE

**Emblemariopsis pricei** Greenfield, 1975  
Copeia, 1975(4): 713.  
Holotype: FMNH 77481. Belize: Glover's Reef.  
Paratype. Same locality.  
CHAENOPSIDAE

**Encaeura evides** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 303.  
Holotype: FMNH 58821 (CM 7931). Japan:  
Wakanoura.  
ELEOTRIDAE

**Enchelycore carychroa** Böhlke and Böhlke, 1976  
Proc. Acad. Nat. Sci. Phila., 127(13): 138.  
Paratypes: 2. Haiti: Gulf of Gonâve to Port-au-  
Prince Bay.  
MURAENIDAE

**Engyprosopon xenandrus** Gilbert, 1905  
Bull. U.S. Fish Comm. 1903, 23(2): 687.  
Paratypes: 2. Hawaiian Islands.  
BOTHIDAE

**Enneapterygius punctulatus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 432.  
Holotype: FMNH 17388. New Hebrides Is-  
lands: Wala Island.  
Paratypes: 2. New Hebrides Islands: 2 localities.  
TRIPTERYGIIDAE

**Enneapterygius waigiensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 434.  
Holotype: FMNH 17494. Indonesia: Waigiu Is-  
land.  
TRIPTERYGIIDAE

**Entomacrodus randalli** Springer, 1967  
Proc. U.S. Natl. Mus., 122(3582): 71.  
Paratypes: 2. Marquesas Islands: Nuku Hiva Is-  
land.  
BLENNIIDAE

**Entomocorus benjamini** Eigenmann, 1917  
Ann. Carnegie Mus., 11(3-4): 403.  
Holotype: FMNH 58019 (CM 7006). Bolivia:  
San Joaquín.  
Paratypes: 9. Bolivia: 2 localities.  
AUCHENIPTERIDAE

**Entonanthias pascalus** Jordan and Tanaka, 1927  
Ann. Carnegie Mus., 17(12): 385.  
Holotype: FMNH 59184 (CM 8327). Japan:  
Okinawa, Naha.  
SERRANIDAE

**Epapterus blohmi** Vari, Jewett, Taphorn, and Gil-  
bert, 1984  
Proc. Biol. Soc. Wash., 97(2): 463.  
Paratypes: 2. Venezuela: Guarico, Laguna Los  
Guácimos.  
AUCHENIPTERIDAE

**Epinephelus elongatus** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 345.  
Paratypes: 3. Marshall Islands: Bikini Atoll, 2  
localities.  
SERRANIDAE

**Epinephelus emoryi** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 340.  
Paratype: Marianas Islands: Rota Island.  
SERRANIDAE

**Epinephelus ionthas** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 32.  
Holotype: FMNH 55797 (CM 4559). Korea:  
Fusan.  
SERRANIDAE

- Epinephelus melanostigma** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 348.  
Paratypes: 2. Samoan Islands: Swains Island.  
SERRANIDAE
- Epinephelus spilotoceps** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 357.  
Paratypes: 2. Marshall Islands: Bikini Atoll, 2 localities.  
SERRANIDAE
- Epinnula orientalis americana** Grey, 1953  
Copeia, 1953(3): 137.  
Holotype: FMNH 46391. Gulf of Mexico.  
Paratypes: 8. Gulf of Mexico: 2 localities.  
GEMPYLIDAE
- Epinnula orientalis pacifica** Grey, 1953  
Copeia, 1953(3): 138.  
Holotype: FMNH 59528. Japan: Kumano-Nada, off Owase.  
Paratype. Same locality.  
GEMPYLIDAE
- Ernstichthys intonsus** Stewart, 1985  
Fieldiana Zool., New Ser., 25: 16.  
Holotype: FMNH 94603. Ecuador: Río Napo at Añangu.  
ASPREDINIDAE
- Etelis evurus** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 184.  
Paratype. Hawaiian Islands: Hilo.  
LUTJANIDAE
- Etheostoma obeyense** Kirsch, 1892  
Bull. U.S. Fish Comm. 1890, 10: 292.  
Paratypes: 36. Kentucky.  
PERCIDAE
- Eucinostomus californiensis elongatus** Meek and Hildebrand, 1925  
Field Mus. Nat. Hist. Zool. Ser., 15(2): 586.  
Paratypes: 52. Panama: Taboga Island.  
GERREIDAE
- Euctenogobius liolepis** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 875.  
Paratypes: 7. Panama: Canal Zone, on beach near Balboa.  
GOBIIDAE
- Euctenogobius panamensis** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 874.  
Paratypes: 3. Panama: Río Culebra at Culebra.  
GOBIIDAE
- Euleptoeleotris clarki** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 352.  
Paratypes: 2. Panama: Canal Zone, Miraflores locks.  
ELEOTRIDAE
- Euleptoeleotris shropshirei** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 353.  
Paratype. Panama: Puerto Pilón.  
ELEOTRIDAE
- Eumegistus illustris** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 36.  
Holotype: FMNH 59325 (CM 3899). Hawaiian Islands: Honolulu Market.  
BRAMIDAE
- Eupomacentrus chrysus** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 32.  
Holotype: FMNH 5025. Bermuda: White's Flat Channel.  
POMACENTRIDAE
- Eupomacentrus rocasensis** Emery, 1972  
Copeia, 1972(2): 330.  
Paratypes: 12. Atlantic Ocean: off Brazil, Atol das Rocas.  
POMACENTRIDAE
- Evarra tlahuacensis** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 85.  
Holotype: FMNH 3681. Mexico: Lago Chalco, Tlahuac.  
CYPRINIDAE
- Evermannella ahlstromi** Johnson and Glodek, 1975  
Copeia, 1975(4): 716.  
Paratypes: 19. Pacific Ocean: 7 localities.  
EVERMANNELLIDAE
- Evermannella sicaria** Rofen, 1963  
Aquatica, Aquatic Res. Inst. Stockton, 1: 1.  
Paratypes: 2. Western North Atlantic: off Bermuda.  
EVERMANNELLIDAE

**Eviota japonica** Jewett and Lachner, 1983  
Proc. Biol. Soc. Wash., 96(4): 788.  
Paratypes: 19. Japan: Ryukyu Islands, Aikawa  
Rikuzen.  
GOBIIDAE

**Evorthodus minutus** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 870.  
Paratype. Panama: Canal Zone, Corozal.  
GOBIIDAE

**Exerpes asper earli** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 94.  
Paratypes: 5. Mexico: Baja California, Laguna  
San Ignacio.  
LABRISOMIDAE

**Farlowella azygia** Eigenmann and Vance, 1917  
Ann. Carnegie Mus., 11(1-2): 299.  
Holotype: FMNH 55091 (CM 3782). Brazil:  
Santarém.  
LORICARIIDAE

**Farlowella hasemani** Eigenmann and Vance, 1917  
Ann. Carnegie Mus., 11(1-2): 301.  
Holotype: FMNH 55089 (CM 3781). Brazil:  
Pará.  
Paratype. Same locality.  
LORICARIIDAE

**Farlowella jauruensis** Eigenmann and Vance, 1917  
Ann. Carnegie Mus., 11(1-2): 300.  
Holotype: FMNH 55088 (CM 3780). Brazil:  
Jaurú.  
LORICARIIDAE

**Felichthys amblops** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 77.  
Holotype: FMNH 7576. Panama: Río Tuyra,  
Marriganti.  
Paratypes: 3. Panama: 2 localities.  
ARIIDAE

**Fitzroyia eigenmanni** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 385.  
Holotype: FMNH 69527 (CM 2866). Brazil:  
creek near Serrinha Paraná.  
Paratypes: 2 (mixed with 8 topotypes). Same  
locality.  
JENYNSIIDAE

**Flammeo scythrops** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 174.  
Paratype. Hawaiian Islands: Honolulu.  
HOLOCENTRIDAE

**Formosania gilberti** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 194.  
Holotype: FMNH 59085 (CM 8222). Taiwan:  
Tamusui River near Shinten.  
CYPRINIDAE

**Fowlerina franciscensis** Eigenmann, 1914  
Indiana Univ. Stud., 20: 45.  
Holotype: FMNH 56426 (CM 5240). Brazil: La-  
goas of Rio Grande, Barreiras.  
Paratypes: 143. Brazil: 10 localities.  
Redescribed as "*Ephippicharax franciscoen-  
sis*, sp. nov." by Eigenmann and Myers  
(1928, Mem. Mus. Comp. Zool. Harv.  
Coll., 43(4): 506).  
CHARACIDAE

**Franzia ardens** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 251.  
Holotype: FMNH 57509 (CM 6451). Japan:  
Misaki.  
SERRANIDAE

**Fundulus cubensis** Eigenmann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 222.  
Paratype. Cuba: Pinar del Río.  
CYPRINODONTIDAE

**Fundulus neumanni** Hilgendorf, 1905  
Zool. Jahrb. Abt. Syst. Geogr. Biol. Tiere,  
22(4): 417.  
Syntype: FMNH 54290 (CM 2921). Tanzania:  
North Ugogo.  
CYPRINODONTIDAE

**Fundulus oaxacae** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 90.  
Holotype: FMNH 3721. Mexico: Río Verde at  
Oaxaca.  
Paratypes: 40. Same locality.  
CYPRINODONTIDAE

**Galeichthys aguadulce** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 9.  
Holotype: FMNH 4678. Mexico: Río Tesecho-  
cán at Pérez.  
ARIIDAE

**Gambusia alvarezi** Hubbs and Springer, 1957  
Tex. J. Sci., 9(3): 310.  
Paratypes: 34. Mexico: Chihuahua, El Ojo de  
San Gregorio.  
POECILIIDAE

***Gambusia bonita* Meek, 1904**

Field Mus. Nat. Hist. Zool. Ser., 5: 132.

Holotype (?): FMNH 4630. Mexico: Veracruz, Río Tonto at Refugio.

Allotype and paratypes: 57. Mexico: 2 localities. FMNH 4630 has 2 specimens. Both smaller than the published size of the holotype; thus both may be paratypes and the holotype lost or mixed with a paratype lot.

POECILIIDAE

***Gambusia caliensis* Eigenmann and Henn, 1916**

In: Henn, Ann. Carnegie Mus., 10(1–2): 113.

Holotype: FMNH 57721 (CM 6700). Colombia: Cali.

POECILIIDAE

***Gambusia cana* Meek and Hildebrand, 1913**

Field Mus. Nat. Hist. Zool. Ser., 10(8): 87.

Holotype: FMNH 7596. Panama: Río Satiganti, Cana.

Allotype and paratypes: 35. Same locality.

POECILIIDAE

***Gambusia cascajalensis* Meek and Hildebrand, 1913**

Field Mus. Nat. Hist. Zool. Ser., 10(8): 86.

Holotype: FMNH 7594. Panama: Río Cascajal at Porto Bello.

Allotype and paratypes: 17. Same locality.

POECILIIDAE

***Gambusia darienensis* Meek and Hildebrand, 1913**

Field Mus. Nat. Hist. Zool. Ser., 10(8): 88.

Holotype: FMNH 7597. Panama: Darién, Río Capetí.

Allotype and paratypes: 61. Panama: 8 localities.

POECILIIDAE

***Gambusia fasciata* Meek, 1904**

Field Mus. Nat. Hist. Zool. Ser., 5: 129.

Holotype: FMNH 4715. Mexico: Oaxaca, San Gerónimo.

Paratypes: 19. Mexico: Oaxaca, 2 localities.

POECILIIDAE

***Gambusia heterochir* Hubbs, 1957**

Tulane Stud. Zool., 5: 3.

Paratypes: 7. Texas: Menard County, Clear Creek.

POECILIIDAE

***Gambusia hurtadoi* Hubbs and Springer, 1957**

Tex. J. Sci., 9(3): 307.

Paratypes: 30. Mexico: Chihuahua, El Ojo de la Hacienda Dolores.

POECILIIDAE

***Gambusia infans* Woolman, 1894**

Bull. U.S. Fish Comm. 1894, 14: 62.

Lectoparatypes: 6. Mexico: Río de Lerma at Salamanca.

Lectotype designated by Jordan and Evermann (1896, Bull. U.S. Natl. Mus., 47(1): 680).

POECILIIDAE

***Gambusia latipunctata* Meek and Hildebrand, 1913**

Field Mus. Nat. Hist. Zool. Ser., 10(8): 87.

Holotype: FMNH 7595. Panama: Arriján.

Paratypes: 16. Same locality.

POECILIIDAE

***Gambusia nigroventralis* Eigenmann and Henn, 1912**

In: Eigenmann, Indiana Univ. Stud., 16: 26.

Holotype: FMNH 56045 (CM 4835). Colombia: Río San Juan at Itsmina.

Paratypes: 5. Same locality.

POECILIIDAE

***Gambusia panuco* Hubbs, 1926**

Misc. Publ. Mus. Zool. Univ. Mich., 16: 30.

Holotype: FMNH 14060. Mexico: San Luis Potosí, Valles.

Allotype and paratypes: 8. Same locality.

POECILIIDAE

***Gambusia parismina* Meek, 1912**

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.

Holotype: FMNH 7678. Costa Rica: Parismina.

Paratypes: 39. Same locality.

POECILIIDAE

***Gambusia regani* Hubbs, 1926**

Misc. Publ. Mus. Zool. Univ. Mich., 16: 28.

Holotype: FMNH 14033. Mexico: Tamaulipas, Forlón.

Allotype and paratypes: 12. Same locality.

POECILIIDAE

***Gambusia turrubarensis* Meek, 1912**

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.

Holotype: FMNH 7676. Costa Rica: Turrubares.

Allotype and paratypes: 95. Same locality.

POECILIIDAE

**Gambusia umbratilis** Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 70.  
Holotype: FMNH 7684. Costa Rica: Guapilis.  
Paratypes: 47. Same locality.  
POECILIIDAE

**Gambusia vittata** Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 26.  
Holotype: FMNH 14046. Mexico: Tamaulipas,  
Forlón.  
Paratypes: 12. Same locality.  
POECILIIDAE

**Gambusia xanthosoma** Greenfield, 1983

Copeia, 1983(2): 459.  
Holotype: FMNH 94188. Grand Cayman Is-  
land: West Bay.  
Paratypes: 10. Same locality.  
POECILIIDAE

**Gambusia yucatana australis** Greenfield, 1985

Copeia, 1985(2): 375.  
Holotype: FMNH 94901. Belize: Saint George's  
Cay, Barrier Reef.  
Paratypes: 58. Belize: 4 localities.  
POECILIIDAE

**Gastromyzon fasciatus** Inger and Chin, 1961

Copeia, 1961(2): 173.  
Holotype: FMNH 68119. Malaysia: Sarawak,  
Sungei Dapu River near mouth of Sungei  
Putai.  
Paratypes: 29. Malaysia: Sarawak, 3 localities.  
HOMALOPTERIDAE

**Gastromyzon lepidogaster** Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 509.  
Paratypes: 8. Malaysia: Sabah, 2 localities.  
HOMALOPTERIDAE

**Gastromyzon megalepis** Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 510.  
Holotype: FMNH 68126. Malaysia: Sarawak,  
tributary of Baleh River.  
Paratypes: 8. Malaysia: Sarawak, 2 localities.  
HOMALOPTERIDAE

**Gastromyzon pauciradiatus** Inger and Chin, 1961

Copeia, 1961(2): 174.  
Holotype: FMNH 68121. Malaysia: Sarawak,  
tributary of Baleh River.  
Paratypes: 21. Malaysia: Sarawak, 2 localities.  
HOMALOPTERIDAE

**Gastromyzon punctulatus** Inger and Chin, 1961

Copeia, 1961(2): 173.  
Holotype: FMNH 68116. Malaysia: Sarawak,  
Sungei Dapu River near mouth of Sungei  
Putai.  
Paratypes: 16. Malaysia: Sarawak, 3 localities.  
HOMALOPTERIDAE

**Gastromyzon ridens** Roberts, 1982

Proc. Calif. Acad. Sci., 42(20): 515.  
Paratype. Indonesia: Kapuas River basin, Sun-  
gei Pinoh.  
HOMALOPTERIDAE

**Gazza achlamys** Jordan and Starks, 1917

Ann. Carnegie Mus., 11: 446.  
Holotype: FMNH 58939 (CM 8074). Sri Lanka:  
Colombo.  
Paratypes: 2. Same locality.  
LEIOGNATHIDAE

**Gelanoglanis stroudi** Böhlke, 1980

Proc. Acad. Nat. Sci. Phila., 132: 152.  
Paratypes: 6. Colombia: Río Manacacias near  
Puerto Gaitán.  
AUCHENIPTERIDAE

**Genycharax tarpon** Eigenmann, 1912

Indiana Univ. Stud., 16: 22.  
Holotype: FMNH 56018 (CM 4808). Colombia:  
Cartago.  
Paratypes: 15. Colombia: 3 localities.  
CHARACIDAE

**Geophagus brasiliensis iporangensis** Haseman,  
1911

Ann. Carnegie Mus., 7(3-4): 364.  
Holotype: FMNH 54202 (CM 2792). Brazil:  
Iporanga.  
Paratypes: 3. Same locality.  
CICHLIDAE

**Geophagus brasiliensis itapicuruensis** Haseman,  
1911

Ann. Carnegie Mus., 7(3-4): 365.  
Holotype: FMNH 54365 (CM 2793). Brazil: Rio  
Itapicuru at Queimadas.  
Paratypes: 7. Brazil: 3 localities.  
CICHLIDAE

**Gephyrocharax caucanus** Eigenmann, 1912

Indiana Univ. Stud., 16: 24.  
Holotype: FMNH 56012 (CM 4802). Colombia:  
Cartago.  
Paratypes: 37. Colombia: 3 localities.  
CHARACIDAE



- Gephyrocharax choacoensis** Eigenmann, 1912  
Indiana Univ. Stud., 16: 23.  
Holotype: FMNH 56016 (CM 4806). Colombia: Istmina.  
Paratypes: 25. Same locality.  
CHARACIDAE
- Gephyrocharax intermedius** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 278.  
Holotype: FMNH 8945. Panama: Río Chame.  
Paratypes: 28. Same locality.  
CHARACIDAE
- Gephyrocharax melanocheir** Eigenmann, 1912  
Indiana Univ. Stud., 16: 24.  
Holotype: FMNH 56049 (CM 4839). Colombia: Bernal Creek near Honda.  
Paratypes: 19. Colombia: 2 localities.  
CHARACIDAE
- Gephyrocharax whaleri** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 254.  
Paratypes: 8. Panama: between Campana and La Venta.  
CHARACIDAE
- Gerhardinus nudus** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 889.  
Paratypes: 7. Panama: tide pools at Panama City.  
GOBIIDAE
- Gibbonsia elegans erroli** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 151.  
Paratypes: 6. Mexico: Baja California, Isla Guadalupe.  
CLINIDAE
- Gibbonsia elegans montereyensis** Hubbs, 1927  
Pap. Mich. Acad. Sci. Arts and Lett., 7: 354.  
Paratypes: 8. California: 2 localities.  
CLINIDAE
- Gibbonsia elegans velifera** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 149.  
Paratypes: 17. California: 2 localities.  
CLINIDAE
- Gibbonsia metzi** Hubbs, 1927  
Pap. Mich. Acad. Sci. Arts and Lett., 7: 360.  
Paratypes: 8. California: 4 localities.  
CLINIDAE
- Gibbonsia norae** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 126.  
Paratypes: 9. Mexico: Baja California, Isla Guadalupe.  
CLINIDAE
- Gila minacae** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 80.  
Holotype: FMNH 3573. Mexico: Río Papigochic at Miñaca.  
CYPRINIDAE
- Gillellus greyae** Kanazawa, 1952  
Fieldiana Zool., 34(7): 88.  
Paratypes: 2. Bermuda and Florida: 2 localities.  
DACTYLOSCOPIDAE
- Girardinus garmani** Eigenmann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 226.  
Paratype: Cuba: Palacios.  
POECILIIDAE
- Gladiogobius ensifer** Herre, 1933  
Copeia, 1933(1): 23.  
Holotype: FMNH 17412. Indonesia: Waigiu Island.  
GOBIIDAE
- Glandulocauda inequalis** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 169.  
Holotype: FMNH 54893 (CM 3555). Brazil: Porto Alegre.  
Paratypes: 4. Same locality.  
CHARACIDAE
- Glandulocauda melanogenys** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 168.  
Holotype: FMNH 54891 (CM 3553). Brazil: São Paulo, Alto da Serra.  
Paratypes: 20. Same locality.  
CHARACIDAE
- Glandulocauda melanopleura** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 170.  
Holotype: FMNH 54895 (CM 3557). Brazil: Rio Iguaçu at Serrinha Paraná.  
Paratypes: 3. Same locality.  
CHARACIDAE
- Glanidium ribeiroi** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 381.  
Holotype: FMNH 54253 (CM 2877). Brazil: Rio Iguaçu, Porto União da Victoria.  
Paratypes: 5. Same locality.  
AUCHENIPTERIDAE

- Glaridichthys falcatus** Eigenmann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 224.  
Paratypes: 4. Cuba: 3 localities.  
POECILIIDAE
- Glossanodon pygmaeus** Cohen, 1958  
Bull. Fla. State Mus. Biol. Sci., 3(3): 156.  
Holotype: FMNH 64345. Gulf of Mexico: off Tortugas.  
Paratypes: 7. Gulf of Mexico: 3 localities.  
ARGENTINIDAE
- Glossogobius abacopus** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 200.  
Holotype: FMNH 52210 (CM 357). Taiwan: Takao.  
Paratype: Same locality.  
GOBIIDAE
- Glossogobius koragensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 419.  
Holotype: FMNH 17365. New Guinea: Sepik River at Koragu.  
Paratype: New Guinea: Kerame River at Ambot.  
GOBIIDAE
- Glossogobius parvus** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 305.  
Holotype: FMNH 59138 (CM 8276). Taiwan: Kizanto Island near Giran.  
GOBIIDAE
- Glossogobius sandakanensis** Inger, 1957  
Fieldiana Zool., 36(3): 393.  
Holotype: FMNH 59526. Malaysia: Sabah, Mankala River at Sandakan.  
GOBIIDAE
- Glyphisodon troscheli** Gill, 1863  
Proc. Acad. Nat. Sci. Phila., 1862: 150.  
Paratype: Mexico: Baja California, Cabo San Lucas.  
Misspelled *Glyphidodon* in Gill (1863).  
POMACENTRIDAE
- Glyptothorax housei** Herre, 1942  
Stanford Ichthyol. Bull., 2(4): 117.  
Paratypes: 3. India: stream in Anaimallai Hills, east of Valparai.  
SISORIDAE
- Gnathodentex oculomaculatus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 407.  
Holotype: FMNH 17335. Solomon Islands: Isabel Island, Tenibuli.  
Paratype: Same locality.  
LETHRINIDAE
- Gnatholepis corlettei** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 418.  
Holotype: FMNH 17367. New Hebrides Islands: Malekula Island, Bushman's Bay.  
Paratypes: 3. Same locality.  
GOBIIDAE
- Gnatholepis knighti** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 204.  
Paratypes: 5. Hawaiian Islands: Hilo.  
GOBIIDAE
- Gnathopogon iijimae** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 219.  
Holotype: FMNH 59096 (CM 8234). Taiwan: Tozen River.  
CYPRINIDAE
- Gnathopogon ishikawae** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 215.  
Paratypes: 3. Japan: Chikugo River at Kurume.  
FMNH 57077 (CM 6014), the published number for the lot which included the holotype has 3 specimens, all too small to be the holotype; perhaps the holotype went to Stanford University.  
CYPRINIDAE
- Gnathopogon longifilis** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 169.  
Holotype: FMNH 58711 (CM 7817). Korea: Ping-yang River.  
Published number CM 7819 is an error.  
CYPRINIDAE
- Ganthopogon majimae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 167.  
Holotype: FMNH 58710 (CM 7816). Korea: Ping-yang River.  
CYPRINIDAE
- Gnathopogon suwae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 166.  
Holotype: FMNH 58708 (CM 7814). Japan: Lake Suwa at Kamisuwa.  
CYPRINIDAE

- Gnathopogon tsuchigae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 170.  
Holotype: FMNH 58712 (CM 7818). Korea:  
Ping-yang River.  
CYPRINIDAE
- Gobiella birtwistlei** Herre, 1934  
Bull. Raffles Mus., 9: 85.  
Syntypes: 18, FMNH 47158 and 47230. Singapore.  
GOBIIDAE
- Gobiesox costaricensis** Meek, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(7): 74.  
Holotype: FMNH 7677. Costa Rica: Zent.  
Paratypes: 4. Costa Rica: 2 localities.  
GOBIESOCIDAE
- Gobiesox fluviatilis** Briggs and Miller, 1960  
Occas. Pap. Mus. Zool. Univ. Mich., 616: 2.  
Paratype. Mexico: Jalisco, Barranca de Barranquitas.  
GOBIESOCIDAE
- Gobiesox fulvus** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(5): 149.  
Holotype: FMNH 6035. Costa Rica: Cocos Island.  
GOBIESOCIDAE
- Gobiesox marijeanae** Briggs, 1960  
Copeia, 1960(3): 215.  
Paratype. Mexico: Islas Tres Marías, Isla San Juanito.  
GOBIESOCIDAE
- Gobiesox mexicanus** Briggs and Miller, 1960  
Occas. Pap. Mus. Zool. Univ. Mich., 616: 7.  
Paratype. Mexico: Oaxaca, Río Tonameca.  
GOBIESOCIDAE
- Gobiesox paradiseus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 432.  
Holotype: FMNH 17404. Ecuador: Islas Galápagos, Isla Eden.  
GOBIESOCIDAE
- Gobiesox potamius** Briggs, 1955  
Stanford Ichthyol. Bull., 6: 93.  
Holotype: FMNH 7815. Costa Rica: Turru-bares.  
Paratype. Same locality.  
Both are from the paratype series of *Gobiesox costaricensis*.  
GOBIESOCIDAE
- Gobiocichla ethelwynnae** Roberts, 1982  
Proc. K. Ned. Akad. Wet. Ser. C Biol. Med. Sci., 85(4): 577.  
Paratypes: 15. Cameroon: Cross River, downstream from Mamfé.  
CICHLIDAE
- Gobiocichla wonderi** Kanazawa, 1951  
Ann. Mag. Nat. Hist., Ser. 12, 4: 378.  
Holotype: FMNH 50423. Mali: Niger River at Sotuba.  
CICHLIDAE
- Gobionellus saepepallens** Gilbert and Randall, 1968  
Not. Nat. (Phila.), 411: 2.  
Paratypes: 2. The Bahamas: Green Turtle Cay, east of Little Abaco.  
GOBIIDAE
- Gobiopsis aporia** Lachner and McKinney, 1978  
Smithson. Contrib. Zool., 262: 12.  
Paratype. Sri Lanka: Galle.  
GOBIIDAE
- Gobiopsis woodsi** Lachner and McKinney, 1978  
Smithson. Contrib. Zool., 262: 21.  
Paratypes: 39. India: Gulf of Mannar, Musal Tivu.  
GOBIIDAE
- Gobio coriparoides** Nichols, 1925  
Am. Mus. Nov., 181: 4.  
Paratypes: 2. China: Shansi, Chin-ssu.  
CYPRINIDAE
- Gobio kesleri banaticus** Banarescu, 1953  
Stud. Cercet. Stiint., 4(1-2): 300.  
Paratypes: 22. Romania: Banat, Timis River.  
CYPRINIDAE
- Gobio rivuloides** Nichols, 1925  
Am. Mus. Nov., 181: 5.  
Paratypes: 3. China: Shansi, Niang-tze-kwan.  
CYPRINIDAE
- Gobio soldatovi minulus** Nichols, 1925  
Am. Mus. Nov., 181: 3.  
Paratypes: 4. China: Shansi, Kweihwa.  
CYPRINIDAE
- Gobius claytonii** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 121.  
Holotype: FMNH 3740. Mexico: Río San Francisco at La Antigua.  
Paratypes: 8. Same locality.  
GOBIIDAE

- Gobius cliffordpopei** Nichols, 1925  
Am. Mus. Nov., 185: 5.  
Paratypes: 4. China: Hunan, Tungting Lake.  
GOBIIDAE
- Gobius daguae** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 685.  
Holotype: FMNH 58479 (CM 7481). Colombia: mouth of Río Dagua.  
Paratype. Same locality.  
GOBIIDAE
- Gobius parvus** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 121.  
Holotype: FMNH 3738. Mexico: Río San Francisco at La Antigua.  
Paratypes: 29. Same locality.  
GOBIIDAE
- Gonostoma ebelingi** Grey, 1960  
Bull. Mus. Comp. Zool. Harv. Coll., 122(2): 109.  
Paratype. Western Pacific Ocean: Marshall Islands area.  
CLUPEIDAE
- Goodea captiva** Hubbs, 1924  
Occas. Pap. Mus. Zool. Univ. Mich., 148: 4.  
Holotype: FMNH 5557. Mexico: San Luis Potosí, Jesús María.  
Paratypes: 172. Same locality.  
GOODEIDAE
- Goodea toweri** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 138.  
Holotype: FMNH 4519. Mexico: San Luis Potosí, Río Verde.  
Paratypes: 26. Same locality.  
GOODEIDAE
- Goodea whitei** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 137.  
Holotype: FMNH 4547. Mexico: Morelos, Yau-tepec.  
Paratypes: 420. Mexico: 2 localities.  
GOODEIDAE
- Grammonus mowbrayi** Grey, 1951  
Copeia, 1951(2): 154.  
Holotype: FMNH 49185. Western North Atlantic: off Bermuda.  
Paratypes: 4. Same locality.  
OPHIDIIDAE
- Grasseichthys gabonensis** Géry, 1964  
C. R. Seances Acad. Sci. Paris, 259: 4806.  
Paratypes: 6. Gabon: tributary of Ntsimy River near Nzingmeyong.  
KNERIIDAE
- Grimatrocetes bullisi** Grey, 1958  
Fieldiana Zool., 39(16): 150.  
Paratypes: 3. Gulf of Mexico.  
ALEPOCEPHALIDAE
- Gunnellichthys viridesens** Dawson, 1968  
Proc. Biol. Soc. Wash., 81: 61.  
Paratype. Maldive Islands: Addu Atoll, Bushy Island.  
MICRODESMIDAE
- Gymnapogon gracilicauda** Lachner, 1953  
Bull. U.S. Natl. Mus., 202(1): 497.  
Paratype. Marshall Islands: Bikini Atoll, Airi Island.  
APOGONIDAE
- Gymnapogon urospilotus** Lachner, 1953  
Bull. U.S. Natl. Mus., 202(1): 494.  
Paratypes: 2. Marshall Islands: 2 localities.  
APOGONIDAE
- Gymnorhamphichthys hypostomus** Ellis, 1912  
Mem. Carnegie Mus., 5: 436.  
Holotype: FMNH 54554 (CM 3182). Bolivia: San Joaquín.  
Paratypes: 4. Brazil: 3 localities.  
RHAMPHICHTHYIDAE
- Gymnothorax bayeri** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 124.  
Paratype. Marshall Islands: Bikini Atoll, Erik Island.  
MURAENIDAE
- Gymnothorax bikiniensis** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 116.  
Paratype. Marshall Islands: Bikini Atoll, Boro Island.  
MURAENIDAE
- Gymnothorax efatensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 384.  
Holotype: FMNH 17193. New Hebrides Islands: Efate Island, Vila.  
MURAENIDAE

**Gymnothorax leucostigma** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 174.  
Holotype: FMNH 52124 (CM 253). Taiwan: Takao.  
MURAENIDAE

**Hadropterus scierus apristis** Hubbs and Hubbs, 1954  
In: Hubbs, Am. Midl. Nat., 52(1): 211.  
Paratypes: 13. Texas: San Marcos River.  
PERCIDAE

**Halichoeres fijiensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 410.  
Holotype: FMNH 17341. Fiji Islands: Ovalau Island.  
Paratypes: 3. Same locality.  
LABRIDAE

**Hampala macrolepidota sabana** Inger and Chin, 1962  
Fieldiana Zool., 45: 81.  
Holotype: FMNH 68218. Malaysia: Sabah, tributary of Kinabatangan River near Deramakot.  
Paratypes: 134. Malaysia: Sabah, 5 localities.  
CYPRINIDAE

**Hasemania bilineata** Ellis, 1911  
Ann. Carnegie Mus., 8(2): 150.  
Holotype: FMNH 54383 (CM 3001). Brazil: São Paulo, Alto da Serra.  
Paratypes: 6. Brazil: Rio Tietê at Mogi das Cruces.  
CHARACIDAE

**Hasemania maxillaris** Ellis, 1911  
Ann. Carnegie Mus., 8(2): 148.  
Holotype: FMNH 54303 (CM 2937). Brazil: Rio Iguaçu at Porto União.  
CHARACIDAE

**Hasemania melanura** Ellis, 1911  
Ann. Carnegie Mus., 8(2): 149.  
Holotype: FMNH 54384 (CM 3002). Brazil: Rio Iguaçu at Porto União.  
Paratypes: 39. Same locality.  
CHARACIDAE

**Helogenes unidorsalis** Glodek and Carter, 1978  
Fieldiana Zool., 72(6): 77.  
Holotype: FMNH 80463. Ecuador: Río Bobonaza between Sarayacu and Montalvo.  
Paratypes: 5. Same locality.  
HELOGENIDAE

**Helotes lorentzi** Weber, 1910  
Notes Leiden Mus., 32: 236.  
Syntype: FMNH 52495 (CM 753). New Guinea: Lorentz River.  
TERAPONIDAE

**Hemiancistrus daguae** Eigenmann, 1912  
Indiana Univ. Stud., 16: 11.  
Holotype: FMNH 56052 (CM 4842). Colombia: Caldas.  
Paratypes: 11. Colombia: 2 localities.  
LORICARIIDAE

**Hemiancistrus mayoloi** Eigenmann, 1912  
Indiana Univ. Stud., 16: 10.  
Holotype: FMNH 56036 (CM 4826). Colombia: Istmina.  
Paratypes: 2. Same locality.  
LORICARIIDAE

**Hemiancistrus wilsoni** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 678.  
Holotype: FMNH 58550 (CM 7570). Colombia: Truandó.  
Paratypes: 3. Same locality.  
LORICARIIDAE

**Hemibrycon boquiae** Eigenmann, 1913  
Indiana Univ. Stud., 18: 20.  
Holotype: FMNH 56259 (CM 5059). Colombia: Boquia.  
Paratypes: 10. Same locality.  
Published originally as "*Bryconamericus* or *Hemibrycon*".  
CHARACIDAE

**Hemibrycon colombianus** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 8.  
Holotype: FMNH 56653 (CM 5470). Colombia: Río San Gil, Santander.  
Paratypes: 7. Colombia: 3 localities.  
CHARACIDAE

**Hemibrycon dariensis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 285.  
Holotype: FMNH 8947. Panama: Río Tuyra basin, Río Yape.  
Paratypes: 28. Panama: 7 localities.  
CHARACIDAE

**Hemibrycon helleri** Eigenmann, 1927  
Mem. Mus. Comp. Zool. Harv. Coll., 43(4): 406.  
Paratypes: 3. Peru: Río Comerciato.  
CHARACIDAE

**Hemicarax rhomboides** Meek and Hildebrand, 1925

Field Mus. Nat. Hist. Zool. Ser., 15(2): 345.  
Paratype. Panama: Fox Bay, Colón.  
CARANGIDAE

**Hemicetopsis amphioxus** Eigenmann, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 14.  
Holotype: FMNH 56519 (CM 5332). Colombia: Río Patía basin, creek near San Lorenzo.  
CETOPSIDAE

**Hemicetopsis macilentus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 211.  
Holotype: FMNH 53260 (CM 1726). Guyana: creek below Potaro Landing.  
CETOPSIDAE

**Hemicetopsis minutus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 211.  
Holotype: FMNH 53262 (CM 1728). Guyana: Amaturuk.  
CETOPSIDAE

**Hemicetopsis othonops** Eigenmann, 1912

Indiana Univ. Stud., 16: 17.  
Holotype: FMNH 56040 (CM 4830). Colombia: Girardot.  
Paratypes: 16. Colombia: 2 localities.  
CETOPSIDAE

**Hemiculterella engraulis** Nichols, 1925

Am. Mus. Nov., 182: 7.  
Paratypes: 2. China: Hunan, Tungting Lake.  
CYPRINIDAE

**Hemidoras leporhinus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 195.  
Holotype: FMNH 53185 (CM 1624). Guyana: Tumatumari.  
Paratype. Guyana: Crab Falls.  
DORADIDAE

**Hemidoras micropoeus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 195.  
Holotype: (CM 1636) MISSING. Guyana: Wismar.  
Paratype. Guyana: Lama Stop-off.  
DORADIDAE

**Hemidoras microstomus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 193.  
Holotype: FMNH 53206 (CM 1650). Guyana: Rockstone.  
Paratypes: 3. Guyana: Essequibo River, Crab Falls.  
DORADIDAE

**Hemidoras notospilus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 196.  
Holotype: FMNH 53184 (CM 1623). Guyana: Crab Falls.  
DORADIDAE

**Hemieleotris levis** Eigenmann, 1917

Proc. Am. Philos. Soc., 56(7): 684.  
Paratypes: 5. Colombia: pools in Buenaventura.  
ELEOTRIDAE

**Hemigrammus analis** Durbin, 1909

Ann. Carnegie Mus., 6(1): 64.  
Holotype: FMNH 53041 (CM 1466). Guyana: Rockstone.  
Paratypes: 45. Guyana: 3 localities.  
CHARACIDAE

**Hemigrammus brevis** Ellis, 1911

Ann. Carnegie Mus., 8(2): 161.  
Holotype: FMNH 54444 (CM 3068). Brazil: Lagoas of Rio Grande at Barreiras.  
Paratypes: 37. Brazil: 3 localities.  
CHARACIDAE

**Hemigrammus compressus** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 87.  
Holotype: FMNH 4641. Mexico: Río Papaloapan basin at El Hule.  
Paratypes: 28. Mexico: 2 localities.  
CHARACIDAE

**Hemigrammus cylindricus** Durbin, 1909

Ann. Carnegie Mus., 6(1): 62.  
Holotype: FMNH 53426 (CM 1461). Guyana: Tumatumari.  
Paratypes: 20. Guyana: 4 localities.  
CHARACIDAE

**Hemigrammus erythrozonus** Durbin, 1909

Ann. Carnegie Mus., 6(1): 56.  
Holotype: FMNH 53546 (CM 1448). Guyana: Erukin.  
Paratypes: 7. Same locality.  
CHARACIDAE

**Hemigrammus iota** Durbin, 1909

Ann. Carnegie Mus., 6(1): 60.  
Holotype: FMNH 53034 (CM 1458). Guyana: Gluck Island.  
Paratypes: 4. Guyana: 2 localities.  
CHARACIDAE

- Hemigrammus kennedyi** Eigenmann, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 520.  
Paratypes: 4. Paraguay: Campo Grande.  
CHARACIDAE
- Hemigrammus marginatus** Ellis, 1911  
Ann. Carnegie Mus., 8(2): 159.  
Holotype: FMNH 54430 (CM 3053). Brazil: Rio Itapicurú at Queimadas.  
Paratypes: 130. Brazil and Paraguay: 18 localities.  
CHARACIDAE
- Hemigrammus melanopterus** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 518.  
Paratypes: 2. Paraguay: Arroyo Pypucú.  
Spelled *melasopterus* in Eigenmann and Kennedy (1903), apparently a typographic error.  
CHARACIDAE
- Hemigrammus minutus** Meek and Hildebrand, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(6): 67.  
Holotype: FMNH 7572 MISSING or mixed.  
Panama: Canal Zone, Río Agua Clara.  
Paratypes: 71. Panama: 7 localities.  
CHARACIDAE
- Hemigrammus orthus** Durbin, 1909  
Ann. Carnegie Mus., 6(1): 61.  
Holotype: FMNH 53052 (CM 1477). Guyana: Tukeit.  
Paratypes: 19. Guyana: 3 localities.  
CHARACIDAE
- Hemigrammus riddlei** Meek, 1907  
In: Eigenmann and Ogle, Proc. U.S. Natl. Mus., 33(1556): 13.  
Holotype: MISSING. Venezuela: Los Castillos.  
See comment in Grey (1947, p. 156); this was published as at the Field Museum, but neither the catalogue entry nor the fish have been found.  
CHARACIDAE
- Hemigrammus rodwayi** Durbin, 1909  
Ann. Carnegie Mus., 6(1): 58.  
Holotype: FMNH 53026 (CM 1450). Guyana: Georgetown trenches.  
Paratypes: 248. Guyana: 4 localities.  
CHARACIDAE
- Hemiodus orthonops** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 511.  
Paratype. Paraguay: Asunción.  
HEMIODONTIDAE
- Hemiodus parnaguae** Eigenmann and Henn, 1916  
Ann. Carnegie Mus., 10(1–2): 87.  
Holotype: FMNH 56886 (CM 5701). Brazil: Lagoa de Parnagua.  
Paratypes: 2. Same locality.  
HEMIODONTIDAE
- Hemiodus quadrimaculatus** Pellegrin, 1908  
Bull. Mus. d'Hist. Nat., 14(7): 343.  
Paratype: MISSING. French Guiana: Camopi River.  
HEMIODONTIDAE
- Hemipimelodus papillifer** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 390.  
Holotype: FMNH 17211. New Guinea: Sepik River at Timbunke.  
Paratype. Same locality.  
ARIIDAE
- Hemipimelodus velutinus** Weber, 1908  
Nova Guinea, 5(2): 225.  
Syntype: FMNH 52392 (CM 648). New Guinea: Tawarin River.  
ARIIDAE
- Hemipteronotus baldwini** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 192.  
Paratypes: 2. Hawaiian Islands: Honolulu.  
LABRIDAE
- Hemipteronotus evides** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 196.  
Holotype: FMNH 52197 (CM 343). Taiwan: Takao.  
LABRIDAE
- Hemiramphus archipelagicus** Collette, 1978  
Proc. Biol. Soc. Wash., 91(3): 732.  
Paratypes: 2. Indonesia: Java, Batavia.  
HEMIRAMPHIDAE
- Hemiramphus bermudensis** Collette, 1962  
Bull. Mar. Sci. Gulf Caribb., 12(3): 432.  
Holotype: FMNH 48115. Bermuda: Saint George's Island, Whalebone Bay.  
Paratypes: 6. Bermuda: 6 localities.  
HEMIRAMPHIDAE

**Heptapterus stewarti** Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 376.

Holotype: FMNH 54234 (CM 2850). Brazil: creek at Serrinha Paraná.

PIMELODIDAE

**Heros beani** Jordan, 1889

Proc. U.S. Natl. Mus. 1888, 11: 332.

Paratype. Mexico: Río Presidio, Mazatlán.

CICHLIDAE

**Heterandria colombianus** Eigenmann and Henn, 1912

In: Eigenmann, Indiana Univ. Stud., 16: 27.

Holotype: FMNH 56047 (CM 4837). Colombia: mouth of Río Dagua.

Paratypes: 2. Same locality.

POECILIIDAE

**Heterandria cubensis** Eigenmann, 1903

Bull. U.S. Natl. Mus. 1902, 22: 227.

Paratype. Cuba: Pinar del Río.

POECILIIDAE

**Heterandria hasemani** Henn, 1916

Ann. Carnegie Mus., 10(1-2): 116.

Holotype: FMNH 55874 (CM 4663). Bolivia: Puerto Suárez.

Paratypes: 3. Same locality.

POECILIIDAE

**Heterandria lutzi** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 106.

Holotype: FMNH 3718. Mexico: Río Quioatepec at Cuicatlán.

Paratypes: 381. Mexico: 3 localities.

POECILIIDAE

**Heterenchelys biaggii** Böhlke, 1956

Not. Nat. (Phila.), 289: 8.

Holotype: FMNH 61774. Puerto Rico: Mayagüez pier.

Paratype. Same locality.

HETERENCHELYIDAE

**Heterocharax macrolepis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 406.

Holotype: FMNH 53669 (CM 2142). Guyana: Rockstone.

Paratypes: 2. Guyana: 2 localities.

CHARACIDAE

**Heterogramma corumbae** Eigenmann and Ward, 1907

Ann. Carnegie Mus., 4(2): 146.

Paratype. Paraguay: Corumbá.

CICHLIDAE

**Heterogramma ortmanni** Eigenmann, 1912

Mem. Carnegie Mus., 5: 506.

Holotype: FMNH 53801 (CM 2306). Guyana: Erukin.

Paratypes: 25. Guyana: 6 localities.

CICHLIDAE

**Heterogramma ritense** Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 362.

Holotype: FMNH 54187 (CM 2765). Brazil: Rio Santa Rita at Santa Rita.

Paratypes: 3. Same locality.

CICHLIDAE

**Heterogramma taeniatum pertense** Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 359.

Holotype: FMNH 54171 (CM 2741). Brazil: Manaus.

Paratype: MISSING. Brazil: Rio Tapajós at Santarém.

CICHLIDAE

**Heterogramma trifasciatum maciliense** Haseman, 1911

Ann. Carnegie Mus., 7(3-4): 360.

Syntypes: 2, FMNH 54177 (CM 2751) MISSING. Brazil: São Antonio de Guaporé.

CICHLIDAE

**Hildadoras boliviensis** Fernández-Yépez, 1968

Bol. Inst. Oceanogr. Univ. Oriente, 7(1): 43.

Holotype: FMNH 84067. Venezuela: Orinoco system, Quebrada El Pilón.

Paratype. Same locality.

DORADIDAE

**Hippocampus brunneus** Bean, 1906

Proc. Biol. Soc. Wash., 19: 32.

Holotype: FMNH 5494. Bermuda: Long Bird Island.

SYNGNATHIDAE

**Hippocampus fisheri** Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 169.

Paratype. Hawaiian Islands: Hilo.

SYNGNATHIDAE

**Holacanthus lunatus** Blosser, 1909

Ann. Carnegie Mus., 6(1): 299.

Holotype: FMNH 52831 (CM 1235). Lesser Antilles: Saint Croix Island.

Misspelled *Holocanthus* in Blosser (1909).

POMACANTHIDAE



- Holesthes heterodon** Eigenmann, 1915  
 Mem. Carnegie Mus., 7(1): 87.  
 Holotype: FMNH 57890 (CM 6875). Brazil: Jaguará.  
 Paratypes: 86. Brazil: 13 localities.  
 CHARACIDAE
- Holocentrus bullisi** Woods, 1955  
 Fieldiana Zool., 37(4): 113.  
 Paratypes: 16. Gulf of Mexico and North Atlantic: 6 localities.  
 HOLOCENTRIDAE
- Holocentrus ensifer** Jordan and Evermann, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 176.  
 Paratype. Hawaiian Islands: Honolulu.  
 HOLOCENTRIDAE
- Holocentrus meeki** Bean, 1906  
 Proc. Biol. Soc. Wash., 19: 31.  
 Lectotype: FMNH 5079. Bermuda: Saint David's Island, Hayward's Bay.  
 Lectoparatype. Same locality.  
 Lectotype designated by Grey (1947, p. 157).  
 HOLOCENTRIDAE
- Holocentrus xantherythrus** Jordan and Evermann, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 175.  
 Paratypes: 2. Hawaiian Islands: Honolulu.  
 HOLOCENTRIDAE
- Hololepis saludae** Hubbs and Cannon, 1935  
 Misc. Publ. Mus. Zool. Univ. Mich., 30: 50.  
 Paratype. South Carolina: Saluda County, Richland Creek.  
 PERCIDAE
- Holopristis riddlei** Meek, 1907  
 In: Eigenmann and Ogle, Proc. U.S. Natl. Mus., 33(1556): 11.  
 Paratypes: 3. Venezuela: Los Castillos.  
 See comment in Grey (1947, p. 158); misspelled *Holopristes* in Meek (1907).  
 CHARACIDAE
- Homaloptera hingi** Herre, 1934  
 Lingnan Sci. J. Canton, 13(2): 287.  
 Paratypes: 4. Hong Kong: stream at Pok Fu Lam.  
 HOMALOPTERIDAE
- Hoplomyzon papillatus** Stewart, 1985  
 Fieldiana Zool., New Ser., 25: 8.  
 Holotype: FMNH 94908. Ecuador: Río Aguarico, 1 km upstream from confluence with Río Shushufindi.  
 ASPREDINIDAE
- Hoplosternum magdalenae** Eigenmann, 1913  
 In: Ellis, Ann. Carnegie Mus., 8(3-4): 412.  
 Holotype: FMNH 56281 (CM 5081). Colombia: Soplaviento.  
 Paratypes: 12. Colombia: 2 localities.  
 CALLICHTHYIDAE
- Hoplosternum punctatum** Meek and Hildebrand, 1916  
 Field Mus. Nat. Hist. Zool. Ser., 10(15): 264.  
 Holotype: FMNH 8943. Panama: Río Marte. Armade east of Panama City.  
 Paratype. Same locality.  
 CALLICHTHYIDAE
- Hoplostethus occidentalis** Woods, 1973  
 Mem. Sears Fnd. Mar. Res., 1(6): 319.  
 Holotype: FMNH 66790. Gulf of Mexico.  
 Paratypes: 110. Tropical Western Atlantic: 37 localities.  
 TRACHICHTHYIDAE
- Hybognathus hankinsoni** Hubbs, 1929  
 In: Jordan, Manual of Vertebrate Animals of Northeast United States, p. 88.  
 Paratypes: 13. Iowa: 4 localities.  
 CYPRINIDAE
- Hybopsis x-punctata trautmani** Hubbs and Crowe, 1956  
 Occas. Pap. Mus. Zool. Univ. Mich., 578: 7.  
 Paratype. Indiana: Park County, Sugar Creek.  
 CYPRINIDAE
- Hybopsis x-punctata x-punctata** Hubbs and Crowe, 1956  
 Occas. Pap. Mus. Zool. Univ. Mich., 578: 7.  
 Paratypes: 2. Missouri: Arlington.  
 Two additional specimens with localities "Illinois" and "no data" were also labeled paratypes of *H. x-punctata* by Hubbs, but the subspecies designations were not given.  
 CYPRINIDAE
- Hymenocephalus aterrimus** Gilbert, 1905  
 Bull. U.S. Fish Comm. 1903, 23(2): 666.  
 Paratype. Hawaiian Islands.  
 MACROURIDAE
- Hymenocephalus grimaldii** Weber, 1913  
 Die Fische der Siboga-Expedition, Leiden, p. 169.  
 Syntype: FMNH 52450 (CM 708). Indonesia: Bali Sea.  
 MACROURIDAE

- Hyphessobrycon bifasciatus* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 156.  
 Holotype: FMNH 54404 (CM 3026). Brazil: Rio Parahyba at Campos.  
 Paratypes: 201. Brazil: 8 localities.  
 CHARACIDAE
- Hyphessobrycon duragenys* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 155.  
 Holotype: (CM 3023) MISSING. Brazil: Rio Tietê at Mogy das Cruces.  
 Paratypes: 14. Brazil: 2 localities.  
 CHARACIDAE
- Hyphessobrycon ecuadoriensis* Eigenmann and Henn, 1914**  
 In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 9.  
 Paratypes: 107. Ecuador: forest pool near Vines.  
 CHARACIDAE
- Hyphessobrycon eos* Durbin, 1909**  
 Ann. Carnegie Mus., 6(1): 69.  
 Holotype: FMNH 52795 (CM 1194). Guyana: creek between Potaro Landing and Kangaruma.  
 Paratypes: 23. Guyana: 2 localities.  
 CHARACIDAE
- Hyphessobrycon melanopleurus* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 157.  
 Holotype: FMNH 54413 (CM 3035). Brazil: São Paulo, Alto da Serra.  
 Paratypes: 2. Same locality.  
 CHARACIDAE
- Hyphessobrycon minimus* Durbin, 1909**  
 Ann. Carnegie Mus., 6(1): 68.  
 Holotype: FMNH 52794 (CM 1193). Guyana: Cane Grove Corner.  
 CHARACIDAE
- Hyphessobrycon minor* Durbin, 1909**  
 Ann. Carnegie Mus., 6(1): 65.  
 Holotype: FMNH 52789 (CM 1189) MISSING.  
 Guyana: Konawaruk.  
 CHARACIDAE
- Hyphessobrycon panamensis daguae* Eigenmann, 1922**  
 Mem. Carnegie Mus., 9(1): 141.  
 Syntypes: 9, FMNH 56526–56529 and 58316.  
 Colombia: Río Patía basin.  
 CHARACIDAE
- Hyphessobrycon parvulus* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 153.  
 Holotype: FMNH 54392 (CM 3011). Brazil: Rio Catu at Alagoinhas.  
 Paratypes: 13. Brazil: 4 localities.  
 CHARACIDAE
- Hyphessobrycon poecilioides* Eigenmann, 1913**  
 Indiana Univ. Stud., 18: 29.  
 Holotype: FMNH 56290 (CM 5091). Colombia: Cali.  
 Paratypes: 36. Same locality.  
 CHARACIDAE
- Hyphessobrycon proteus* Eigenmann, 1913**  
 Indiana Univ. Stud., 18: 28.  
 Holotype: FMNH 56293 (CM 5094). Colombia: Quibdó.  
 Paratypes: 198. Colombia: 8 localities.  
 CHARACIDAE
- Hyphessobrycon reticulatus* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 153.  
 Holotype: FMNH 54397 (CM 3018). Brazil: Rio Parahyba at Campos.  
 Paratypes: 17. Brazil: 6 localities.  
 CHARACIDAE
- Hyphessobrycon rosaceus* Durbin, 1909**  
 Ann. Carnegie Mus., 6(1): 67.  
 Holotype: FMNH 52791 (CM 1190). Guyana: Glück Island.  
 Paratypes: 10. Guyana: 2 localities.  
 CHARACIDAE
- Hyphessobrycon stictus* Durbin, 1909**  
 Ann. Carnegie Mus., 6(1): 71.  
 Holotype: FMNH 52798 (CM 1197). Guyana: Lama Stop-off.  
 Paratypes: 146. Guyana: 5 localities.  
 CHARACIDAE
- Hyphessobrycon taurocephalus* Ellis, 1911**  
 Ann. Carnegie Mus., 8(2): 151.  
 Holotype: FMNH 54389 (CM 3007). Brazil: Rio Iguaçu at Serrinha Paraná.  
 Paratypes: 59. Brazil: 2 localities.  
 CHARACIDAE
- Hypoatherina barnesi* Schultz, 1953**  
 Bull. U.S. Natl. Mus., 202(1): 304.  
 Paratypes: 4. Marshall Islands: 2 localities.  
 ATHERINIDAE

**Hyporhamphus hildebrandi** Jordan and Evermann, 1927

Proc. Calif. Acad. Sci., 16(15): 503.

Lectotype: FMNH 8260. Panama: Colón, Fox Bay at Toro Point.

Lectoparatypes: 6. Same locality.

Lectotype designated by Weed (1933, Field Mus. Nat. Hist. Zool. Ser., 20: 60).

HEMIRAMPHIDAE

**Hyporhamphus melanopterus** Collette and Parin, 1978

Proc. Biol. Soc. Wash., 91(3): 738.

Paratypes: 2. Malaysia: Sabah, Sandakan, fish market.

HEMIRAMPHIDAE

**Hyporhamphus snyderi** Meek and Hildebrand, 1923

Field Mus. Nat. Hist. Zool. Ser., 15(1): 240.

Paratypes: 2. Panama: Chame Point.

HEMIRAMPHIDAE

**Hypsoblennius lignus** Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 946.

Paratype. Panama: Chame Point.

BLENNIIDAE

**Hysteronotus megalostomus** Eigenmann, 1911

Ann. Carnegie Mus., 8(1): 171.

Holotype: FMNH 54889 (CM 3551). Brazil: Rio das Velhas.

Paratypes: 2. Same locality.

CHARACIDAE

**Iburiella kasawae** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 291.

Holotype: FMNH 58796 (CM 7906). Japan: Tomakomai, near Mororan.

AGONIDAE

**Ichthyomyzon unicuspis** Hubbs and Trautman, 1937

Misc. Publ. Mus. Zool. Univ. Mich., 35: 53.

Paratypes: 8. Illinois and Ohio: 3 localities.

PETROMYZONTIDAE

**Icticus ischanus** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 242.

Holotype: FMNH 57097 (CM 6036). Japan: Ryukyu Islands, Okinawa.

NOMEIDAE

**Ilisha argentata** Meek and Hildebrand, 1923

Field Mus. Nat. Hist. Zool. Ser., 15(1): 190.

Paratypes: 15. Panama: 5 localities.

CLUPEIDAE

**Imparfinis hollandi** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 383.

Holotype: FMNH 54244 (CM 2864). Brazil: Rio Iguaçu at Pôrto União da Victoria.

Haseman stated “Type unique”, but lot CM 2864 included 3 specimens. The 2 topotypes are now FMNH 54245.

PIMELODIDAE

**Imparfinis microps** Eigenmann and Fisher, 1916

In: Eigenmann, Ann. Carnegie Mus., 10(1–2): 82.

Holotype: FMNH 57793 (CM 6776). Colombia: Río Negro at Villavicencio.

Paratypes (?): 5. Same locality.

Published number CM 8778 is an error. Original description was based on “type only”; but 5 paratypes were listed in Eigenmann (1922, Mem. Carnegie Mus., 9(1): 221).

PIMELODIDAE

**Imparfinis mirini** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 318.

Holotype: FMNH 54335 (CM 2981). Brazil: Rio Piracicaba-mirini near Piracicaba.

Paratypes: 6. Same locality.

PIMELODIDAE

**Iridio bimaculata** Wilson, 1916

Ann. Carnegie Mus., 10(1–2): 68.

Holotype: FMNH 56466 (CM 5280). Colombia: Río Dagua at Buenaventura.

Paratype. Same locality.

LABRIDAE

**Iridio decoratus** Bean, 1906

Proc. Biol. Soc. Wash., 19: 29.

Lectotype: FMNH 5128. Bermuda: Nonsuch Island.

Lectoparatype. Same locality.

Lectotype designated by Grey (1947, p. 160).

LABRIDAE

**Iridio elegans** Bean, 1906

Proc. Biol. Soc. Wash., 19: 30.

Lectotype: FMNH 5129. Bermuda: Nonsuch Island, Castle Harbor.

Lectoparatypes: 3. Bermuda: Cooper's Island, Jones Bay.

Lectotype designated by Grey (1947, p. 160).

LABRIDAE

- Iridio meyeri** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 29.  
Holotype: FMNH 4938, Bermuda: Nonsuch Island.  
Published number FMNH 5496 is an error.  
LABRIDAE
- Iridio microstomus** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 30.  
Holotype: FMNH 5138, Bermuda: Nonsuch Island.  
LABRIDAE
- Ischnomembras gabunensis** Fowler, 1903  
Proc. Acad. Nat. Sci. Phila., 55: 731.  
Paratype. Gabon: Gabon River.  
Locality probably wrong since this is synonym of *Menidia beryllina*, a New World taxon; see remarks in Böhlke (1984, Spec. Publ. Acad. Nat. Sci. Phila., 14: 23).  
ATHERINIDAE
- Jenkinsia majua majua** Whitehead, 1963  
Bull. Brit. Mus. (Nat. Hist.) Zool., 10(6): 356.  
Paratypes: 54. Gulf of Mexico: Campeche Bank.  
CLUPEIDAE
- Jenkinsia majua woodsi** Whitehead, 1963  
Bull. Brit. Mus. (Nat. Hist.) Zool., 10(6): 356.  
Paratypes: 10. Caribbean Sea: off Quita Sueño Bank.  
CLUPEIDAE
- Joturus daguae** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 681.  
Holotype: FMNH 58457 (CM 7458). Colombia: Río Dagua at Caldas.  
MUGILIDAE
- Kaupichthys atlanticus** Böhlke, 1956  
Proc. Acad. Nat. Sci. Phila., 108: 66.  
Paratype. Bermuda: Castle Harbor.  
XENOCOGRIDAE
- Kaupichthys brachychirus** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 67.  
Paratype. Marshall Islands: Bikini Atoll, Romuk Island.  
XENOCOGRIDAE
- Kryptopterus parvanalis** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 284.  
Holotype: FMNH 68014, Malaysia: Sabah, Kinabatangan River at Deramakot.  
Paratypes: 25. Malaysia: Sabah, 14 localities.  
SILURIDAE
- Kuhlia bilunulata** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 404.  
Holotype: FMNH 17313, Fiji Islands: Viti Levu Island, Suva Harbor.  
Paratypes: 9. Same locality.  
KUHLIIDAE
- Labeo fisheri** Jordan and Starks, 1917  
Ann. Carnegie Mus., 11: 436.  
Holotype: FMNH 58911 (CM 8044). Sri Lanka: Mahaweli River at Kandy.  
CYPRINIDAE
- Labeo jordani** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 204.  
Holotype: FMNH 59089 (CM 8226). Taiwan: hatchery at Shori.  
CYPRINIDAE
- Labrisomus lentiginosus** Bean, 1906  
Proc. Biol. Soc. Wash., 19: 30.  
Holotype: FMNH 5142, Bermuda: Paget Parish, tide pool at Doe's Rock.  
LABRISOMIDAE
- Labrisomus multiporosus** Hubbs, 1953  
Zoologica (N.Y.), 38(9): 131.  
Paratypes: 2. California: 2 localities.  
LABRISOMIDAE
- Lampanyctus jordani** Gilbert, 1913  
Mem. Carnegie Mus., 6(2): 104.  
Holotype: FMNH 55837 (CM 4617). Japan: Hokkaido, Nemuro.  
MYCTOPHIDAE
- Lampanyctus micropunctatus** Chapman, 1939  
Proc. U.S. Natl. Mus., 86(3062): 527.  
Paratype. North Pacific: off Washington.  
MYCTOPHIDAE
- Lamprologus obliquus** Nichols and Griscom, 1917  
Bull. Am. Mus. Nat. Hist., 37(25): 731.  
Paratypes: 3. Zaire: Stanleyville.  
CICHLIDAE
- Landonia latidens** Eigenmann and Henn, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 2.  
Paratypes: 2. Ecuador: Vinces.  
CHARACIDAE

**Lasiancistrus brevispinis** Heitmans, Nijssen, and Isbrucker, 1983  
Bijdr. Dierkd., 53(1): 38.  
Paratypes: 2. Surinam: Fallawatra River at Stondansie Fall.  
LORICARIIDAE

**Lasiancistrus caucanus** Eigenmann, 1912  
Indiana Univ. Stud., 16: 11.  
Holotype: FMNH 56034 (CM 4824). Colombia: Cartago.  
Paratypes: 4. Same locality.  
LORICARIIDAE

**Leiocassis robustus** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 290.  
Holotype: FMNH 68001. Malaysia: Sabah, Kinabatangan River at Deramakot.  
Paratypes: 5. Malaysia: Sabah, 2 localities.  
BAGRIDAE

**Leiogaster melanopus** Weber, 1913  
Die Fische der Siboga-Expedition, Leiden, p. 180.  
Syntype: FMNH 52454 (CM 712). Indonesia: Ceram Sea.  
BAGRIDAE

**Lepidaplois strophodes** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 190.  
Paratype. Hawaiian Islands: Honolulu.  
Misspelled *Lepidoplois* in Jordan and Evermann (1903).  
LABRIDAE

**Lepidomeda mollispinis mollispinis** Miller and Hubbs, 1960  
Misc. Publ. Mus. Zool. Univ. Mich., 115: 18.  
Paratypes: 12. Nevada: Clark County, Virgin River.  
CYPRINIDAE

**Lepomis occidentalis** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 118.  
Holotype: FMNH 3560. Mexico: Río Conchos at Jiménez.  
Paratypes: 28. Mexico: 2 localities.  
CENTRARCHIDAE

**Lepophidium aporrhox** Robins, 1961  
Copeia, 1961(2): 218.  
Paratypes: 4. Atlantic Ocean: off Guyana and Surinam.  
OPHIDIIDAE

**Lepophidium pheromystax** Robins, 1960  
Bull. Mar. Sci. Gulf Caribb., 10(1): 84.  
Paratypes: 19. Western Atlantic Ocean: 6 localities.  
OPHIDIIDAE

**Lepophidium staurophor** Robins, 1958  
Bull. Mar. Sci. Gulf Caribb., 8(4): 360.  
Holotype: FMNH 64383. Western Caribbean Sea.  
Paratype. Caribbean Sea.  
OPHIDIIDAE

**Leporinodus retropinnis** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 116.  
Holotype: FMNH 55174 (CM 3875). Brazil: Piracicaba.  
ANOSTOMIDAE

**Leporinus alternus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 307.  
Holotype: FMNH 53361 (CM 1827). Guyana: Tukeit.  
Paratypes: 5. Guyana: 2 localities.  
ANOSTOMIDAE

**Leporinus arcus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 300.  
Holotype: FMNH 53366 (CM 1832). Guyana: Tukeit.  
Paratypes: 4. Guyana: 4 localities.  
ANOSTOMIDAE

**Leporinus brunneus** Myers, 1950  
Stanford Ichthyol. Bull., 3(4): 195.  
Paratype. Venezuela: Casiquiare, Chemoni, near bifurcation of Río Orinoco.  
ANOSTOMIDAE

**Leporinus ecuadoriensis** Eigenmann and Henn, 1916  
Ann. Carnegie Mus., 10(1–2): 88.  
Paratypes: 2. Ecuador: Río Barranca Alta near Naranjito.  
ANOSTOMIDAE

**Leporinus granti** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 307.  
Holotype: FMNH 53383 (CM 1851). Guyana: Maripicru Creek.  
Paratypes: 2. Same locality.  
ANOSTOMIDAE

- Leporinus steyermarki** Inger, 1956  
Fieldiana Zool., 34(37): 431.  
Holotype: FMNH 45701. Venezuela: Río Abácapa, west side of Chimantá-tepui.  
ANOSTOMIDAE
- Leptocephalus ectenurus** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 171.  
Holotype: FMNH 52117 (CM 245). Taiwan: Takao.  
CONGRIDAE
- Leptocephalus harringtonensis** Mowbray, 1931  
Fauna Bermudensis, 1: 1.  
Holotype: FMNH 48442. Bermuda: Harrington Sound.  
CONGRIDAE
- Leptocottus armatus australis** Hubbs, 1921  
Occas. Pap. Mus. Zool. Univ. Mich., 94: 5.  
Paratypes: 62. California: 10 localities.  
COTTIDAE
- Leptodoras linnelli** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 191.  
Holotype: FMNH 53561 (CM 1626). Guyana: Tumatumari.  
Paratypes: 16. Guyana: 4 localities.  
DORADIDAE
- Leptoglanis essequibensis** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 158.  
Holotype: FMNH 53331 (CM 1652). Guyana: Crab Falls.  
Paratypes: 3. Guyana: 2 localities.  
PIMELODIDAE
- Leptophilypnus fluviatilis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(5): 361.  
Holotype: FMNH 8952. Panama: Canal Zone, creek at Mindi.  
Paratypes: 20. Same locality.  
ELEOTRIDAE
- Leucichthys alpenae** Koelz, 1924  
Occas. Pap. Mus. Zool. Univ. Mich., 146: 1.  
Paratypes: 2. Michigan: 2 localities.  
SALMONIDAE
- Leucichthys kiya** Koelz, 1921  
Occas. Pap. Mus. Zool. Univ. Mich., 104: 1.  
Paratypes: 2. Michigan: Lake Michigan at Ludington.  
SALMONIDAE
- Leucichthys reighardi** Koelz, 1924  
Occas. Pap. Mus. Zool. Univ. Mich., 146: 5.  
Paratypes: 2. Wisconsin and Indiana: 2 localities.  
SALMONIDAE
- Leuciscus carletoni** Kendall, 1903  
Bull. U.S. Fish Comm. 1902, 22: 357.  
Paratypes: 9. Maine: East Branch, Penobscot River.  
CYPRINIDAE
- Leucogobio polytaenia** Nichols, 1925  
Am. Mus. Nov., 181: 6.  
Paratypes: 3. China: Shansi, Niang-tze-kwan.  
CYPRINIDAE
- Limia hollandi** Henn, 1916  
Ann. Carnegie Mus., 10(1-2): 138.  
Holotype: FMNH 55861 (CM 4643). Brazil: Rio São Francisco at Penedo.  
Paratypes: 56. Brazil: 11 localities.  
POECILIIDAE
- Limia rivasi** Franz and Burgess, 1983  
Northeast Gulf Sci., 6(1): 51.  
Paratypes: 10. Haiti: Ile de la Gonâve.  
POECILIIDAE
- Liobagrus anguillicauda** Nichols, 1926  
Am. Mus. Nov., 224: 1.  
Paratypes: 2. China: Fukien, Chungan Hsien.  
BAGRIDAE
- Liobagrus nantoensis** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 183.  
Holotype: FMNH 59080 (CM 8217). Taiwan: Dainansho, Nanto.  
BAGRIDAE
- Liopempheris sasakii** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 228.  
Holotype: FMNH 58751 (CM 7860). Japan: Toba market.  
Published number CM 7861 is an error.  
PEMPHERIDAE
- Liopropoma mowbrayi** Woods and Kanazawa, 1951  
Fieldiana Zool., 31(53): 633.  
Holotype: FMNH 48544. Bermuda: south shore.  
SERRANIDAE

- Lithogenes villosus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 6.  
Holotype: FMNH 52960 (CM 1002). Guyana:  
Potaro River at Aruataima Falls.  
LORICARIIDAE
- Lithoxus lithoides** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 242.  
Holotype: FMNH 53557 (CM 1527). Guyana:  
Amatuk.  
Paratypes: 122. Guyana: 6 localities.  
LORICARIIDAE
- Liza formosae** Oshima, 1922  
Ann. Carnegie Mus., 13(3–4): 251.  
Holotype: FMNH 59145 (CM 8283). Taiwan:  
Anpin.  
MUGILIDAE
- Liza parva** Oshima, 1922  
Ann. Carnegie Mus., 13(3–4): 253.  
Holotype: FMNH 59146 (CM 8284). Taiwan:  
Anpin.  
MUGILIDAE
- Liza pescadorensis** Oshima, 1922  
Ann. Carnegie Mus., 13(3–4): 254.  
Holotype: FMNH 59147 (CM 8285). Taiwan:  
Pescadores Island at Bako.  
MUGILIDAE
- Lonchopisthus higmani** Mead, 1959  
Stud. Fauna Suriname Other Guyanas, 2(5):  
104.  
Paratype. Western Atlantic Ocean: off Surinam.  
OPISTOGNATHIDAE
- Lophiobagrus aquilus** Bailey and Stewart, 1984  
Misc. Publ. Mus. Zool. Univ. Mich., 168: 24.  
Paratypes: 5. Zambia: Lake Tanganyika.  
BAGRIDAE
- Lophiobagrus brevispinis** Bailey and Stewart, 1984  
Misc. Publ. Mus. Zool. Univ. Mich., 168: 35.  
Paratypes: 4. Zambia: Lake Tanganyika.  
BAGRIDAE
- Lophiodes beroe** Caruso, 1981  
Copeia, 1981(3): 542.  
Paratypes: 2. Western North Atlantic: 2 localities.  
LOPHIIDAE
- Lophiodes reticulatus** Caruso and Suttkus, 1979  
Bull. Mar. Sci., 29(4): 491.  
Paratypes: 7. Gulf of Mexico.  
LOPHIIDAE
- Loricaria capetensis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 80.  
Holotype: FMNH 7582. Panama: Río Capetí.  
LORICARIIDAE
- Loricaria filamentosa latiura** Eigenmann and Vance, 1912  
In: Eigenmann, Indiana Univ. Stud., 16: 13.  
Syntypes: 7, FMNH 55115 (CM 3806). Colombia:  
Boca de Certegui.  
LORICARIIDAE
- Loricaria filamentosa seminuda** Eigenmann and Vance, 1912  
In: Eigenmann, Indiana Univ. Stud., 16: 13.  
Holotype: FMNH 55116 (CM 3807). Colombia:  
Girardot.  
LORICARIIDAE
- Loricaria fimbriata** Eigenmann and Vance, 1912  
In: Eigenmann, Indiana Univ. Stud., 16: 12.  
Holotype: FMNH 55117 (CM 3808). Colombia:  
Boca de Certegui.  
Paratype. Same locality.  
LORICARIIDAE
- Loricaria griseus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 8.  
Holotype: FMNH 53077 (CM 1504). Guyana:  
Konawaruk.  
Paratypes: 13. Guyana: 2 localities.  
LORICARIIDAE
- Loricaria gymnogaster** Eigenmann and Vance, 1912  
In: Eigenmann, Indiana Univ. Stud., 16: 12.  
Paratypes: 29. Colombia: Girardot.  
LORICARIIDAE
- Loricaria gymnogaster lagoichthys** Schultz, 1944  
Proc. U.S. Natl. Mus., 94(3172): 331.  
Paratypes: 5. Venezuela: Río Motatán north of  
Motatán.  
LORICARIIDAE
- Loricaria microdon** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 7.  
Holotype: FMNH 53555 (CM 1507). Guyana:  
Rupununi.  
LORICARIIDAE

- Loricaria stewarti** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 9.  
Holotype: FMNH 53330 (CM 1508). Guyana:  
Ireng River system, Chipoo Creek.  
Paratypes: 4. Same locality.  
LORICARIIDAE
- Loricaria submarginatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 10.  
Holotype: FMNH 53080 (CM 1510). Guyana:  
creek below Potaro Landing.  
LORICARIIDAE
- Loricaria tuyrensis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 81.  
Holotype: FMNH 7583. Panama: Río Tuyra,  
Boca de Cupe.  
Paratypes: 7. Same locality.  
LORICARIIDAE
- Loricaria uracantha rupestre** Schultz, 1944  
Proc. U.S. Natl. Mus., 94(3172): 325.  
Paratypes: 5. Venezuela: Río San Juan south of  
Mene Grande.  
LORICARIIDAE
- Loricaria variegata venezuelae** Schultz, 1944  
Proc. U.S. Natl. Mus., 94(3172): 329.  
Paratypes: 5. Venezuela: Río Sucuy, above its  
mouth.  
LORICARIIDAE
- Luciobrotula corethromycter** Cohen, 1954  
Bull. Mar. Sci. Gulf Caribb., 14(3): 391.  
Paratype. Gulf of Mexico.  
OPHIDIIDAE
- Lumpenus longirostris** Evermann and Goldsborough, 1907  
Bull. Bur. Fish. 1906, 26: 340.  
Paratypes: 2. Alaska: Albatross Station 4252.  
STICHAETIDAE
- Lycodes tanakae** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 299.  
Holotype: FMNH 57068 (CM 6004). Japan:  
Hondo, Noto.  
ZOARCIDAE
- Macgregorella badia** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 415.  
Holotype: FMNH 17373. Fiji Islands: Ovalau  
Island.  
GOBIIDAE
- Macgregorella santa** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 416.  
Holotype: FMNH 17374. New Hebrides Is-  
lands: Espiritu Santo Island, Hog Harbor.  
Paratypes: 7. New Hebrides Islands: 2 localities.  
GOBIIDAE
- Macrodonogobius wilburi** Herre, 1936  
Philipp. J. Sci., 59: 279.  
Paratype (?). Palau Islands.  
GOBIIDAE
- Macropodus filamentosus** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 278.  
Holotype: FMNH 59123 (CM 8261). Taiwan:  
Kotosho (Botel Tobago Island).  
BELONTIIDAE
- Macropsobrycon uruguayanae** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 48.  
Holotype: FMNH 57910 (CM 6895). Brazil:  
Cacequy.  
Paratypes: 5. Brazil: 2 localities.  
CHARACIDAE
- Macrostoma quercinum japonicum** Tanaka, 1908  
J. Coll. Sci. Imp. Univ. Tokyo, 23(13): 5.  
Holotype: FMNH 80459. Japan: Sagami Sea.  
MYCTOPHIDAE
- Macrurus richardi** Weber, 1913  
Die Fische der Siboga-Expedition, Leiden, p.  
154.  
Syntype: FMNH 52442 (CM 700). Indonesia:  
Flores Sea.  
MACROURIDAE
- Malacoctenus afueriae multipunctatus** Springer, 1958  
Inst. Mar. Sci., 5: 462.  
Paratypes: 52. Mexico: Golfo de California, Ba-  
hía Santa Inez.  
LABRISOMIDAE
- Malacoctenus ebisui** Springer, 1958  
Inst. Mar. Sci., 5: 478.  
Paratype. Costa Rica: Golfo de Nicoya, Isla Tor-  
tuga.  
LABRISOMIDAE
- Malacoctenus gigas** Springer, 1958  
Inst. Mar. Sci., 5: 457.  
Paratypes: 48. Mexico: Golfo de California, Isla  
Tiburón.  
LABRISOMIDAE



- Malaccoctenus hubbsi hubbsi** Springer, 1958  
Inst. Mar. Sci., 5: 465.  
Paratypes: 125. Mexico: Guaymas, 2 localities.  
LABRISOMIDAE
- Malaccoctenus hubbsi polyporosus** Springer, 1958  
Inst. Mar. Sci., 5: 468.  
Paratypes: 32. Mexico: Golfo de California, Isla Pájaro.  
LABRISOMIDAE
- Malaccoctenus margaritae mexicanus** Springer, 1958  
Inst. Mar. Sci., 5: 449.  
Paratypes: 47. Mexico: Golfo de California, Bahía Santa Inez.  
LABRISOMIDAE
- Malaccoctenus puertoricensis** Evermann and Marsh, 1900  
Rept. U.S. Fish Comm. 1899, 25: 358.  
Paratypes: 2. Puerto Rico: Culebra.  
LABRISOMIDAE
- Malaccoctenus triangulatus** Springer, 1958  
Inst. Mar. Sci., 5: 477.  
Paratypes: 8. Gulf of Mexico: 2 localities.  
LABRISOMIDAE
- Malaccoctenus zacae** Springer, 1958  
Inst. Mar. Sci., 5: 470.  
Paratypes: 12. Mexico: Golfo de California, Bahía San Lucas.  
LABRISOMIDAE
- Malacochthys wakiyai** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 233.  
Holotype: FMNH 58754 (CM 7863). Japan: Kagoshima Bay.  
PERCICHTHYIDAE
- Mastacembelus keithi** Herre, 1940  
Bull. Raffles Mus., 16: 24.  
Paratype. Malaysia: Sabah, Kabili River.  
MASTACEMBELIDAE
- Mastacembelus oatesii** Boulenger, 1893  
Ann. Mag. Nat. Hist., Ser. 6, 12: 199.  
Syntype: FMNH 5707. Burma: Fort Stedman.  
MASTACEMBELIDAE
- Megalamphodus megalopterus** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 50.  
Holotype: FMNH 57823 (CM 6806). Brazil: Cáceres.  
Paratypes: 2. Same locality.  
CHARACIDAE
- Megalamphodus micropterus** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 54.  
Holotype: FMNH 57916 (CM 6900). Brazil: Rio São Francisco at Lagoa do Porto.  
Paratypes: 54. Brazil: 4 localities.  
CHARACIDAE
- Megalonema platycephalum** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 150.  
Holotype: FMNH 53224 (CM 1684). Guyana: Tumatumari.  
Paratypes: 2. Guyana: Crab Falls.  
PIMELODIDAE
- Megalonema punctatum** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 77.  
Holotype: FMNH 7577. Panama: Río Tuyra, Boca de Cupe.  
PIMELODIDAE
- Megalonema robustum** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 78.  
Holotype: FMNH 7578. Panama: Río Tuyra, Marriganti.  
Paratypes: 3. Panama: 2 localities.  
PIMELODIDAE
- Megalonema xanthum** Eigenmann, 1912  
Indiana Univ. Stud., 16: 16.  
Holotype: FMNH 56032 (CM 4822). Colombia: Girardot.  
Paratypes: 90. Same locality.  
PIMELODIDAE
- Melamphaes pumilus** Ebeling, 1962  
Dana-Rep. Carlsberg Found. 58: 106.  
Paratypes: 31. Western North Atlantic: 6 localities.  
MELAMPHAIDAE
- Melaniris balsanus** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 117.  
Holotype: FMNH 3706. Mexico: Río Balsas at Balsas.  
Paratypes: 99. Same locality.  
ATHERINIDAE

- Melaniris sardina** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 114.  
Holotype: FMNH 5937. Nicaragua: Lago Managua at Managua.  
Paratypes: 309. Nicaragua: 3 localities.  
ATHERINIDAE
- Melanotaenia kabia** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 397.  
Holotype: FMNH 17230. New Guinea: Sepik River at Nyaurangai.  
Paratypes: 14. New Guinea: 2 localities.  
MELANOTAENIIDAE
- Melanotaenia rosacea** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 398.  
Holotype: FMNH 17245. New Guinea: Sepik River at Koragu.  
Paratypes: 19. New Guinea: 2 localities.  
MELANOTAENIIDAE
- Menidia chagresi** Meek and Hildebrand, 1914  
Field Mus. Nat. Hist. Zool. Ser., 10(10): 119.  
Paratypes: 59. Panama and Costa Rica: 3 localities.  
ATHERINIDAE
- Menidia lisa** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 182.  
Holotype: FMNH 4633. Mexico: Veracruz, Refugio.  
Paratypes: 48. Mexico: 2 localities.  
ATHERINIDAE
- Menidia peninsulae atrimentis** Kendall, 1902  
Rept. U.S. Fish Comm. 1901, 27: 258.  
Paratypes: 5. Florida: Lake Monroe.  
ATHERINIDAE
- Menidia starksi** Meek and Hildebrand, 1923  
Field Mus. Nat. Hist. Zool. Ser., 15(1): 267.  
Paratypes: 4. Panama: Taboga Island.  
ATHERINIDAE
- Menticirrhus focaliger** Ginsburg, 1952  
J. Wash. Acad. Sci., 42(3): 97.  
Paratypes: 2. Florida: Cape San Blas.  
SCIAENIDAE
- Metynnis guaporensis** Eigenmann, 1915  
Ann. Carnegie Mus., 9(3-4): 267.  
Holotype: FMNH 56913 (CM 5729). Brazil: Rio Guaporé at Maciel.  
Paratypes: 2. Brazil and Bolivia: 2 localities.  
CHARACIDAE
- Metynnis mola** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 528.  
Paratype. Paraguay: Arroyo Trementina.  
CHARACIDAE
- Metynnis roosevelti** Eigenmann, 1915  
Ann. Carnegie Mus., 9(3-4): 268.  
Holotype: FMNH 56925 (CM 5740). Brazil: Manaus.  
Paratypes: 8. Brazil: 3 localities.  
CHARACIDAE
- Microbrycon minutus** Eigenmann and Wilson, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 3.  
Holotype: FMNH 56606 (CM 5422). Colombia: Río Truando.  
CHARACIDAE
- Microdesmus carri** Gilbert, 1966  
Copeia, 1966(2): 326.  
Paratypes: 2. Costa Rica: Limón Province near mouth of Tortuguero.  
MICRODESMIDAE
- Microdesmus woodsi** Kanazawa, 1952  
Fieldiana Zool., 34(7): 93.  
Holotype: FMNH 48876. Bermuda: Saint David's Island.  
MICRODESMIDAE
- Microeleotris mindii** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 364.  
Holotype: FMNH 8954. Panama: Canal Zone, creek at Mindi.  
Paratypes: 25. Panama: 2 localities.  
ELEOTRIDAE
- Microeleotris panamensis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 363.  
Holotype: FMNH 8953. Panama: Río Juan Díaz.  
Paratypes: 18. Panama: 2 localities.  
ELEOTRIDAE
- Microgenys minutus** Eigenmann, 1913  
Indiana Univ. Stud., 18: 22.  
Holotype: FMNH 56215 (CM 5007). Colombia: Piedra de Moler.  
Paratype. Same locality.  
CHARACIDAE

- Microglanis iheringi** Gomes, 1946  
Occas. Pap. Mus. Zool. Univ. Mich., 494: 9.  
Holotype: FMNH 35350. Venezuela: Aragua, Rio Turmero.  
Paratypes: 2. Same locality.  
PIMELODIDAE
- Microglanis poecilus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 155.  
Holotype: FMNH 47365 (CM 1676). Guyana: below Packeo Falls.  
Paratype. Same locality.  
PIMELODIDAE
- Microglanis secundus** Mees, 1974  
Zool. Verh. (Leiden), 132: 235.  
Paratypes: 3. Surinam: Coppename River.  
PIMELODIDAE
- Microglanis variegatus** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 14.  
Paratype. Ecuador: Vines.  
PIMELODIDAE
- Micrognathus suvensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 396.  
Holotype: FMNH 17229. Fiji Islands: Viti Levu Island, Suva Harbor.  
SYNGNATHIDAE
- Micropterus pseudaplites** Hubbs, 1927  
Occas. Pap. Mus. Zool. Univ. Mich., 184: 8.  
Paratypes: 13. Arkansas: 6 localities.  
CENTRARCHIDAE
- Microschemobrycon guaporensis** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 56.  
Holotype: FMNH 57926 (CM 6910). Brazil: Rio Guaporé at Maciel.  
CHARACIDAE
- Microspathodon frontatus** Emery, 1970  
Stud. Trop. Oceanogr. Inst. Mar. Sci. Miami, 4(2): 294.  
Paratypes: 35. Gulf of Guinea: 3 localities.  
POMACENTRIDAE
- Mionorus pacificus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 403.  
Holotype: FMNH 17311. Ecuador: Islas Galápagos, Isla Albemarle, Tagus Cove.  
Paratype. Same locality.  
APOGONIDAE
- Misgurnus anguillicaudatus tungting** Nichols, 1925  
Am. Mus. Nov., 169: 3.  
Paratypes: 2. China: Hunan, Tungting Lake.  
COBITIDAE
- Misgurnus mizolepis fukien** Nichols, 1925  
Am. Mus. Nov., 169: 4.  
Paratype. China: Fukien.  
COBITIDAE
- Misgurnus mizolepis grangeri** Nichols, 1925  
Am. Mus. Nov., 169: 5.  
Paratypes: 2. China: Szechwan.  
COBITIDAE
- Misgurnus mohoity yunnan** Nichols, 1925  
Am. Mus. Nov., 169: 5.  
Paratype. China: Yunnan.  
COBITIDAE
- Mnierpes macrocephalus catherinae** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 61.  
Paratypes: 3. Nicaragua and Costa Rica: 2 localities.  
LABRISOMIDAE
- Moenkhausia browni** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 13.  
Holotype: FMNH 52733 (CM 1106). Guyana: Potaro River at Aruataima Falls.  
Paratypes: 28. Guyana: 7 localities.  
CM 1106 was received from Carnegie Museum with 2 specimens and 2 labels, one of each reads "1106 type (a)". FMNH 53732 (64 mm total length) is most likely the holotype, but we defer a final determination to a revisor.  
CHARACIDAE
- Moenkhausia lepidura hasemani** Eigenmann, 1917  
Mem. Carnegie Mus., 43(1): 102.  
Holotype: FMNH 55055 (CM 3746). Brazil: Santarém.  
CHARACIDAE
- Moenkhausia metae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 234.  
Paratypes: 4. Colombia: Villavicencio.  
CHARACIDAE
- Moenkhausia phaeonota** Fink, 1979  
Breviora, 450: 2.  
Paratype. Brazil: Mato Grosso, Rio Arinos.  
CHARACIDAE

- Moenkhausia profunda** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 322.  
 Holotype: FMNH 53717 (CM 2207). Guyana:  
 Issorora Rubber Plantation.  
 CHARACIDAE
- Moenkhausia shideleri** Eigenmann, 1909  
 Ann. Carnegie Mus., 6(1): 15.  
 Holotype: FMNH 52961 (CM 1012). Guyana:  
 Bartica.  
 CHARACIDAE
- Monacanthus tuckeri** Bean, 1906  
 Proc. Biol. Soc. Wash., 19: 33.  
 Lectotype: FMNH 5183. Bermuda: The Flatts.  
 Lectoparatypes: 2. Bermuda: 2 localities.  
 Lectotype designated by Grey (1947, p. 168).  
 BALISTIDAE
- Monolene megalepis** Woods, 1961  
 Copeia, 1961(2): 192.  
 Holotype: FMNH 65746. Puerto Rico: west of  
 Arecibo.  
 Paratypes: 9. Caribbean Sea: 4 localities.  
 BOTHIDAE
- Monomitopus kumae** Jordan and Hubbs, 1925  
 Mem. Carnegie Mus., 10(2): 324.  
 Holotype: FMNH 58844 (CM 7954). Japan:  
 Misaki.  
 OPHIDIIDAE
- Moroco yamamotis** Jordan and Hubbs, 1925  
 Mem. Carnegie Mus., 10(2): 182.  
 Holotype: FMNH 58723 (CM 7829). Korea:  
 Lake Yamanaka.  
 CYPRINIDAE
- Moxostoma rubreques** Hubbs, 1930  
 Misc. Publ. Mus. Zool. Univ. Mich., 20: 24.  
 Paratypes: 2. Michigan: Raisin River at Dundee.  
 CATOSTOMIDAE
- Mugil anpinensis** Oshima, 1922  
 Ann. Carnegie Mus., 13(3-4): 245.  
 Holotype: FMNH 59143 (CM 8281). Taiwan:  
 Anpin.  
 MUGILIDAE
- Mugilops cyanellus** Meek and Hildebrand, 1923  
 Field Mus. Nat. Hist. Zool. Ser., 15(1): 271.  
 Paratypes: 2. Panama: Canal Zone, 2 localities.  
 ATHERINIDAE
- Mulloidies flammeus** Jordan and Evermann, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 186.  
 Paratype: MISSING. Hawaiian Islands: Hilo.  
 MULLIDAE
- Myctophum hollandi** Jordan and Jordan, 1922  
 Mem. Carnegie Mus., 10(1): 11.  
 Holotype: FMNH 55191 (CM 3897). Hawaiian  
 Islands: Honolulu Market.  
 MYCTOPHIDAE
- Mylossoma ocellatum** Eigenmann, 1915  
 Ann. Carnegie Mus., 9(3-4): 265.  
 Holotype: FMNH 56816 (CM 5629). Paraguay:  
 Río Confuso at Villa Hays.  
 Paratypes: 15. Same locality.  
 CHARACIDAE
- Myoglanis potaroensis** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 159.  
 Holotype: FMNH 53333 (CM 1664). Guyana:  
 Tukeit.  
 Paratypes: 14. Guyana: 5 localities.  
 PIMELODIDAE
- Myripristis argyromus** Jordan and Evermann,  
 1903  
 Bull. U.S. Fish Comm. 1902, 22: 172.  
 Paratype. Hawaiian Islands: Honolulu.  
 HOLOCENTRIDAE
- Myripristis berndti** Jordan and Evermann, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 170.  
 Paratype. Hawaiian Islands: Honolulu.  
 HOLOCENTRIDAE
- Myripristis bowditchae** Woods, 1953  
 Bull. U.S. Natl. Mus., 202(1): 202.  
 Paratypes: 4. Marshall Islands: 3 localities.  
 HOLOCENTRIDAE
- Myripristis chryseres** Jordan and Evermann, 1903  
 Bull. U.S. Fish Comm. 1902, 22: 171.  
 Paratype. Hawaiian Islands: Hilo.  
 HOLOCENTRIDAE
- Myripristis mooreanus** Herre, 1935  
 Field Mus. Nat. Hist. Zool. Ser., 18(12): 394.  
 Holotype: FMNH 17225. Society Islands: Moo-  
 rea Island.  
 Paratypes: 2. Same locality.  
 HOLOCENTRIDAE

- Myripristis randalli** Greenfield, 1974  
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 34.  
Paratype. Austral Islands: Rurutu.  
HOLOCENTRIDAE
- Myripristis symmetricus** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 173.  
Paratype. Hawaiian Islands: Honolulu.  
HOLOCENTRIDAE
- Myripristis tiki** Greenfield, 1974  
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 9.  
Paratype. Chile: Easter Island, Hanga Pico.  
HOLOCENTRIDAE
- Myripristis undecimalis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 393.  
Holotype: FMNH 17228. Tuamotu Islands: Takaroa Island.  
HOLOCENTRIDAE
- Myripristis woodsi** Greenfield, 1974  
Nat. Hist. Mus. Los Ang. Cty. Sci. Bull., 19: 32.  
Paratypes: 3. Marshall Islands: Eniwetok Atoll, Giriinien Island.  
HOLOCENTRIDAE
- Mystus sabanus** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 294.  
Holotype: FMNH 68088. Malaysia: Sabah, Kinabatangan River at Deramakot.  
Paratypes: 13. Malaysia: Sabah, 3 localities.  
BAGRIDAE
- Myxine biniplicata** Richardson and Jowett, 1951  
Zool. Publ. Victoria Univ. Wellington, 12: 3.  
Paratype. New Zealand: Cook Strait.  
MYXINIDAE
- Myxus malayanus** Herre, 1936  
Philipp. J. Sci., 59: 286.  
Paratypes: 2. Sulawesi Island (Celebes): Lembeh Strait.  
MUGILIDAE
- Nannacara bimaculata** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 487.  
Holotype: FMNH 53799 (CM 2304). Guyana: Erukin.  
CICHLIDAE
- Nannorhamdia nemacheir** Eigenmann and Fisher, 1916  
In: Eigenmann, Ann. Carnegie Mus., 10(1–2): 83.  
Holotype: FMNH 58127 (CM 7125). Colombia: Girardot.  
Paratype. Same locality.  
PIMELODIDAE
- Nannostomus marginatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 41.  
Holotype: FMNH 53550 (CM 1171). Guyana: Maduni Creek.  
Paratypes: 17. Guyana: 7 localities.  
LEBIASINIDAE
- Nannostomus minimus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 42.  
Holotype: FMNH 52771 (CM 1165). Guyana: Erukin.  
Paratype. Guyana: Amatuik.  
LEBIASINIDAE
- Nannostomus simplex** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 42.  
Holotype: FMNH 52773 (CM 1167). Guyana: Lama Stop-off.  
LEBIASINIDAE
- Nanochromis transvestitus** Stewart and Roberts, 1984  
Copeia, 1984(1): 82.  
Holotype: FMNH 93584. Zaïre: Lac Mai-ndombe near Ipeque.  
Paratypes: 80. Same locality.  
CICHLIDAE
- Nansenia ardesiaca** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 210.  
Holotype: FMNH 57086 (CM 6023). Japan: Sagami Sea, Okinose.  
BATHYLAGIDAE
- Neetroplus carpintis** Jordan and Snyder, 1900  
Bull. U.S. Fish Comm. 1899, 19: 146.  
Paratype. Mexico: Laguna del Carpinte near Tampico.  
CICHLIDAE
- Neetroplus fluviatilis** Meek, 1912  
Field Mus. Nat. Hist. Zool. Ser., 10(7): 74.  
Holotype: FMNH 7685. Costa Rica: Costa Rica River at Guapilis.  
Paratype. Same locality.  
CICHLIDAE

- Neetroplus panamensis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 90.  
Holotype: FMNH 7601. Panama: Río Mandingo at Bas Obispo.  
Paratypes: 20. Panama: 6 localities.  
CICHLIDAE
- Nemaclinus atelestos** Böhlke and Springer, 1975  
Proc. Acad. Nat. Sci. Phila., 127(7): 58.  
Paratype. Caribbean Sea: off Virgin Islands.  
LABRISOMIDAE
- Nematabramis alestes borneensis** Inger and Chin, 1962  
Fieldiana Zool., 45: 47.  
Holotype: FMNH 44791. Malaysia: Sabah, Kota Belud.  
Paratypes: 7. Same locality.  
CYPRINIDAE
- Nematobrycon amphioxus** Eigenmann and Wilson, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 13.  
Holotype: (CM 5050) MISSING. Colombia: Boca de Raspadura.  
Paratypes: 15. Colombia: 3 localities.  
CHARACIDAE
- Nematobrycon lacortei** Weitzman and Fink, 1971  
Beaufortia, 19(248): 59.  
Paratype. Aquarium specimen, probably from Colombia: Río Calima.  
CHARACIDAE
- Neoharriotta carri** Bullis and Carpenter, 1966  
Copeia, 1966(3): 443.  
Paratypes: 2. Southern Caribbean Sea: 2 localities.  
RHINOCHIMAERIDAE
- Neoheterandria elegans** Henn, 1916  
Ann. Carnegie Mus., 10(1-2): 118.  
Holotype: FMNH 57007 (CM 5823). Colombia: Truandó.  
Paratypes: 7. Same locality.  
POECILIIDAE
- Neostethus borneensis** Herre, 1939  
Proc. Biol. Soc. Wash., 52: 143.  
Paratypes: 12. Malaysia: Sabah, Kibili River.  
NEOSTETHIDAE
- Nesiotes purpurascens** De Vis, 1884  
Proc. Linn. Soc. N. S. Wales, 8(4): 453.  
Neotype: FMNH 17340. New Hebrides Islands: Efate Island, Vila.  
Neotype designated by Herre (1936, Field Mus. Nat. Hist. Zool. Ser., 21: 165).  
PSEUDOCROMIDAE
- Netuma osakae** Jordan and Kasawa, 1925  
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 157.  
Holotype: FMNH 59388 (CM 7808). Japan: market at Osaka.  
ARIIDAE
- Netuma planifrons** Higuchi, Reis, and Araújo, 1982  
Atlântica, Rio Grande, 5(1): 12.  
Paratypes: 5. Brazil: Rio Grande do Sul, Lagoa dos Patos.  
ARIIDAE
- Nototropis forlonensis** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 70.  
Holotype: FMNH 4478. Mexico: Tamaulipas, Forlón.  
Paratypes: 28. Mexico: 2 localities.  
CYPRINIDAE
- Nototropis nazas** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 70.  
Holotype: FMNH 4375. Mexico: Río Nazas at Santiago Papasquiaro.  
Paratypes: 133. Same locality.  
CYPRINIDAE
- Notropis aztecus** Woolman, 1894  
Bull. U.S. Fish Comm. 1894, 14: 63.  
Syntypes: 11, FMNH 6604. Mexico: Canal in Mexico City.  
CYPRINIDAE
- Notropis chihuahua** Woolman, 1894  
Am. Nat., 26(3): 260.  
Syntype: FMNH 1889. Mexico: Chihuahua, Río de los Conchos.  
CYPRINIDAE
- Notropis edwardraneyi** Suttkus and Clemmer, 1968  
Tulane Stud. Zool. Bot., 15(1): 19.  
Paratypes: 100. Alabama: Alabama River, at Yellow Jacket Bar.  
CYPRINIDAE

- Notropis greenei** Hubbs and Ortenburger, 1929  
 Publ. Univ. Okla. Biol. Surv., 1: 78.  
 Paratypes: 37. Arkansas: Kinderhook.  
 CYPRINIDAE
- Notropis muskoka** Meek, 1899  
 Field Mus. Nat. Hist. Zool. Ser., 1(17): 308.  
 Holotype: FMNH 2964. Canada: Ontario, Gull Lake.  
 Paratypes: 20. Same locality.  
 CYPRINIDAE
- Notropis panarcys** Hubbs and Miller, 1978  
 Copeia, 1978(4): 582.  
 Paratypes: 10. Mexico: Chihuahua, Río San Pedro at Meoqui.  
 CYPRINIDAE
- Notropis robustus** Meek, 1902  
 Field Mus. Nat. Hist. Zool. Ser., 3(6): 82.  
 Holotype: FMNH 3548. Mexico: Río Conchos at Jiménez.  
 Paratypes: 19. Mexico: 2 localities.  
 CYPRINIDAE
- Notropis saladonis** Hubbs and Hubbs, 1958  
 Copeia, 1958(4): 297.  
 Paratypes: 18. Mexico: 2 localities.  
 CYPRINIDAE
- Notropis santarosaliae** Meek, 1902  
 Field Mus. Nat. Hist. Zool. Ser., 3(6): 85.  
 Holotype: FMNH 3535. Mexico: Río Noavaco at Santa Rosalia.  
 Paratypes: 17. Same locality.  
 CYPRINIDAE
- Noturus albater** Taylor, 1969  
 U.S. Natl. Mus. Bull., 282: 144.  
 Paratypes: 4. Arkansas: Main Fork White River at Fayetteville.  
 ICTALURIDAE
- Noturus flavater** Taylor, 1969  
 U.S. Natl. Mus. Bull., 282: 104.  
 Paratype. Missouri: Shannon County, Montier.  
 ICTALURIDAE
- Noturus munitus** Suttkus and Taylor, 1965  
 Proc. Biol. Soc. Wash., 78: 171.  
 Paratypes: 10. Mississippi: Marion County, Pearl River.  
 ICTALURIDAE
- Novumbra hubbsi** Schultz, 1929  
 Publ. Univ. Wash. Coll. Fish., 2(6): 4.  
 Paratype. Washington: one-half mile west of Satsop.  
 UMBRIDAE
- Ochmacanthus flabelliferus** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 213.  
 Holotype: FMNH 53263 (CM 1729). Guyana: Konawaruk.  
 TRICHOMYCTERIDAE
- Odontognathus compressus** Meek and Hildebrand, 1923  
 Field Mus. Nat. Hist. Zool. Ser., 15(1): 194.  
 Paratypes: 23. Panama: Fox Bay, Colón.  
 CLUPEIDAE
- Odontorhamphus cancellori** Weed, 1933  
 Field Mus. Nat. Hist. Zool. Ser., 20(7): 52.  
 Holotype: FMNH 16208. Cook Archipelago: Aitutaki Island.  
 HEMIRAMPHIDAE
- Odontostilbe hastatus** Eigenmann, 1913  
 Indiana Univ. Stud., 18: 27.  
 Holotype: FMNH 56383 (CM 5103). Colombia: Soplaviento.  
 Paratypes: 13. Colombia: Bernal Creek.  
 CHARACIDAE
- Odontostilbe melandetus** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 312.  
 Holotype: (CM 1878) MISSING. Guyana.  
 CHARACIDAE
- Odontostilbe paraguayensis** Eigenmann and Kennedy, 1903  
 Proc. Acad. Nat. Sci. Phila., 59: 512.  
 Paratype. Paraguay: Arroyo Trementina.  
 CHARACIDAE
- Odontostilbe trementinae** Eigenmann and Kennedy, 1903  
 Proc. Acad. Nat. Sci. Phila., 59: 513.  
 Paratypes: 2. Paraguay: Arroyo Trementina.  
 CHARACIDAE
- Ogocephalus corniger** Bradbury, 1980  
 Proc. Calif. Acad. Sci., 42(7): 274.  
 Paratypes: 23. Caribbean Sea and Atlantic Ocean: 12 localities.  
 OGCOEPHALIDAE

- Ogcocephalus declivirostris** Bradbury, 1980  
Proc. Calif. Acad. Sci., 42(7): 269.  
Holotype: FMNH 64136. Gulf of Mexico: off Texas.  
Paratypes: 21. Gulf of Mexico: 14 localities.  
OGCOEPHALIDAE
- Ogcocephalus pantostictus** Bradbury, 1980  
Proc. Calif. Acad. Sci., 42(7): 264.  
Holotype: FMNH 50765. Florida: Pensacola, Santa Rosa Sound.  
Paratypes: 8. Gulf of Mexico: 6 localities.  
OGCOEPHALIDAE
- Ogcocephalus pumilus** Bradbury, 1980  
Proc. Calif. Acad. Sci., 42(7): 272.  
Paratypes: 75. Caribbean Sea: 14 localities.  
OGCOEPHALIDAE
- Ogcocephalus rostellum** Bradbury, 1980  
Proc. Calif. Acad. Sci., 42(7): 267.  
Paratypes: 17. Atlantic Ocean: North Carolina to Florida, 4 localities.  
OGCOEPHALIDAE
- Oligobrycon microstomus** Eigenmann, 1915  
Mem. Carnegie Mus., 7(1): 57.  
Holotype: FMNH 57913 (CM 6898). Brazil: Rio Paratyba at Jacarehy.  
Paratype. Same locality.  
CHARACIDAE
- Ompok sabanus** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 282.  
Holotype: FMNH 44828. Malaysia: Sabah, Segama River near Lahad Datu.  
Paratypes: 106. Malaysia: Sabah, 15 localities.  
SILURIDAE
- Oncorhynchus adonis** Jordan and McGregor, 1925  
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 127.  
Holotype: FMNH 58680 (CM 7784). Japan: Sagami, Lake Hakone.  
SALMONIDAE
- Oncorhynchus ishikawae** Jordan and McGregor, 1925  
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 132.  
Holotype: FMNH 58682 (CM 7786). Japan: Lake Biwa.  
SALMONIDAE
- Oncorhynchus kawamurae** Jordan and McGregor, 1925  
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 128.  
Holotype: FMNH 58681 (CM 7785). Japan: Ugo, Lake Toyama.  
SALMONIDAE
- Oncorhynchus rhodurus** Jordan and McGregor, 1925  
In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 137.  
Holotype: FMNH 59389 (CM 7794). Japan: Sagami, Lake Hakone.  
Paratype. Same locality.  
SALMONIDAE
- Ophichthus evermanni** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 172.  
Holotype: FMNH 52118 (CM 246). Taiwan: Takao.  
OPHICHTHIDAE
- Ophichthus hyposagmatus** McCosker and Böhlke, 1984  
Proc. Acad. Nat. Sci. Phila., 136: 24.  
Paratype. Caribbean Sea: between Honduras and Jamaica.  
OPHICHTHIDAE
- Ophioscion punctatissimus** Meek and Hildebrand, 1925  
Field Mus. Nat. Hist. Zool. Ser., 15(2): 644.  
Paratypes: 2. Panama: 2 localities.  
SCIAENIDAE
- Ophioscion venezuelae** Schultz, 1945  
Proc. U.S. Natl. Mus., 96(3192): 131.  
Paratype. Venezuela: Caño de Sagua north of Sinamaica.  
SCIAENIDAE
- Opisthonema berlangai** Berry and Barrett, 1963  
Inter-Am. Trop. Tuna Comm. Bull., 7(2): 119.  
Paratypes: 6. Ecuador: Islas Galápagos, Isla Barrington (Santa Fe).  
CLUPEIDAE
- Opisthonema medirastre** Berry and Barrett, 1963  
Inter-Am. Trop. Tuna Comm. Bull., 7(2): 118.  
Paratypes: 5. Panama and Mexico: 2 localities.  
CLUPEIDAE



- Optonurus atherodon** Gilbert and Cramer, 1896  
Proc. U.S. Natl. Mus., 19(1114): 431.  
Paratype. Hawaiian Islands.  
MACROURIDAE
- Oreias furcatus** Banareescu and Nalbant, 1976  
Nimphaca, 4: 188.  
Paratypes: 4. China: Szechwan, Baurong.  
COBITIDAE
- Orestias mundus** Parenti, 1984  
Bull. Am. Mus. Nat. Hist., 178(2): 199.  
Holotype: FMNH 41135. Peru: Cuzco.  
Paratypes: 110. Same locality.  
CYPRINODONTIDAE
- Ostichthys acanthorhinus** Randall, Shimizu, and Yamakawa, 1982  
Jpn. J. Ichthyol., 29(1): 10.  
Holotype: FMNH 70251. Arabian Sea: Gulf of Oman.  
Paratypes: 3. Arabian Sea: Gulf of Oman, 3 localities.  
HOLOCENTRIDAE
- Otolithes lini** Herre, 1935  
Lingnan Sci. J. Canton, 14(4): 603.  
Paratype. China: Canton Market.  
SCIAENIDAE
- Otophidium schmidtii** Woods and Kanazawa, 1951  
Fieldiana Zool., 31(53): 640.  
Holotype: FMNH 48175. Bermuda: Castle Harbor, Cock Rock.  
OPHIDIIDAE
- Oxyloricaria citurensis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 82.  
Holotype: FMNH 7585. Panama: Río Cupe at Cituro.  
Paratypes: 150. Panama: 5 localities.  
LORICARIIDAE
- Oxyloricaria dariensis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 81.  
Holotype: FMNH 7584 MISSING or mixed.  
Panama: Río Tuyra, Boca de Cupe.  
Paratypes: 39. Panama: 7 localities.  
LORICARIIDAE
- Pangasius tubbi** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 287.  
Holotype: FMNH 68047. Malaysia: Sabah, confluence of Deramakot River with Kinabatangan River.  
Paratypes: 18. Malaysia: Sabah, 4 localities.  
PANGASIIDAE
- Pantosteus jordani** Evermann, 1893  
Bull. U.S. Fish Comm. 1892, 12: 51.  
Paratypes: 2 (disintegrated). South Dakota: Hat Creek, Ardmore.  
CATOSTOMIDAE
- Pantosteus santaanae** Snyder, 1908  
Proc. U.S. Natl. Mus., 34(1595): 33.  
Paratypes: 6. California: Santa Ana River near Riverside.  
CATOSTOMIDAE
- Paraclinus integripinnis zacae** Hubbs, 1952  
Stanford Ichthyol. Bull., 4(2): 87.  
Paratypes: 18. Mexico: Baja California, Punta Aberrojos.  
LABRISOMIDAE
- Paracrossochilus acerus** Inger and Chin, 1962  
Fieldiana Zool., 45: 100.  
Holotype: FMNH 68240. Malaysia: Sarawak, Sungei Dapu River.  
Paratypes: 600. Malaysia: Sarawak, 10 localities.  
CYPRINIDAE
- Parahollardia schmidtii** Woods, 1959  
Copeia, 1959(3): 223.  
Paratypes: 7. Western Caribbean Sea: 3 localities.  
TRIACANTHODIDAE
- Paralepis philippinus** Fowler, 1934  
Proc. Acad. Nat. Sci. Phila., 85: 281.  
Paratypes: 2. Philippine Islands: Leyte.  
PARALEPIDAE
- Parapelecus eigenmanni** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 21.  
Holotype: FMNH 55802 (CM 4565). Korea: Suigen.  
CYPRINIDAE
- Parascolopsis townsendi** Boulenger, 1901  
Ann. Mag. Nat. Hist., Ser. 7, 7: 262.  
Syntype: FMNH 5736. Sea of Oman.  
NEMIPTERIDAE
- Parastremma sadina** Eigenmann, 1912  
Indiana Univ. Stud., 16: 20.  
Holotype: FMNH 56022 (CM 4812). Colombia: Istmina.  
Paratypes: 11. Same locality.  
CHARACIDAE

**Paratrachichthys argyrophanus** Woods, 1961

Fieldiana Zool., 39(45): 525.

Paratypes: 13. Atlantic Ocean: off Brazil, 3 localities.

TRACHICHTHYIDAE

**Parecbasis cyclolepis** Eigenmann, 1914

Indiana Univ. Stud., 20: 45.

Holotype: FMNH 56677 (CM 5495). Brazil: São Antonio de Rio Madeira.

Paratype. Same locality.

CHARACIDAE

**Parodon bifasciatus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 274.

Holotype: FMNH 53582 (CM 1925). Guyana: Maripicru Creek.

PARODONTIDAE

**Parodon dariensis** Meek and Hildebrand, 1913

Field Mus. Nat. Hist. Zool. Ser., 10(8): 83.

Holotype: FMNH 7587. Panama: Río Cupe at Cituro.

Paratype. Same locality.

PARODONTIDAE

**Parodon ecuadoriensis** Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 12.

Holotype: FMNH 56597 (CM 5413). Ecuador: Vinces.

Paratypes: 8. Ecuador: 2 localities.

PARODONTIDAE

**Parodon terminalis** Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 12.

Holotype: FMNH 56599 (CM 5415). Ecuador: Vinces.

Paratype. Same locality.

PARODONTIDAE

**Parotocinclus jimi** Garavello, 1976

Arq. Zool. (São Paulo), 28(4): 9.

Paratypes: 3. Brazil: Bahia, Rio do Peixe at Fazenda Pedra Branca, Itagiba.

LORICARIIDAE

**Pellonula tenuis** Nichols and Griscom, 1917

Bull. Am. Mus. Nat. Hist., 37(25): 675.

Paratypes: 5. Zaïre: Stanleyville.

CLUPEIDAE

**Pempheris dispar** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 407.

Holotype: FMNH 17326. Solomon Islands: Isabel Island, Tenibuli.

Paratypes: 6. Same locality.

PEMPHERIDAE

**Peristedion gilberti** Jordan, 1922

Proc. U.S. Natl. Mus., 59(2392): 655.

Paratype. Hawaiian Islands: deep water of Oahu.

TRIGLIDAE

**Peristedion greyae** Miller, 1967

Bull. Mar. Sci., 17(1): 27.

Paratypes: 2. Western North Atlantic and Caribbean Sea: 2 localities.

TRIGLIDAE

**Petersius barnardi** Herre, 1936

Proc. Biol. Soc. Wash., 49: 100.

Paratypes: 8. Mozambique: near Beira.

CHARACIDAE

**Petersius conserialis** Hilgendorf, 1894

Sitzungsber. Ges. naturf. Freunde Berl., 1894: 173.

Paratype. Tanzania: Kingani River at Dunda.

CHARACIDAE

**Petroscirtes kulambangrae** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 436.

Holotype: FMNH 17392. Solomon Islands: Kulambangra Island.

BLENNIIDAE

**Phalloptychus eigenmanni** Henn, 1916

Ann. Carnegie Mus., 10(1-2): 121.

Holotype: FMNH 55876 (CM 4665). Brazil: Rio Catu at Alagoinhas.

Paratypes: 10. Same locality.

POECILIIDAE

**Phallotorynus fasciolatus** Henn, 1916

Ann. Carnegie Mus., 10(1-2): 129.

Holotype: FMNH 55061 (CM 3752). Brazil: São Paulo, Jacarehy.

Paratypes: 10. Same locality.

POECILIIDAE

**Phenacogaster beni** Eigenmann, 1911

Ann. Carnegie Mus., 8(1): 174.

Holotype: FMNH 54597 (CM 3229). Bolivia: Río Beni at Villa Bella.

Paratype. Same locality.

CHARACIDAE

**Phenacogaster franciscoensis** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 173.  
Holotype: FMNH 54599 (CM 3231). Brazil: Boqueirão, near mouth of Rio Preto.  
Paratypes: 16. Brazil: 4 localities.  
CHARACIDAE

**Phenacogaster megalostictus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 29.  
Holotype: FMNH 53530 (CM 1056). Guyana: Tumatumari.  
Paratypes: 47. Guyana: 6 localities.  
CHARACIDAE

**Phenacogaster microstictus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 30.  
Holotype: FMNH 52970 (CM 1063). Guyana: Potaro River at Tumatumari.  
Paratypes: 5. Guyana: 3 localities.  
CHARACIDAE

**Phenagoniates wilsoni** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 2.  
Holotype: FMNH 56540 (CM 5354). Colombia: Manigrú.  
Paratypes: 2. Colombia: 2 localities.  
CHARACIDAE

**Phoxinus cumberlandensis** Starnes and Starnes, 1978  
Copeia, 1978(3): 509.  
Paratypes: 16. Kentucky: South Fork, Dog Slaughter Creek.  
CYPRINIDAE

**Phoxinus orcuttii** Eigenmann and Eigenmann, 1890  
Proc. Calif. Acad. Sci., Ser. 2, 3: 2.  
Syntypes: 10, FMNH 54985 (CM 3675). California: Temecula River at Riverside.  
CYPRINIDAE

**Phoxiscus kikuchii** Oshima, 1919  
Ann. Carnegie Mus., 12(2–4): 226.  
Holotype: FMNH 59099 (CM 8237). Taiwan: Bokusekikaku.  
CYPRINIDAE

**Phrynelox atra** Schultz, 1957  
Proc. U.S. Natl. Mus., 107(3383): 76.  
Holotype: FMNH 21705. Australia: Sydney.  
Paratype. Same locality.  
ANTENNARIIDAE

**Phrynelox zebrinus** Schultz, 1957  
Proc. U.S. Natl. Mus., 107(3383): 75.  
Paratypes: 2. Australia: 2 localities.  
ANTENNARIIDAE

**Physailia villiersi** Boulenger, 1912  
Ann. Mus. Congo Belge Ser. Quarto Zool., Ser. 1, 2(3): 17.  
Syntypes: 2, FMNH 56151 (CM 4942). Zaïre: Luculla River.  
SCHILBIDAE

**Physiculus grinnelli** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 22.  
Holotype: FMNH 55192 (CM 3898). Hawaiian Islands: Honolulu Market.  
MORIDAE

**Piabina analis** Eigenmann, 1914  
In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 8.  
Holotype: FMNH 56661 (CM 5478). Brazil: Cáceres.  
CHARACIDAE

**Pikea aurora** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 178.  
Paratype. Hawaiian Islands: Honolulu.  
SERRANIDAE

**Pikea mexicanus atlanticus** Schultz, 1958  
Proc. U.S. Natl. Mus., 108(3405): 326.  
Paratype. Atlantic Ocean: off Surinam.  
SERRANIDAE

**Pikea mexicanus mexicanus** Schultz, 1958  
Proc. U.S. Natl. Mus., 108(3405): 323.  
Paratype. Gulf of Mexico: off Florida, south of Panama City.  
SERRANIDAE

**Pimelodella avanhandavae** Eigenmann, 1917  
Mem. Carnegie Mus., 7(4): 240.  
Holotype: FMNH 57981 (CM 6969). Brazil: Rio Tietê above Salto Avanhandava.  
Paratypes: 30. Brazil: 2 localities.  
PIMELODIDAE

**Pimelodella boliviana** Eigenmann, 1917  
Mem. Carnegie Mus., 7(4): 245.  
Holotype: FMNH 57976 (CM 6964). Bolivia: Santa Cruz de la Sierra.  
Paratype. Same locality.  
PIMELODIDAE

- Pimelodella eigenmanni* Meek, 1905**  
 Proc. Biol. Soc. Wash., 18: 241.  
 Holotype: FMNH 3400. Brazil: São Paulo.  
 Paratypes: 5. Same locality.  
 PIMELODIDAE
- Pimelodella griffini* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 250.  
 Holotype: FMNH 57974 (CM 6962). Paraguay:  
 near Sapucay.  
 Paratypes: 7. Same locality.  
 PIMELODIDAE
- Pimelodella hasemani* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 241.  
 Holotype: FMNH 57980 (CM 6968). Brazil: São  
 Antonio de Rio Madeira.  
 PIMELODIDAE
- Pimelodella itapicuruensis* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 247.  
 Holotype: FMNH 57986 (CM 6974). Brazil: Rio  
 Itapicuru at Queimadas.  
 Paratypes: 21. Brazil: 4 localities.  
 PIMELODIDAE
- Pimelodella laticeps* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 243.  
 Holotype: FMNH 57969 (CM 6957). Brazil:  
 Uruguayana.  
 Paratypes: 3. Same locality.  
 PIMELODIDAE
- Pimelodella laticeps australis* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 243.  
 Holotype: FMNH 57962 (CM 6950). Brazil:  
 Uruguayana.  
 Paratypes: 43. Same locality.  
 PIMELODIDAE
- Pimelodella macturki* Eigenmann, 1912**  
 Mem. Carnegie Mus., 5: 170.  
 Holotype: FMNH 53234 (CM 1695). Guyana:  
 creek in Mora Passage.  
 Paratypes: 11. Guyana: 3 localities.  
 PIMELODIDAE
- Pimelodella megalops* Eigenmann, 1912**  
 Mem. Carnegie Mus., 5: 169.  
 Holotype: FMNH 53231 (CM 1692). Guyana:  
 Tumatumari.  
 Paratypes: 35. Guyana: 2 localities.  
 PIMELODIDAE
- Pimelodella metae* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 244.  
 Holotype: FMNH 58441 (CM 7441). Colombia:  
 Río Negro at Villavicencio.  
 Paratypes: 2. Colombia: Río Meta at Barrigona.  
 PIMELODIDAE
- Pimelodella notomelas* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 244.  
 Holotype: FMNH 57967 (CM 6955). Brazil:  
 Cáceres.  
 Paratypes: 9. Brazil: 2 localities.  
 PIMELODIDAE
- Pimelodella serrata* Eigenmann, 1917**  
 Mem. Carnegie Mus., 7(4): 235.  
 Holotype: FMNH 57979 (CM 6967). Bolivia:  
 San Joaquín.  
 Paratype. Same locality.  
 PIMELODIDAE
- Pimelodus clarias coprophagus* Schultz, 1944**  
 Proc. U.S. Natl. Mus., 94(3172): 203.  
 Paratypes: 4. Venezuela: Río Palmar near To-  
 tuma.  
 PIMELODIDAE
- Pimelodus heteropleurus* Eigenmann, 1912**  
 Mem. Carnegie Mus., 5: 176.  
 Holotype: FMNH 53336 (CM 1734). Guyana:  
 Rupununi Pan.  
 PIMELODIDAE
- Pimelodus ortmanni* Eigenmann, 1911**  
 Ann. Carnegie Mus., 7(3-4): 379.  
 Holotype: FMNH 54240 (CM 2856). Brazil: Rio  
 Iguaçu at Pôrto União da Victoria.  
 Paratypes: 5. Same locality.  
 PIMELODIDAE
- Pimephales anuli* Kendall, 1903**  
 Bull. U.S. Fish Comm. 1902, 22: 360.  
 Paratypes: 15. Maine: Lunkasoos Lake.  
 CYPRINIDAE
- Plagiotremus phenax* Smith-Vaniz, 1976**  
 Monogr. Acad. Nat. Sci. Phila., 19: 131.  
 Paratypes: 11. Maldive Islands: 3 localities.  
 BLENNIIDAE
- Platophrys inermis* Gilbert, 1905**  
 Bull. U.S. Fish Comm. 1903, 23(2): 685.  
 Paratype. Hawaiian Islands.  
 BOTHIDAE

**Platypoecilus tropicus** Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 146.  
Holotype: FMNH 6027. Costa Rica: Turrialba.  
Paratypes: 2. Same locality.  
POECILIIDAE

**Platypoecilus variatus** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 146.  
Holotype: FMNH 4501. Mexico: San Luis Potosí, Valles.  
Paratypes: 109. Mexico: 5 localities.  
POECILIIDAE

**Platysilurus barbatus** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 320.  
Holotype: FMNH 60306 (CM 2987). Brazil: São Antonio de Rio Madeira.  
Paratypes: 2. Bolivia: Río Machupo near San Joaquín.  
PIMELODIDAE

**Plecostomus derbyi** Haseman, 1911

Ann. Carnegie Mus., 7(3–4): 384.  
Holotype: FMNH 54246 (CM 2865). Brazil: Rio Iguaçu at Pôrto União da Victoria.  
LORICARIIDAE

**Plecostomus hemiurus** Eigenmann, 1912

Mem. Carnegie Mus., 5: 224.  
Holotype: FMNH 53110 (CM 1544). Guyana: Amatuik.  
Paratypes: 10. Guyana: 6 localities.  
LORICARIIDAE

**Plectranthias garrupellus** Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 295.  
Paratype. Western North Atlantic: no station data.  
SERRANIDAE

**Plesiops cephalotaenia** Inger, 1955

Pac. Sci., 9(3): 272.  
Holotype: FMNH 44708. Philippine Islands: Sit-tankai.  
Paratypes: 3. Same locality.  
PLESIOPIDAE

**Pleurogrammus azonus** Jordan and Metz, 1913

Mem. Carnegie Mus., 6(1): 47.  
Holotype: FMNH 55796 (CM 4558). Korea: Chinnampo.  
HEXAGRAMMIDAE

**Plotosus papuensis** Weber, 1910

Notes Leiden Mus., 32: 228.  
Syntype: FMNH 52494 (CM 752). New Guinea: Lorentz River.  
PLOTOSIDAE

**Pneumatophorus peruanus** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 211.  
Holotype: FMNH 58739 (CM 7847). Peru: Callao.  
SCOMBRIDAE

**Poblana ferdebueni** Solórzano and López, 1965

Ciencia (Mex. City), 23(5): 209.  
Paratypes: 3. Mexico: Puebla, Laguna de Al-moyala o Chignahuapan.  
ATHERINIDAE

**Poecilia amates** Miller, 1907

Bull. Am. Mus. Nat. Hist., 23(2): 108.  
Paratypes: 6. Guatemala: Los Amates.  
POECILIIDAE

**Poecilia caudata** Meek, 1909

Field Mus. Nat. Hist. Zool. Ser., 7(7): 209.  
Holotype: FMNH 6360. Costa Rica: Turru-bares.  
Allotype and paratypes: 10. Costa Rica: 3 localities.  
POECILIIDAE

**Poecilia fria** Eigenmann and Henn, 1914

In: Eigenmann, Henn, and Wilson, Indiana Univ. Stud., 19: 13.  
Holotype: FMNH 56603 (CM 5420). Ecuador: Vinces.  
Paratypes: 112. Same locality.  
POECILIIDAE

**Poecilia latipunctata** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 150.  
Holotype: FMNH 4484. Mexico: Tamaulipas, Forlón.  
Paratypes: 12. Same locality.  
POECILIIDAE

**Poecilia pittieri** Meek, 1912

Field Mus. Nat. Hist. Zool. Ser., 10(7): 71.  
Holotype: FMNH 7680. Costa Rica: Río Re-ventazón at La Junta.  
Allotype and paratypes: 15. Same locality.  
POECILIIDAE

**Poecilia tenuis** Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 147.  
Holotype (?): FMNH 6028. Costa Rica: Tiribi.  
Paratypes: 4. Costa Rica: 2 localities.

See comment in Grey (1947, p. 175). One of  
2 specimens in FMNH 6029 is also la-  
beled "type", and is closer to the size  
given for the type in Meek (1907).

POECILIIDAE

**Poeciliopsis balsas** Hubbs, 1926

Misc. Publ. Mus. Zool. Univ. Mich., 16: 66.  
Holotype: FMNH 3702. Mexico: Balsas, Gue-  
rrero.

Paratypes: 91. Mexico: 6 localities.

POECILIIDAE

**Poecilobrycon auratus** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 43.  
Holotype: FMNH 52768 (CM 1161). Guyana:  
Konawaruk.

Paratypes: 8. Guyana: 3 localities.

LEBIASINIDAE

**Poecilobrycon erythrurus** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 44.  
Holotype: FMNH 52774 (CM 1168). Guyana:  
Rockstone sand-bank.

Paratypes: 6. Guyana: 4 localities.

LEBIASINIDAE

**Poecilobrycon harrisoni** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 43.  
Holotype: FMNH 52767 (CM 1160). Guyana:  
canal at Christianburg.

LEBIASINIDAE

**Poecilobrycon ocellatus** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 45.  
Holotype: FMNH 52975 (CM 1179). Guyana:  
Wismar.

Paratypes: 25. Guyana: 6 localities.

LEBIASINIDAE

**Poecilocharax bovalii** Eigenmann, 1909

Ann. Carnegie Mus., 6(1): 34.  
Holotype: FMNH 53544 (CM 1136). Guyana:  
creek at Savanna Landing.

Paratypes: 171. Guyana: 4 localities.

CHARACIDAE

**Poecilurichthys moenkhausii** Eigenmann and  
Kennedy, 1903

Proc. Acad. Nat. Sci. Phila., 59: 522.  
Paratypes: 2. Paraguay: brook near Arroyo Tre-  
mentina.

CHARACIDAE

**Pogoneleotris microps** Weber, 1908

Nova Guinea, 5(2): 258.

Syntypes: 2, FMNH 53459 (CM 717). New  
Guinea: Merauke River.

ELEOTRIDAE

**Polyacanthonotus merretti** Sulak, Crabtree, and  
Hureau, 1984

Cybiurn, 8(4): 59.

Paratype. Atlantic Ocean: off Virginia, Norfolk  
Canyon.

NOTACANTHIDAE

**Polyipnus nuttingii** Gilbert, 1905

Bull. U.S. Fish Comm. 1903, 23(2): 609.

Paratypes: 6. Hawaiian Islands.

STERNOPTYCHIDAE

**Pomacentrus cranei** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 409.

Holotype: FMNH 17338. Solomon Islands: Is-  
abel Isand, Tenibuli.

Paratype. Same locality.

POMACENTRIDAE

**Pomacentrus jenkinsi** Jordan and Evermann, 1903

Bull. U.S. Fish Comm. 1902, 22: 189.

Paratypes: 3. Hawaiian Islands.

POMACENTRIDAE

**Pomacentrus reidi** Fowler and Bean, 1928

Bull. U.S. Natl. Mus., 100(7): 99.

Paratypes: 2. Philippine Islands: Dalaganem Is-  
land.

POMACENTRIDAE

**Pomacentrus stigma** Fowler and Bean, 1928

Bull. U.S. Natl. Mus., 100(7): 100.

Paratype. Philippine Islands: Mindoro, Paluan  
Bay.

POMACENTRIDAE

**Pomadasy grandis** Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(4): 116.

Holotype: FMNH 5939. Nicaragua: Lago Ni-  
caragua at Granada.

HAEMULIDAE

**Pomadasy starri** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 200.

Holotype: FMNH 4693. Mexico: Veracruz, Pé-  
rez.

HAEMULIDAE

- Pomadasy templei** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 201.  
Holotype: FMNH 4504. Mexico: San Luis Potosí, Valles.  
Paratypes: 2. Mexico: Veracruz, Pérez.  
HAEMULIDAE
- Porichthys bathoiketes** Gilbert, 1968  
Bull. Mar. Sci., 18(3): 710.  
Paratype. Caribbean Sea: off Honduras.  
BATRACHOIDIDAE
- Porichthys kymosemum** Gilbert, 1968  
Bull. Mar. Sci., 18(3): 724.  
Holotype: FMNH 66688. Atlantic Ocean: off Brazil.  
BATRACHOIDIDAE
- Porichthys myriaster** Hubbs and Schultz, 1939  
Proc. U.S. Natl. Mus., 86(3060): 478.  
Paratypes: 4. California: off Mira Mar pier.  
BATRACHOIDIDAE
- Porogadus silus** Carter and Sulak, 1984  
Bull. Mar. Sci., 34(3): 361.  
Paratypes: 2. Western North Atlantic: The Bahamas region.  
OPHIDIIDAE
- Porotergus gimbeli** Ellis, 1912  
Mem. Carnegie Mus., 5: 441.  
Holotype: FMNH 54566 (CM 3197). Brazil: Pará.  
Paratypes: 2. Same locality.  
APTERONOTIDAE
- Porotergus gymnotus** Ellis, 1912  
Mem. Carnegie Mus., 5: 441.  
Holotype: FMNH 53575 (CM 1759). Guyana: Amatuk.  
Paratype. Guyana: Konawaruk.  
APTERONOTIDAE
- Priapichthys annectens hesperis** Hubbs, 1924  
Misc. Publ. Mus. Zool. Univ. Mich., 13: 22.  
Holotype: FMNH 14106. Costa Rica: San José.  
Paratypes: 630. Costa Rica: 6 localities.  
POECILIIDAE
- Priapichthys fosteri** Hildebrand, 1925  
Bull. Bur. Fish., 41: 260.  
Paratypes: 18. El Salvador: 2 localities.  
POECILIIDAE
- Priapichthys letonai** Hildebrand, 1925  
Bull. Bur. Fish., 41: 258.  
Paratypes: 45. El Salvador: 5 localities.  
POECILIIDAE
- Priapichthys olomina** Meek, 1914  
Field Mus. Nat. Hist. Zool. Ser., 10(10): 114.  
Holotype: FMNH 7827. Costa Rica: Orotina.  
Allotype and paratypes: 132. Costa Rica: 4 localities.  
The 16 "Pacific Side" paratypes which Grey (1947, p. 178) noted as MISSING are probably among the 7 lots (29 specimens) collected by Alfaro that are now in our main collection as *Brachyrhaphis olomina*.  
POECILIIDAE
- Priapichthys panamensis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 322.  
Holotype: FMNH 8950. Panama: Chame Point.  
Allotype and paratypes: 17. Panama: 2 localities.  
POECILIIDAE
- Prionodes baldwini** Evermann and Marsh, 1899  
Rept. U.S. Fish Comm. 1899, 25: 353.  
Paratypes: 5. Puerto Rico: 2 localities.  
SERRANIDAE
- Prionotus aspersus** Meek and Hildebrand, 1928  
Field Mus. Nat. Hist. Zool. Ser., 15(3): 856.  
Paratypes: 3. Panama: 3 localities.  
TRIGLIDAE
- Pristella aubynei** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 24.  
Holotype: FMNH 52698 (CM 1042). Guyana: Lama Stop-off.  
Paratypes: 47. Guyana: 3 localities.  
CHARACIDAE
- Pristipomoides freemani** Anderson, 1966  
Bull. Mar. Sci., 16(4): 816.  
Paratype. Caribbean Sea: off Colombia.  
LUTJANIDAE
- Pristiurus sauteri** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 160.  
Holotype: FMNH 55193 (CM 219). Taiwan: Takao.  
Paratype. Same locality.  
SCYLIORHINIDAE

- Probolodus heterostomus** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 164.  
Holotype: FMNH 54328 (CM 2973). Brazil: Campos.  
Paratypes: 5. Brazil: 2 localities.  
CHARACIDAE
- Prochilodus maripicru** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 271.  
Holotype: FMNH 53597 (CM 2066). Guyana: Maripicru Creek.  
Paratype. Same locality.  
PROCHILODONTIDAE
- Profundulus candalarius** Hubbs, 1924  
Misc. Publ. Mus. Zool. Univ. Mich., 13: 15.  
Holotype: FMNH 9071. Guatemala: spring at Candalaria.  
Allotype. Same locality.  
CYPRINODONTIDAE
- Protomyzon aphelocheilus** Inger and Chin, 1962  
Fieldiana Zool., 45: 110.  
Holotype: FMNH 68166. Malaysia: Sabah, Sungei Kaingeran.  
Paratypes: 9. Same locality.  
HOMALOPTERIDAE
- Psalidodon gymnodontus** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 166.  
Holotype: FMNH 54574 (CM 3204). Brazil: Rio Iguaçu at Pôrto União.  
CHARACIDAE
- Psectrogaster curviventris** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 509.  
Paratype. Paraguay: Asunción.  
CURIMATIDAE
- Pseudancistrus carnegiei** Eigenmann, 1916  
Ann. Carnegie Mus., 10(1-2): 85.  
Holotype: FMNH 58350 (CM 7346). Colombia: Río San Gil, Santander.  
Paratypes: 5. Colombia: Quebrada de Honda.  
LORICARIIDAE
- Pseudancistrus nigrescens** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 234.  
Holotype: FMNH 53105 (CM 1539). Guyana: Amatuk.  
LORICARIIDAE
- Pseudancistrus pediculatus** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 679.  
Holotype: FMNH 58352 (CM 7348). Colombia: Río Negro at Villavicencio.  
Paratypes: 7. Colombia: 2 localities.  
LORICARIIDAE
- Pseudanos irinae** Winterbottom, 1980  
Life Sci. Contrib. R. Ont. Mus., 123: 27.  
Paratypes: 3. Guyana: 3 localities.  
ANOSTOMIDAE
- Pseudaspius atrilatus** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 231.  
Holotype: FMNH 57075 (CM 6012). Japan: Lake Biwa at Matsubara.  
CYPRINIDAE
- Pseudaspius bergi** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 22.  
Holotype: FMNH 55800 (CM 4563). Korea: Chinnampo.  
Paratypes: 2. Same locality.  
CYPRINIDAE
- Pseudaspius modestus** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 23.  
Holotype: FMNH 55799 (CM 4561). Korea: Chinnampo.  
Paratypes: 4. Same locality.  
CYPRINIDAE
- Pseudauschenipterus nigrolineatus** Fowler, 1911  
Proc. Acad. Nat. Sci. Phila., 63: 434.  
Paratype. Venezuela: Pedernales.  
AUCHENIPTERIDAE
- Pseudobagrus adiposalis** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 181.  
Holotype: FMNH 59079 (CM 8216). Taiwan: Tamusui River near Shinten.  
BAGRIDAE
- Pseudobagrus taiwanensis** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 180.  
Holotype: FMNH 59078 (CM 8215). Taiwan: Tozen River near Taichu.  
BAGRIDAE
- Pseudocheiroidon affinis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 275.  
Holotype: FMNH 8944. Panama: Río Gatún, Monte Liria.  
Paratypes: 251. Panama: 13 localities.  
Published number FMNH 8941 is an error.  
CHARACIDAE



**Pseudochromis aurea marshallensis** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 392.  
Paratypes: 2. Marshall Islands: 2 localities.  
PSEUDOCROMIDAE

**Pseudocorynopoma heterandria** Eigenmann, 1914.  
Indiana Univ. Stud., 20: 39.  
Holotype: FMNH 56407 (CM 5222). Brazil:  
Xiririca.  
Paratypes: 8. Same locality.  
CHARACIDAE

**Pseudogobio chinssuensis** Nichols, 1926  
Am. Mus. Nov., 214: 3.  
Paratype. China: Shansi, Chin-ssu.  
CYPRINIDAE

**Pseudoperilampus hondae** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 20.  
Holotype: FMNH 55803 (CM 4566). Korea:  
Suigen.  
CYPRINIDAE

**Pseudopimelodus albomarginatus** Eigenmann,  
1912  
Mem. Carnegie Mus., 5: 153.  
Holotype: FMNH 53221 (CM 1680). Guyana:  
Tukeit.  
Paratypes: 10. Guyana: 2 localities.  
PIMELODIDAE

**Pseudopimelodus villosus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 152.  
Holotype: FMNH 53219 (CM 1677). Guyana:  
Potaro Landing.  
Paratypes: 2. Guyana: 2 localities.  
PIMELODIDAE

**Pseudoplesiops sargenti** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 405.  
Paratype. Philippine Islands: Endeavor Straits.  
PSEUDOCROMIDAE

**Pseudoraja fischeri** Bigelow and Schroeder, 1954  
Breviora, 24: 4.  
Paratype. Gulf of Mexico: near Campeche Bank.  
RAJIDAE

**Pseudorasbora altipinna** Nichols, 1925  
Am. Mus. Nov., 182: 5.  
Paratypes: 2. China: Szechwan, Yen-ching-kao.  
CYPRINIDAE

**Pseudorasbora depressirostris** Nichols, 1925  
Am. Mus. Nov., 182: 5.  
Paratypes: 2. China: Shansi, Chin-ssu.  
CYPRINIDAE

**Pseudotatia parva** Mees, 1974  
Zool. Verh. (Leiden), 132: 105.  
Holotype: FMNH 70580. Brazil: Joazeiro.  
Paratypes: 13. Same locality.  
AUCHENIPTERIDAE

**Pseudotolithus brunneolus** Jordan and Richardson,  
1909  
Mem. Carnegie Mus., 4(4): 191.  
Holotype: FMNH 52174 (CM 327). Taiwan:  
Takao.  
Paratype. Same locality.  
SCIAENIDAE

**Pseudupeneus chrysonemus** Jordan and Evermann,  
1903  
Bull. U.S. Fish Comm. 1902, 22: 186.  
Paratypes: 2. Hawaiian Islands: Hilo.  
MULLIDAE

**Psilorhynchus gracilis** Rainboth, 1983  
Proc. Calif. Acad. Sci., 43(6): 69.  
Paratypes: 5. Bangladesh: Mahananda River at  
Tetulia.  
PSILORHYNCHIDAE

**Psychichthys eidolon** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 117.  
Holotype: FMNH 59519 (CM 7779). Japan: Sa-  
gami Bay, Izu off Mishima.  
CHIMAERIDAE

**Pterobrycon landoni** Eigenmann, 1913  
Indiana Univ. Stud., 18: 3.  
Holotype: FMNH 56250 (CM 5051). Colombia:  
Boca de Raspadura.  
CHARACIDAE

**Pterois elliotti** Meek, 1897  
Field Mus. Nat. Hist. Zool. Ser., 1(8): 169.  
Holotype: FMNH 484. Somali: Berbera.  
SCORPAENIDAE

**Pterolebias hoignei** Thomerson, 1974  
Copeia, 1974(1): 31.  
Paratypes: 9. Venezuela: Río Pao near Caño  
Benito.  
APLOCHEILIDAE

**Pterolebias maculipinnis** Weibezahn, 1964  
 In: Radda, Die Aquar. un Terr. Zeitschrift,  
 17: 39–41 (not seen).  
 Neotype: FMNH 85267. Venezuela: Río Pao  
 drainage, Cojedes.  
 Neotype designated by Taphorn and Thom-  
 erson (1978, Acta Biol. Venez., 9(4): 395);  
 perhaps authorship should be attributed  
 to Radda (1964).

APLOCHEILIDAE

**Pterygoplichthys anisitsi** Eigenmann and Kenne-  
 dy, 1903  
 Proc. Acad. Nat. Sci. Phila., 59: 503.  
 Paratype. Paraguay: Asunción.  
 Misspelled *Pteryophlichthys* in Eigenmann and  
 Kennedy (1903).  
 LORICARIIDAE

**Puntius joaquinae** Wood, 1968  
 Philipp. J. Sci., 95(4): 415.  
 Holotype: FMNH 73395. Philippine Islands:  
 Mindanao, Agus River.  
 Paratypes: 5. Philippine Islands: 4 localities.  
 CYPRINIDAE

**Puntius snyderi** Oshima, 1919  
 Ann. Carnegie Mus., 12(2–4): 216.  
 Holotype: FMNH 59093 (CM 8231). Taiwan:  
 Nanto, Rigyokutsu.  
 CYPRINIDAE

**Puntius umalii** Wood, 1968  
 Philipp. J. Sci., 95(4): 412.  
 Holotype: FMNH 73393. Philippine Islands:  
 Mindanao, Agus River.  
 Paratypes: 12. Philippine Islands: 5 localities.  
 CYPRINIDAE

**Pygidium alternatum** Eigenmann, 1917  
 Proc. Am. Philos. Soc., 56(8): 700.  
 Holotype: FMNH 58082 (CM 7079). Brazil: Rio  
 Doce.  
 Paratypes: 62. Same locality.  
 TRICHOMYCTERIDAE

**Pygidium banneaui** Eigenmann, 1912  
 Indiana Univ. Stud., 16: 19.  
 Holotype: FMNH 56025 (CM 4815). Colombia:  
 Bernal Creek near Honda.  
 Paratypes: 54. Same locality.  
 TRICHOMYCTERIDAE

**Pygidium barbouri** Eigenmann, 1911  
 Ann. Carnegie Mus., 7(2): 214.  
 Paratypes: 5. Bolivia: upper Beni River.  
 TRICHOMYCTERIDAE

**Pygidium bogotense** Eigenmann, 1912  
 Indiana Univ. Stud., 16: 18.  
 Holotype: FMNH 56030 (CM 4820). Colombia:  
 Chapinero, near Bogotá.  
 Paratypes: 131. Colombia: 2 localities.  
 TRICHOMYCTERIDAE

**Pygidium caliense** Eigenmann, 1912  
 Indiana Univ. Stud., 16: 18.  
 Holotype: FMNH 56029 (CM 4819). Colombia:  
 Cali.  
 Published number CM 6819 is an error.  
 TRICHOMYCTERIDAE

**Pygidium chapmani** Eigenmann, 1912  
 Indiana Univ. Stud., 16: 18.  
 Holotype: FMNH 56027 (CM 4817). Colombia:  
 Boquia.  
 Paratypes: 26. Same locality.  
 TRICHOMYCTERIDAE

**Pygidium conradi** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 212.  
 Holotype: FMNH 53721 (CM 2212). Guyana:  
 Amatuk.  
 TRICHOMYCTERIDAE

**Pygidium davisii** Haseman, 1911  
 Ann. Carnegie Mus., 7(3–4): 380.  
 Holotype: FMNH 60309 (CM 2862). Brazil:  
 Serrinha Paraná.  
 Paratypes: 9. Brazil: creek near Serrinha Paraná.  
 TRICHOMYCTERIDAE

**Pygidium dorsostriatum** Eigenmann, 1917  
 Proc. Am. Philos. Soc., 56(8): 695.  
 Holotype: FMNH 58096 (CM 7093). Colombia:  
 Villavicencio.  
 Paratype. Same locality.  
 Species name was spelled *dorsotriatum* in Ei-  
 genmann (1917), apparently a typo-  
 graphic error emended in subsequent pa-  
 pers by Eigenmann.  
 TRICHOMYCTERIDAE

**Pygidium gracilior** Eigenmann, 1912  
 Mem. Carnegie Mus., 5: 213.  
 Holotype: FMNH 53264 (CM 1730). Guyana:  
 Erukin.  
 TRICHOMYCTERIDAE

- Pygidium guianensis** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 11.  
Holotype: FMNH 52676 (CM 1003). Guyana: Aruataima Falls.  
TRICHOMYCTERIDAE
- Pygidium hasemani** Eigenmann, 1914  
Indiana Univ. Stud., 20: 48.  
Holotype: FMNH 56424 (CM 5238). Brazil: Santarém.  
Paratypes: 312. Same locality.  
TRICHOMYCTERIDAE
- Pygidium iheringi** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 697.  
Holotype: FMNH 58074 (CM 7071). Brazil: São Paulo, Sapina.  
Paratypes: 2. Same locality.  
TRICHOMYCTERIDAE
- Pygidium latistriatum** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 696.  
Holotype: FMNH 58449 (CM 7450). Colombia: Quebrada de Pinchote, Santander.  
TRICHOMYCTERIDAE
- Pygidium paolence** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 698.  
Holotype: FMNH 58085 (CM 7081). Brazil: Rio Tietê at Alto da Serra.  
Paratype. Brazil: Rio Parahyba.  
TRICHOMYCTERIDAE
- Pygidium proops parahybae** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 332.  
Holotype: FMNH 58576 (CM 7598). Brazil: Rio Parahyba at São João da Barra.  
TRICHOMYCTERIDAE
- Pygidium punctulatum piurae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 63.  
Paratypes: 6. Peru: 3 localities.  
TRICHOMYCTERIDAE
- Pygidium reinhardti** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 699.  
Holotype: FMNH 58081 (CM 7078). Brazil: Rio Itabira at Burmier.  
TRICHOMYCTERIDAE
- Pygidium santaeritae** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 341.  
Holotype: FMNH 58577 (CM 7599). Brazil: Rio Preto at Santa Rita.  
TRICHOMYCTERIDAE
- Pygidium septentrionale** Behre, 1928  
Ann. Carnegie Mus., 18(2): 309.  
Holotype: FMNH 59522 (CM 8515). Panama: Salao Creek into Río Chiriquí del Tire.  
Paratype. Panama: Sombrero Creek.  
TRICHOMYCTERIDAE
- Pygidium stellatum** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 308.  
Holotype: FMNH 58101 (CM 7097). Colombia: Quebrada Sarjento.  
Paratypes: 3. Same locality.  
TRICHOMYCTERIDAE
- Pygidium straminium** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 694.  
Holotype: FMNH 58105 (CM 7101). Colombia: Quebrada del Mango.  
Paratypes: 4. Colombia: 2 localities.  
TRICHOMYCTERIDAE
- Pygidium striatum** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 78.  
Holotype: FMNH 7579. Panama: Río Cana at Cana.  
Paratypes: 18. Same locality.  
TRICHOMYCTERIDAE
- Pygidium triguttatum** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 339.  
Holotype: FMNH 58670 (CM 7600). Brazil: Jacarehy.  
Paratypes: 3. Same locality.  
TRICHOMYCTERIDAE
- Pygidium vermiculatum** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 699.  
Holotype: FMNH 58077 (CM 7074). Brazil: Juiz de Fora.  
TRICHOMYCTERIDAE
- Pygidium zonatum** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 330.  
Holotype: FMNH 58573 (CM 7596). Brazil: Agua Quente.  
Paratypes: 3. Brazil: 2 localities.  
TRICHOMYCTERIDAE
- Pygocentrus bilineatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 47.  
Holotype: FMNH 58075 (CM 1072). Guyana: Aruka River.  
Paratype. Same locality.  
CHARACIDAE

- Pyrrhulina australe** Eigenmann and Kennedy, 1903  
Proc. Acad. Nat. Sci. Phila., 59: 508.  
Paratypes: 2. Paraguay: Arroyo Trementina.  
LEBIASINIDAE
- Quisquilius eugenius** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 203.  
Paratype: MISSING. Hawaiian Islands: Honolulu.  
GOBIIDAE
- Rabula fuscomaculata** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 147.  
Paratypes: 2. Marshall Islands: Bikini Atoll, 2 localities.  
MURAENIDAE
- Rabula marshallensis** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 149.  
Paratype: Marshall Islands: Kwajalein Atoll, Ennylabegan Island.  
MURAENIDAE
- Rachovia pyropunctata** Taphorn and Thomerson, 1978  
Acta Biol. Venez., 9(4): 403.  
Holotype: FMNH 85713. Venezuela: west of Lago de Maracaibo, near San Felipe.  
Allotype and paratypes: 30. Same locality.  
APLOCHEILIDAE
- Raja hollandi** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 163.  
Holotype: FMNH 52101 (CM 224). Taiwan: Takao.  
RAJIDAE
- Rhamdella montana** Eigenmann, 1913  
Ann. Carnegie Mus., 8(3-4): 421.  
Holotype: FMNH 56067 (CM 4858). Peru: Queta, brook southeast of Tarma.  
Paratype. Same locality.  
PIMELODIDAE
- Rhamdia barbata** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(4): 106.  
Holotype: FMNH 5906. Nicaragua: lagoon at San Francisco.  
Paratypes: 8. Nicaragua: 2 localities.  
PIMELODIDAE
- Rhamdia branneri** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 377.  
Holotype: FMNH 54235 (CM 2851). Brazil: creek near Serrinha Paraná.  
Paratypes: 2. Same locality.  
PIMELODIDAE
- Rhamdia branneri voulezi** Haseman, 1911  
Ann. Carnegie Mus., 7(3-4): 378.  
Holotype: FMNH 54238 (CM 2854). Brazil: Rio Iguaçu at Pôrto União da Victoria.  
Paratypes: 2. Same locality.  
PIMELODIDAE
- Rhamdia cabreræ** Meek, 1906  
Field Mus. Nat. Hist. Zool. Ser., 7(3): 93.  
Holotype: FMNH 5500. Guatemala: outlet of Lake Amatitlán.  
Paratypes: 49. Same locality.  
PIMELODIDAE
- Rhamdia laticauda typhla** Greenfield, Greenfield, and Woods, 1982  
Brenesia, 19/20: 564.  
Holotype: FMNH 71605. Belize: cave in Mountain Pine Ridge.  
Paratypes: 12. Same locality.  
PIMELODIDAE
- Rhamdia luigiana** Villa, 1977  
Brenesia, 12/13: 133.  
Paratypes: 4. Nicaragua: Lago Nicaragua.  
Villa (1977) lists 2 FMNH paratypes without catalogue number; we have 4 specimens from the type locality that were labeled paratypes.  
PIMELODIDAE
- Rhamdia microps** Eigenmann, 1917  
Ann. Carnegie Mus., 11(3-4): 394.  
Holotype: FMNH 58285 (CM 7283). Brazil: Uruguayana.  
Paratypes: 3. Same locality.  
PIMELODIDAE
- Rhamdia nasuta** Meek, 1909  
Field Mus. Nat. Hist. Zool. Ser., 7(7): 207.  
Holotype: FMNH 6480. Costa Rica: Buenos Aires de Térraba.  
PIMELODIDAE
- Rhamdia oaxacæ** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 74.  
Holotype: FMNH 3716. Mexico: Cuicatlán, Río Quitepec.  
Paratypes: 9. Same locality.  
PIMELODIDAE
- Rhamdia regani** Meek, 1907  
Field Mus. Nat. Hist. Zool. Ser., 7(5): 144.  
Holotype: FMNH 6019. Costa Rica: Turrialba.  
Paratypes: 9. Costa Rica: 2 localities.  
PIMELODIDAE

- Rhamdiopsis moreirai** Haseman, 1911  
Ann. Carnegie Mus., 7(3–4): 375.  
Holotype: FMNH 54372 (CM 2849). Brazil: Rio Iguaçu at Serrinha Paraná.  
Paratype: Same locality.  
PIMELODIDAE
- Rhaphiobelone robusta** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 164.  
Paratype: Philippine Islands: Luzon, Port Dupon.  
BELONIDAE
- Rhegma bermudensis** Kanazawa, 1952  
Fieldiana Zool., 34(7): 82.  
Holotype: FMNH 48924. Bermuda: south side beaches.  
Paratypes: 4. Bermuda and The Bahamas: 2 localities.  
GRAMMISTIDAE
- Rhigophila dearborni** DeWitt, 1962  
Copeia, 1962(4): 821.  
Paratypes: 6. Antarctica: McMurdo Sound.  
ZOARCIDAE
- Rhinodoras boehlkei** Glodek, Whitmire, and Orcés, 1976  
Fieldiana Zool., 70(1): 3.  
Holotype: FMNH 79203. Ecuador: Río Bobonaza.  
DORADIDAE
- Rhinodoras thomersoni** Taphorn and Lilyestrom, 1984  
Rev. UNELLEZ Cienc. Tec., 2: 87.  
Paratypes: 3. Venezuela: mouth of Río Catatumbo, near El Congo.  
DORADIDAE
- Rhinogobius formosanus** Oshima, 1919  
Ann. Carnegie Mus., 12(2–4): 300.  
Holotype: FMNH 59135 (CM 8273). Taiwan: Shinchiku.  
GOBIIDAE
- Rhinogobius mowbrayi** Bean, 1906  
Field Mus. Nat. Hist. Zool. Ser., 7(2): 81.  
Holotype: FMNH 5030. Bermuda: Challenger Bank.  
GOBIIDAE
- Rhinogobius taiwanus** Oshima, 1919  
Ann. Carnegie Mus., 12(2–4): 298.  
Holotype: FMNH 59134 (CM 8272). Taiwan: Shinchiku.  
GOBIIDAE
- Rhinosardinia serrata** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 445.  
Holotype: FMNH 53927 (CM 2443). Guyana: Morawhanna.  
Paratypes: 215. Guyana: 2 localities.  
CLUPEIDAE
- Rhizoprionodon oligolinx** Springer, 1964  
Proc. U.S. Natl. Mus., 115(3493): 521.  
Paratypes: 3. Singapore, Java and Sri Lanka: 3 localities.  
CARCHARHINIDAE
- Rhoadsia altipinna** Fowler, 1911  
Proc. Acad. Nat. Sci. Phila., 63: 498.  
Paratype: Ecuador: tributary of Chimbo River.  
CHARACIDAE
- Rhodeus chosenicus** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 19.  
Holotype: FMNH 55804 (CM 4567). Korea: Suigen.  
Paratypes: 2. Same locality.  
CYPRINIDAE
- Rhodeus kurumeus** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 229.  
Paratype: Japan: Chikugo River at Kurume.  
CYPRINIDAE
- Rhombatractus affinis** Weber, 1908  
Nova Guinea, 5(2): 234.  
Syntypes: 2, FMNH 52394 (CM 650). New Guinea: Timena River.  
MELANOTAENIIDAE
- Rhombatractus sentaniensis** Weber, 1908  
Nova Guinea, 5(2): 235.  
Syntypes: 4, FMNH 52393 (CM 649). New Guinea: Lake Sentani.  
MELANOTAENIIDAE
- Rhombosoma sepikensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 400.  
Holotype: FMNH 17265. New Guinea: Sepik River at Marienberg.  
Paratypes: 6. Same locality.  
MELANOTAENIIDAE
- Rhynchodoras woodsi** Glodek, 1976  
Copeia, 1976(1): 44.  
Holotype: FMNH 77008. Ecuador: Río Bobonaza at Moreta.  
Paratypes: 2 MISSING. Same locality.  
DORADIDAE

**Rineloricaria formosa** Isbrucker and Nijssen, 1979  
Bijdr. Dierkd., 48(2): 192.  
Holotype: FMNH 83713. Colombia: Río Orinoco basin, lagoon near Puerto Inírida.  
Paratypes: 30. Colombia: 2 localities.  
LORICARIIDAE

**Rivulus breviceps** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 49.  
Holotype: FMNH 52710 (CM 1075). Guyana: Shrimp Creek.  
Paratype: MISSING. Same locality.  
APLOCHEILIDAE

**Rivulus brunneus** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 86.  
Holotype: FMNH 7593. Panama: Canal Zone, Creek at Toro Point.  
Paratypes: 7. Same locality.  
APLOCHEILIDAE

**Rivulus compressus** Henn, 1916  
Ann. Carnegie Mus., 10(1-2): 111.  
Holotype: FMNH 57003 (CM 5819). Brazil: Manaus.  
APLOCHEILIDAE

**Rivulus frenatus** Eigenmann, 1912  
Ann. Carnegie Mus., 5: 51.  
Holotype: FMNH 53538 (CM 1085). Guyana: Essequibo River at Gluck Island.  
APLOCHEILIDAE

**Rivulus holmiae** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 50.  
Holotype: FMNH 53535 (CM 1076). Guyana: creek near Holmia.  
Paratypes: 5. Same locality.  
APLOCHEILIDAE

**Rivulus lanceolatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 51.  
Holotype: FMNH 53537 (CM 1084). Guyana: Rockstone.  
APLOCHEILIDAE

**Rivulus magdalenae** Eigenmann and Henn, 1916  
In: Henn, Ann. Carnegie Mus., 10(1-2): 109.  
Holotype: FMNH 56997 (CM 5813). Colombia: Ibagué.  
Paratypes: 32. Colombia: 5 localities.  
APLOCHEILIDAE

**Rivulus montium** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 319.  
Paratypes: 2. Panama: between Río Boquerón and Río Pequeni.  
APLOCHEILIDAE

**Rivulus stagnatus** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 50.  
Holotype: FMNH 52715 (CM 1082). Guyana: Christianburg.  
Paratypes: 4. Same locality.  
APLOCHEILIDAE

**Rivulus stellifer** Thomerson and Turner, 1973  
Copeia, 1973(4): 783.  
Paratypes: 5. Venezuela: Cojedes near Caño Benito.  
APLOCHEILIDAE

**Rivulus vulcanus** Hildebrand, 1938  
Field Mus. Nat. Hist. Zool. Ser., 22(4): 316.  
Paratypes: 10. Panama: Laguna Grande (Davis), Chiriquí.  
APLOCHEILIDAE

**Rivulus waimacui** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 50.  
Holotype: FMNH 53536 (CM 1078). Guyana: Shrimp Creek.  
Paratypes: 5. Guyana: 3 localities.  
APLOCHEILIDAE

**Roeboides cauae** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 162.  
Paratypes: 298. Colombia: 3 localities.  
CHARACIDAE

**Roeboides hildebrandi** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 162.  
Holotype: FMNH 56884 (CM 5186). Colombia: Istmina.  
Paratypes: 44. Colombia: 6 localities.  
CHARACIDAE

**Roeboides macrolepis** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 84.  
Holotype: FMNH 7590. Panama: Río Cupe, Boca de Cupe.  
Paratypes: 10. Panama: 2 localities.  
CHARACIDAE

**Roeboides magdalenae** Eigenmann, 1922

Mem. Carnegie Mus., 9(1): 161.

Syntypes: 201, FMNH 56386–56395 (CM 5202–5210). Colombia: Soplaviento.

Eigenmann (1922, p. 161) indicated that CM 5202 and IUM 12950 (now at California Academy of Sciences) were the “type and paratypes” but apparently never separated, labeled or figured the holotype.

CHARACIDAE

**Roeboides meeki** Eigenmann, 1922

Mem. Carnegie Mus., 9(1): 163.

Holotype: FMNH 55149 (CM 3849). Colombia: Quibdó.

Paratypes: 59. Colombia: 3 localities.

CHARACIDAE

**Roeboides occidentalis** Meek and Hildebrand, 1916

Field Mus. Nat. Hist. Zool. Ser., 10(15): 293.

Holotype: FMNH 8948 MISSING. Panama: Canal Zone, Río Cárdenas at Corozal.

Paratypes: 223. Panama: 21 localities.

CHARACIDAE

**Roeboides salvadoris** Hildebrand, 1925

Bull. Bur. Fish., 41: 246.

Paratypes: 21. El Salvador: 3 localities.

CHARACIDAE

**Roeboides thurni** Eigenmann, 1912

Mem. Carnegie Mus., 5: 399.

Holotype: FMNH 53676 (CM 2149). Guyana: Rockstone sandbank.

Paratypes: 7. Guyana: 2 localities.

CHARACIDAE

**Sagamia russula** Jordan and Snyder, 1901

Proc. U.S. Natl. Mus., 24(1244): 100.

Paratypes: 10. Japan. Nagasaki.

GOBIIDAE

**Salarias martini** Herre, 1942

Proc. Biol. Soc. Wash., 55: 2.

Paratypes: 7. Philippine Islands: Panay, Estancia.

BLENNIIDAE

**Salarias walensis** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 437.

Holotype: FMNH 17393. New Hebrides Islands: Wala Island.

Paratype. Same locality.

BLENNIIDAE

**Salmo bathoecetor** Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 227.

Holotype: FMNH 2035. Washington: Lake Crescent.

Paratype. Same locality.

SALMONIDAE

**Salmo clarki declivifrons** Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 230.

Holotype: FMNH 2006. Washington: Lake Southerland.

SALMONIDAE

**Salmo clarki jordani** Meek, 1899

Field Mus. Nat. Hist. Zool. Ser., 1(12): 229.

Holotype: FMNH 2012. Washington: Lake Southerland.

Paratypes: 18. Same locality.

SALMONIDAE

**Salmo roosevelti** Evermann, 1906

Bull. Bur. Fish. 1905, 25: 26.

Paratype. California: Volcáno Creek.

SALMONIDAE

**Salvelinus imbrius** Jordan and McGregor, 1925

In: Jordan and Hubbs, Mem. Carnegie Mus., 10(2): 142.

Holotype: FMNH 58692 (CM 7797). Japan: Iwami, stream near Hamada.

SALMONIDAE

**Salvelinus timagamiensis** Henn and Rinkenbach, 1925

Ann. Carnegie Mus., 16: 131.

Holotype: FMNH 70514 (CM 7969). Canada: Ontario, White Pine Lake.

Paratypes: 11. Same locality.

SALMONIDAE

**Sarcocheilichthys morii** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 175.

Holotype: FMNH 58718 (CM 7824). Korea: Ping-yang River.

CYPRINIDAE

**Sardinella marquesensis** Berry and Whitehead, 1968

Proc. Biol. Soc. Wash., 81: 209.

Paratypes: 2. Marquesas Islands: 2 localities.

CLUPEIDAE

**Sargocentron macrosquamis** Golani, 1984

Cybius, 8(2): 40.

Holotype: FMNH 86961. Amirantes Islands: eastern end of D'Arros Island.

HOLOCENTRIDAE

- Scaphesthes tamusuiensis** Oshima, 1919  
Ann. Carnegie Mus., 12(2-4): 209.  
Holotype: FMNH 59091 (CM 8228). Taiwan:  
Tamusui River near Shinten.  
CYPRINIDAE
- Scarus lauia** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 196.  
Paratype. Hawaiian Islands: Honolulu.  
SCARIDAE
- Scepterias fragilis** Jordan and Jordan, 1922  
Mem. Carnegie Mus., 10(1): 45.  
Holotype: FMNH 55204 (CM 3900). Hawaiian  
Islands: Honolulu Market.  
APOGONIDAE
- Schizodontopsis laticeps** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 229.  
Holotype: FMNH 53359 (CM 1825). Guyana:  
Crab Falls.  
ANOSTOMIDAE
- Sciaena bathytatos** Chao and Miller, 1975  
Bull. Mar. Sci., 25(2): 267.  
Paratype. Caribbean Sea: off Venezuela.  
SCIAENIDAE
- Sciaena iharae** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 37.  
Holotype: FMNH 55629 (CM 4371). Korea:  
Fusan.  
SCIAENIDAE
- Sciaena trewavasae** Chao and Miller, 1975  
Bull. Mar. Sci., 25(2): 262.  
Paratype. Caribbean Sea: off Venezuela.  
SCIAENIDAE
- Scleronema operculatum** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 691.  
Holotype: FMNH 58080 (CM 7077). Brazil:  
Cacequy.  
Paratypes: 3. Same Locality.  
TRICHOMYCTERIDAE
- Scolopsis eriomma** Jordan and Richardson, 1909  
Mem. Carnegie Mus., 4(4): 188.  
Holotype: FMNH 52247 (CM 317). Taiwan:  
Takao.  
NEMIPTERIDAE
- Scomberoides formosanus** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 236.  
Holotype: FMNH 59499 (CM 7765). Japan: Kii.  
CARANGIDAE
- Scopelarchus beebe** Rofen, 1963  
Aquatica, Aquatic Res. Inst. Stockton, 3: 3.  
Paratype. Western North Atlantic: off Bermuda.  
SCOPELARCHIDAE
- Scopelarchus sagax** Rofen, 1963  
Aquatica, Aquatic Res. Inst. Stockton, 3: 2.  
Holotype: FMNH 49852. Western North At-  
lantic: off Bermuda.  
SCOPELARCHIDAE
- Scopelengys clarkei** Butler and Ahlstrom, 1976  
Fish. Bull., 74(1): 148.  
Paratype. Central Pacific.  
NEOSCOPELIDAE
- Scorpaenopsis catocala** Jordan and Evermann,  
1903  
Bull. U.S. Fish Comm. 1902, 22: 201.  
Paratype. Hawaiian Islands: Hilo.  
SCORPAENIDAE
- Sebastapistes badiorufus** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 409.  
Holotype: FMNH 17337. Tuamotu Islands:  
Takaroa Island.  
SCORPAENIDAE
- Sebastodes clavilatus** Starks, 1911  
Ann. Carnegie Mus., 7(2): 181.  
Holotype: FMNH 52293 (CM 548). Washing-  
ton: San Juan Islands.  
SCORPAENIDAE
- Sebastodes deani** Starks, 1911  
Ann. Carnegie Mus., 7(2): 178.  
Holotype: FMNH 52290 (CM 545). Washing-  
ton: San Juan Islands.  
SCORPAENIDAE
- Sebastodes emphaeus** Starks, 1911  
Ann. Carnegie Mus., 7(2): 182.  
Holotype: FMNH 52294 (CM 549). Washing-  
ton: San Juan Islands.  
Paratypes: 5. Same locality.  
SCORPAENIDAE



**Sebastodes ijimae** Jordan and Metz, 1913

Mem. Carnegie Mus., 6(1): 49.

Holotype: FMNH 55437 (CM 4172). Korea: Fusan.

SCORPAENIDAE

**Sebastodes owstoni** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 270.

Holotype: FMNH 57088 (CM 6026). Japan: Aomori.

Paratypes: 3. Same locality.

SCORPAENIDAE

**Sebastodes thompsoni** Jordan and Hubbs, 1925

Mem. Carnegie Mus., 10(2): 265.

Paratype: Japan: Osaka.

SCORPAENIDAE

**Serraniculus pumilio** Ginsburg, 1952

J. Wash. Acad. Sci., 42(3): 88.

Paratype. Gulf of Mexico: off Texas.

SERRANIDAE

**Serranus chionaraia** Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 270.

Paratype. Western Caribbean Sea: off Honduras.

SERRANIDAE

**Serranus maytagi** Robins and Starck, 1961

Proc. Acad. Nat. Sci. Phila., 113(11): 288.

Paratype. Western Caribbean Sea: off Honduras.

SERRANIDAE

**Serrasalmo hollandi** Eigenmann, 1915

Ann. Carnegie Mus., 9(3–4): 251.

Holotype: FMNH 56978 (CM 5792). Brazil: Rio Guaporé at Maciel.

CHARACIDAE

**Serrasalmo humeralis gracilior** Eigenmann, 1915

Ann. Carnegie Mus., 9(3–4): 257.

Syntypes: 3, FMNH 56977 (CM 5791). Brazil: Rio Guaporé at Maciel.

CHARACIDAE

**Serrasalmus coccogenis** Fowler, 1911

Proc. Acad. Nat. Sci. Phila., 63: 428.

Paratype. Venezuela: Caño Uracoa at La Pedrita.

CHARACIDAE

**Sicydium altum** Meek, 1907

Field Mus. Nat. Hist. Zool. Ser., 7(5): 149.

Holotype: FMNH 6034. Costa Rica: Turrialba.

Paratypes: 2. Costa Rica: El Guayabo.

GOBIIDAE

**Sicydium hildebrandi** Eigenmann, 1917

Proc. Am. Philos. Soc., 56(7): 685.

Holotype: FMNH 58465 (CM 7466). Colombia:

Río Dagua at Cisnero.

GOBIIDAE

**Sicydium montanum** Hubbs, 1920

Proc. Biol. Soc. Wash., 33: 89.

Holotype: FMNH 9053. Venezuela: Caracas, brook at Macuto.

Paratypes: 10. Same locality.

GOBIIDAE

**Signalosa atchafalayae vanhyningi** Weed, 1925

Field Mus. Nat. Hist. Zool. Ser., 12(11): 145.

Holotype: FMNH 11395. Florida: near Gainesville.

Paratypes: 3. Same locality.

CLUPEIDAE

**Signalosa mexicana campi** Weed, 1925

Field Mus. Nat. Hist. Zool. Ser., 12(11): 143.

Holotype: FMNH 11413. Texas: Brownsville, Resaca de la Guerra.

Paratypes: 493. Texas: 4 localities.

CLUPEIDAE

**Siphamia cuprea** Lachner, 1953

Bull. U.S. Natl. Mus., 202(1): 423.

Paratypes: 2. Sulu Archipelago: Pangasinan Island.

APOGONIDAE

**Siphamia fuscolineata** Lachner, 1953

Bull. U.S. Natl. Mus., 202(1): 425.

Paratypes: 2. Bikini Island.

APOGONIDAE

**Siphostoma brevicaudum** Meek, 1904

Field Mus. Nat. Hist. Zool. Ser., 5: 163.

Holotype: FMNH 4586. Mexico: Veracruz, Boca del Río.

Paratypes: 2. Same locality.

SYNGNATHIDAE

**Siphostoma eigenmanni** Wilson, 1916

Ann. Carnegie Mus., 10(1–2): 60.

Holotype: FMNH 56857 (CM 5672). Brazil: Bahia, Rio Vermelho.

SYNGNATHIDAE

**Siphostoma elcapitanense** Meek and Hildebrand, 1914

Field Mus. Nat. Hist. Zool. Ser., 10(10): 119.  
Paratypes: 4. Panama and Costa Rica: 3 localities.

SYNGNATHIDAE

**Skiffia lermæ** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 102.  
Holotype: FMNH 3616. Mexico: Lago de Patzcuaro at Patzcuaro.

Paratypes: 131. Mexico: 2 localities.

GOODEIDAE

**Skiffia variegata** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 104.  
Holotype: FMNH 3612. Mexico: Lago de Zirahuén at Zirahuén.

Allotype and paratypes: 99. Mexico: 2 localities.  
GOODEIDAE

**Solenostomus paegnius** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 235.

Holotype: FMNH 57083 (CM 6020). Japan: Misaki.

SOLENOSTOMIDAE

**Soleonassus finis** Eigenmann, 1912

Mem. Carnegie Mus., 5: 528.

Holotype: FMNH 53965 (CM 2487). Guyana: Tumatumari.

SOLEIDAE

**Solivomer arenidens** Miller, 1947

Proc. U.S. Natl. Mus., 97(3211): 84.

Paratypes: 2. Philippine Islands: off Balicasag Island.

NEOSCOPELIDAE

**Sonoda megalophthalma** Grey, 1959

Bull. Mus. Comp. Zool. Harv. Coll., 121(4): 181.

Holotype: FMNH 65981. Western Caribbean Sea.

Paratypes: 22. Western Caribbean Sea: 5 localities.

STERNOPTYCHIDAE

**Sovichthys abuelo** Schultz, 1944

Proc. U.S. Natl. Mus., 94(3172): 191.

Paratypes: 4. Venezuela: Río Socuy, above its mouth.

PIMELODIDAE

**Sparisoma rhomaleum** Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 754.

Paratype. Panama: Colón market.

SCARIDAE

**Spectrunculus radcliffei** Jordan and Thompson, 1914

Mem. Carnegie Mus., 6(4): 301.

Holotype: FMNH 57123 (CM 6061). Japan: Misaki.

OPHIDIIDAE

**Sphagebranchus conklini** Eigenmann, 1916

Ann. Carnegie Mus., 10(1-2): 55.

Holotype: FMNH 57731 (CM 6710). The Bahamas: New Providence.

OPHICHTHIDAE

**Spheroides asterias** Blosser, 1909

Ann. Carnegie Mus., 6(1): 300.

Holotype: FMNH 53050 (CM 1475). Lesser Antilles: Saint Croix Island.

TETRAODONTIDAE

**Sphoeroides kendalli** Meek and Hildebrand, 1928

Field Mus. Nat. Hist. Zool. Ser., 15(3): 819.

Paratype. Panama: Chame Point.

TETRAODONTIDAE

**Sphoeroides parvus** Shipp and Yerger, 1969

Proc. Biol. Soc. Wash., 82: 478.

Paratypes: 10. Alabama: Mobile Bay.

TETRAODONTIDAE

**Spinibarbus hollandi** Oshima, 1919

Ann. Carnegie Mus., 12(2-4): 218.

Holotype: FMNH 59095 (CM 8233). Taiwan: Sobun River near Tabani.

CYPRINIDAE

**Spinoblennius spiniger** Herre, 1935

Field Mus. Nat. Hist. Zool. Ser., 18(12): 435.

Holotype: FMNH 17395. Ecuador: Islas Galápagos, Isla Edén.

Paratype. Same locality.

BLENNIIDAE

**Spintherobolus broccae** Myers, 1925

Ann. Carnegie Mus., 16: 143.

Holotype: FMNH 58863 (CM 7979). Brazil: hills behind Rio de Janeiro.

Paratype. Same locality.

CHARACIDAE

**Spintherobolus papilliferus** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 167.  
Holotype: (CM 3582) MISSING. Brazil: São Paulo, Alto da Serra.  
Paratypes: 5. Same locality.  
CHARACIDAE

**Spirinchus dilatus** Schultz and Chapman, 1934  
Ann. Mag. Nat. Hist., Ser. 10, 13: 68.  
Paratype. Washington: Nooksack River.  
OSMERIDAE

**Spirinchus verecundus** Jordan and Metz, 1913  
Mem. Carnegie Mus., 6(1): 11.  
Holotype: FMNH 55807 (CM 4570). Korea: Chinnampo.  
Paratype. Same locality.  
OSMERIDAE

**Squaliolus sarmenti** Noronha, 1926  
Ann. Carnegie Mus., 16(3–4): 385.  
Holotype: FMNH 58862 (CM 7976). Portugal: Madeira Islands, Funchal.  
SQUALIDAE

**Squalogadus intermedius** Grey, 1959  
Fieldiana Zool., 39(29): 330.  
Paratypes: 5. Gulf of Mexico.  
MACROURIDAE

**Starksia occidentalis** Greenfield, 1979  
Fieldiana Zool., 73(2): 27.  
Holotype: FMNH 83716. Belize: Tarpon Cay.  
Paratypes: 41. Belize and Honduras: 11 localities.  
LABRISOMIDAE

**Starksia variabilis** Greenfield, 1979  
Fieldiana Zool., 73(2): 33.  
Holotype: FMNH 83717. Colombia: Santa Marta.  
Paratypes: 7. Same locality.  
LABRISOMIDAE

**Starksia y-lineata** Gilbert, 1965  
Not. Nat. (Phila.), 379: 2.  
Paratypes: 2. British West Indies: Grand Cayman Island.  
LABRISOMIDAE

**Stegastes baldwini** Allen and Woods, 1980  
Rec. West. Aust. Mus., 8(2): 183.  
Paratypes: 54. Clipperton Island.  
POMACENTRIDAE

**Stellifer colonensis** Meek and Hildebrand, 1925  
Field Mus. Nat. Hist. Zool. Ser., 15(2): 623.  
Paratypes: 21. Panama: 4 localities.  
SCIAENIDAE

**Stellifer melanocheir** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 682.  
Holotype: FMNH 58507 (CM 7520). Colombia: Tumaco.  
SCIAENIDAE

**Stellifer mindii** Meek and Hildebrand, 1925  
Field Mus. Nat. Hist. Zool. Ser., 15(2): 626.  
Paratypes: 3. Panama: Canal Zone near Mindi Reef.  
SCIAENIDAE

**Stellistius katsukii** Jordan and Tanaka, 1927  
Ann. Carnegie Mus., 17(12): 389.  
Holotype: FMNH 59186 (CM 8329). Japan: Hokkaido, off Mororan.  
HEXAGRAMMIDAE

**Sternarchus hasemani** Ellis, 1913  
Mem. Carnegie Mus., 6(3): 147.  
Holotype: FMNH 54562 (CM 3191). Brazil: near Santarém.  
Paratypes: 15. Same locality.  
APTERONOTIDAE

**Sternarchus leptorhynchus** Ellis, 1912  
In: Eigenmann, Mem. Carnegie Mus., 5: 439.  
Holotype: FMNH 53294 (CM 1762). Guyana: Amatuk.  
Paratype: Guyana: Warraputa.  
Published number CM 1763 for the holotype is an error.  
APTERONOTIDAE

**Sternarchus mariae** Eigenmann and Fisher, 1914  
Indiana Univ. Stud., 25: 236.  
Holotype: FMNH 56774 (CM 5594). Colombia: Girardot.  
APTERONOTIDAE

**Sternarchus rostratus** Meek and Hildebrand, 1913  
Field Mus. Nat. Hist. Zool. Ser., 10(8): 85.  
Holotype: FMNH 7592. Panama: Río Grande, Cana.  
APTERONOTIDAE

**Sternopygus dariensis** Meek and Hildebrand, 1916  
Field Mus. Nat. Hist. Zool. Ser., 10(15): 309.  
Holotype: FMNH 8949. Panama: Río Tuyra at Marriganti.  
Paratypes: 12. Same locality.  
STERNOPYGIDAE

- Sternopygus pejeraton** Schultz, 1949  
Proc. U.S. Natl. Mus., 99(3235): 60.  
Paratypes: 4. Venezuela: Lago Maracaibo basin,  
Río Apón.  
STERNOPYGIDAE
- Stethaprion crenatus** Eigenmann, 1916  
Ann. Carnegie Mus., 10(1-2): 80.  
Holotype: FMNH 56413 (CM 5228). Bolivia:  
San Joaquín.  
Paratypes: 3. Bolivia and Brazil: 2 localities.  
CHARACIDAE
- Stevardia aliata** Eigenmann, 1914  
Indiana Univ. Stud., 20: 37.  
Holotype: FMNH 56400 (CM 5215). Colombia:  
Río Negro at Villavicencio.  
Paratypes: 13. Colombia: 2 localities.  
CHARACIDAE
- Stizostedion glaucum** Hubbs, 1926  
Misc. Publ. Mus. Zool. Univ. Mich., 15: 58.  
Paratypes: 2. Pennsylvania: Lake Erie off Erie.  
PERCIDAE
- Stolephorus branchiomelas** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(7): 682.  
Holotype: FMNH 58489 (CM 7491). Colombia:  
mouth of Río Dagua.  
Paratypes: 6. Colombia: 2 localities.  
ENGRAULIDAE
- Stolephorus guianensis** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 447.  
Holotype: FMNH 53932 (CM 2448). Guyana:  
Bartica.  
Paratype: Guyana: below Wismar.  
ENGRAULIDAE
- Stolephorus lyolepis** Evermann and Marsh, 1902  
Bull. U.S. Fish Comm. 1900, 20(1): 89.  
Paratypes: 5. Puerto Rico: Culebra.  
ENGRAULIDAE
- Suruga fundicola** Jordan and Snyder, 1901  
Proc. U.S. Natl. Mus., 24(1244): 96.  
Paratypes: 4. Japan: Suruga Bay.  
GOBIIDAE
- Symphysanodon berryi** Anderson, 1970  
U.S. Fish Wildl. Serv. Fish. Bull., 68(2): 338.  
Paratypes: 20. Caribbean Sea: 5 localities.  
SERRANIDAE
- Symphysanodon octoactinus** Anderson, 1970  
U.S. Fish Wildl. Serv. Fish. Bull., 68(2): 342.  
Paratype: Western Caribbean Sea.  
SERRANIDAE
- Synaphobranchus oregoni** Castle, 1960  
Fieldiana Zool., 39(35): 388.  
Paratypes: 6. Gulf of Mexico.  
SYNAPHOBRANCHIDAE
- Synchiropus ijimae** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 295.  
Holotype: FMNH 57078 (CM 6015). Japan:  
Misaki.  
CALLIONYMIDAE
- Synchiropus minutulus** Fricke, 1981  
Theses Zoologicae, 1: 119.  
Holotype: FMNH 78973. Maldives Archipelago:  
southwest of Mafilifleuri.  
CALLIONYMIDAE
- Syngnathus mackayi nesiotes** Herald, 1942  
Stanford Ichthyol. Bull., 2(4): 128.  
Paratypes: 10. Bermuda: 7 localities.  
SYNGNATHIDAE
- Syngnathus scovelli makaxi** Herald and Dawson,  
1972  
Copeia, 1972(4): 781.  
Paratypes: 4. Mexico: Isla Mujeres, Laguna  
Makax.  
SYNGNATHIDAE
- Synodus binotatus** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 35.  
Paratypes: 2. Marshall Islands: 2 localities.  
SYNODONTIDAE
- Synodus ulae** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 38.  
Paratype: Hawaiian Islands: Honolulu Market.  
SYNODONTIDAE
- Taenioides snyderi** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 310.  
Holotype: FMNH 58831 (CM 7941). Japan:  
Wakanoura.  
GOBIOIDIDAE
- Tagusa delicata** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 435.  
Holotype: FMNH 17399. Ecuador: Islas Galápagos,  
Isla Albemarle, Tagus Cove.  
Paratypes: 3. Same locality.  
BLENNIIDAE

- Terapon habbema** Weber, 1910  
Notes Leiden Mus., 32: 234.  
Syntype: FMNH 52496 (CM 754). New Guinea:  
Lorentz River, Alkmaar.  
Spelled *Therapon* in Weber (1910).  
TERAPONIDAE
- Tetragonopterus alleni** Eigenmann and McAtee, 1907  
In: Eigenmann, McAtee, and Ward, Ann.  
Carnegie Mus., 4(2): 126.  
Paratype. Paraguay: Corumbá.  
CHARACIDAE
- Tetraodon kretamensis** Inger, 1953  
Fieldiana Zool., 34(11): 149.  
Holotype: FMNH 51558. Malaysia: Sabah, Pi-  
nang River.  
Paratypes: 56. Malaysia: Sabah, 4 localities.  
TETRAODONTIDAE
- Teuthis atrimentatus** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 198.  
Paratype. Hawaiian Islands: Honolulu.  
ACANTHURIDAE
- Thalassoma cranei** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 412.  
Holotype: FMNH 17347. Solomon Islands: Is-  
abel Island, Tenibuli.  
LABRIDAE
- Thalassophryne quadrizonatus** Eigenmann, 1922  
Mem. Carnegie Mus., 9(1): 217.  
Holotype: (CM 3921) MISSING. Colombia: Río  
Truandó.  
BATRACHOIDIDAE
- Thoracocharax brevis** Eigenmann, 1912  
Indiana Univ. Stud., 16: 25.  
Holotype: FMNH 56055 (CM 4845). Colombia:  
Raspadura.  
GASTEROPELECIDAE
- Thoracocharax magdalenae** Eigenmann, 1912  
Indiana Univ. Stud., 16: 25.  
Holotype: FMNH 56056 (CM 4846). Colombia:  
Girardot.  
GASTEROPELECIDAE
- Thorichthys ellioti** Meek, 1904  
Field Mus. Nat. Hist. Zool. Ser., 5: 223.  
Holotype: FMNH 4627. Mexico: Veracruz,  
Motzorongo.  
Paratypes: 24. Same locality.  
CICHLIDAE
- Thyrina colombiensis** Hubbs, 1920  
Occas. Pap. Mus. Zool. Univ. Mich., 88: 4.  
Paratypes: 49. Colombia: 3 localities.  
ATHERINIDAE
- Thyrina guija** Hildebrand, 1925  
Bull. Bur. Fish., 41: 264.  
Paratypes: 13. El Salvador: 3 localities.  
ATHERINIDAE
- Thyrina meeki** Miller, 1907  
Bull. Am. Mus. Nat. Hist., 23(2): 110.  
Paratypes: 23. Guatemala: 5 localities.  
ATHERINIDAE
- Thysanichthys evides** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 272.  
Holotype: FMNH 57082 (CM 6019). Japan:  
Misaki.  
SCORPAENIDAE
- Tomeurus gracilis** Eigenmann, 1909  
Ann. Carnegie Mus., 6(1): 53.  
Holotype: FMNH 53541 (CM 1093). Guyana:  
Mud Creek in Aruka River.  
Paratypes: 2. Guyana: 2 localities.  
POECILIIDAE
- Toxus riddlei** Eigenmann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 226.  
Allotype: Cuba: San Cristóbal.  
POECILIIDAE
- Trachelyichthys exilis** Greenfield and Glodek, 1977  
Fieldiana Zool., 72(3): 49.  
Holotype: FMNH 80476. Peru: tributary of Río  
Nanay.  
Paratype. Same locality.  
AUCHENIPTERIDAE
- Trachinotus jordani** Wakiya, 1924  
Ann. Carnegie Mus., 15(2–3): 221.  
Holotype: FMNH 59374 (CM 7756). Japan:  
Bonin Islands.  
Spelled *Trachynotus* in Wakiya (1924).  
CARANGIDAE
- Trachipterus fukuzakii** Fitch, 1964  
Calif. Fish Game, 50(4): 236.  
Paratype. Mexico: Baja California, southwest of  
Cabo San Lucas.  
TRACHIPTERIDAE

- Trachurus argenteus** Wakiya, 1924  
Ann. Carnegie Mus., 15(2-3): 145.  
Paratype. Japan: Amakusa, near Nagasaki.  
CARANGIDAE
- Trachycorystes fisheri** Eigenmann, 1916  
Ann. Carnegie Mus., 10(1-2): 82.  
Holotype: FMNH 57695 (CM 6667). Colombia:  
Río Sucio.  
Paratypes: 16. Colombia: 3 localities.  
CARANGIDAE
- Triathalassothia gloverensis** Greenfield and  
Greenfield, 1973  
Copeia, 1973(3): 560.  
Holotype: FMNH 71575. Belize: Glover's Reef.  
Paratypes: 7. Belize: 5 localities.  
BATRACHOIDIDAE
- Trichopsetta caribbaea** Anderson and Guthertz,  
1967  
Bull. Mar. Sci., 17(4): 898.  
Paratypes: 2. Caribbean Sea: off Colombia.  
BOTHIDAE
- Tridentiger kuroiwaie** Jordan and Tanaka, 1927  
Ann. Carnegie Mus., 17(2): 276.  
Holotype: FMNH 59181 (CM 8324). Japan:  
Amami-Oshima.  
Paratypes: 9. Same locality.  
GOBIIDAE
- Tylobronchus maculosus** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 272.  
Holotype: FMNH 53449 (CM 1923). Guyana:  
creek below Potaro Landing.  
Paratypes: 54. Guyana: 6 localities.  
CHILODONTIDAE
- Tympanopleura piperata** Eigenmann, 1912  
Mem. Carnegie Mus., 5: 203.  
Holotype: FMNH 53243 (CM 1708). Guyana:  
Crab Falls.  
Paratype. Same locality.  
AGENEIOSIDAE
- Ulua richardsoni** Jordan and Snyder, 1909  
Mem. Carnegie Mus., 4(2): 39.  
Holotype: FMNH 55365 (CM 413). Taiwan:  
Takao.  
CARANGIDAE
- Umbrina tumacoensis** Wilson, 1916  
Ann. Carnegie Mus., 10(1-2): 67.  
Holotype: FMNH 56840 (CM 5654) MISSING.  
Colombia: Tumaco.  
Paratypes: 2. Same locality.  
SCIAENIDAE
- Upeneus arge** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 187.  
Paratype. Hawaiian Islands: Honolulu.  
MULLIDAE
- Uropterygius fuscoguttatus** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 156.  
Paratype. Marshall Islands: Bikini Atoll, Eman  
Island.  
MURAENIDAE
- Vaimosa balteata** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 419.  
Holotype: FMNH 17386. Indonesia: Waigiu Is-  
land, Majalibit Inlet.  
GOBIIDAE
- Vaimosa horiae** Herre, 1936  
Philipp. J. Sci., 59: 280.  
Syntypes (?): 2, FMNH 47241. Palau Islands:  
freshwater creek on Bab-el-Thaup Island.  
These are from the type locality, but were not  
explicitly cited in the original descrip-  
tion; see comment in Böhlke (1953, Stan-  
ford Ichthyol. Bull., 5(1): 119).  
GOBIIDAE
- Vaimosa osgoodi** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 420.  
Holotype: FMNH 17387. Fiji Islands: Viti Levu  
Island, Suva Harbor.  
GOBIIDAE
- Vandellia hasemani** Eigenmann, 1918  
Mem. Carnegie Mus., 7(5): 363.  
Holotype: FMNH 58523 (CM 7542). Bolivia:  
Río Mamoré.  
Paratypes: 2. Same locality.  
TRICHOMYCTERIDAE
- Vandellia sanguinea** Eigenmann, 1917  
Proc. Am. Philos. Soc., 56(8): 701.  
Holotype: FMNH 58086 (CM 7082). Brazil:  
São Antonio de Rio Madeira.  
TRICHOMYCTERIDAE

- Varicorhinus robustus** Nichols, 1925  
Am. Mus. Nov., 182: 2.  
Paratypes: 2. China: Fukien, Yenping.  
CYPRINIDAE
- Varicus bucca** Robins and Böhlke, 1961  
Copeia, 1961(1): 47.  
Paratype. Antilles Islands: north of Culebra Island.  
GOBIIDAE
- Varicus imswe** Greenfield, 1981  
Copeia, 1981(2): 269.  
Holotype: FMNH 83894. Belize: Carrie-Bow Cay.  
Paratype. Same locality.  
GOBIIDAE
- Vesicatrus tegatus** Eigenmann, 1911  
Ann. Carnegie Mus., 8(1): 174.  
Holotype: (CM 3201) MISSING. Brazil: Upper Rio Paraguay basin, Jaurú.  
Paratypes: 6. Same locality.  
CHARACIDAE
- Vinciguerria mabahiss** Johnson and Feltes, 1984  
Fieldiana Zool., New Ser., 22: 22.  
Paratype. Red Sea: Gulf of Aqaba.  
PHOTICHTHYIDAE
- Wallago maculatus** Inger and Chin, 1959  
Fieldiana Zool., 39(27): 279.  
Holotype: FMNH 68038. Malaysia: Sabah, Kinabatangan River at Deramakot.  
Paratypes: 11. Malaysia: Sabah, 2 localities.  
SILURIDAE
- Xiliphius magdalenae** Eigenmann, 1912  
Indiana Univ. Stud., 16: 10.  
Holotype: FMNH 56039 (CM 4829). Colombia: Girardot.  
Spelled *Xyliphius* and *Xiliphius* in Eigenmann (1912), but spelled as above in subsequent usages by Eigenmann and others.  
ASPREDINIDAE
- Xiphophorus jalapae** Meek, 1902  
Field Mus. Nat. Hist. Zool. Ser., 3(6): 107.  
Holotype: FMNH 3724. Mexico: Veracruz, Jalapa.  
Allotype and paratypes: 110. Same locality.  
POECILIIDAE
- Xyrichthys niveilatus** Jordan and Evermann, 1903  
Bull. U.S. Fish Comm. 1902, 22: 194.  
Paratype. Hawaiian Islands: Honolulu.  
LABRIDAE
- Xyrichthys sciistius** Jordan and Thompson, 1914  
Mem. Carnegie Mus., 6(4): 263.  
Holotype: FMNH 57090 (CM 6028). Japan: Sagami Bay.  
LABRIDAE
- Ypsigramma lineata** Schultz, 1953  
Bull. U.S. Natl. Mus., 202(1): 375.  
Paratype: Marshall Islands: Bikini Atoll, Yuro Island.  
SERRANIDAE
- Zalecopus satsumae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 313.  
Holotype: FMNH 58839 (CM 7949). Japan: Kagoshima Bay.  
URANOSCOPIDAE
- Zalecopus tosae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 312.  
Holotype: FMNH 58835 (CM 7945). Japan: Tosa at Kochi.  
URANOSCOPIDAE
- Zenarchopteris robertsi** Collette, 1982  
Copeia, 1982(2): 266.  
Paratype: Papua New Guinea: tributary of Kumusi River.  
HEMIRAMPHIDAE
- Zenarchopteris sepikensis** Herre, 1935  
Field Mus. Nat. Hist. Zool. Ser., 18(12): 321.  
Holotype: FMNH 17213. New Guinea: Sepik River at Koragu.  
Paratypes: 10. New Guinea: 3 localities.  
HEMIRAMPHIDAE
- Zestichthys tanakae** Jordan and Hubbs, 1925  
Mem. Carnegie Mus., 10(2): 321.  
Holotype: FMNH 58841 (CM 7951). Japan: Kushiro.  
ZOARCIDAE
- Zingel streber nerensis** Banareescu and Nalbant, 1979  
Trav. Mus. D'Hist. Nat. "Grigore Antipa", 20(1): 353.  
Paratype. Romania: Banat, Nera River at Sasca.  
PERCIDAE

**Zoogoneticus diazi** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 93.

Holotype: FMNH 3618. Mexico: Lago de Patzcuaro at Patzcuaro.

Paratypes: 126. Mexico: 2 localities.

GOODEIDAE

**Zoogoneticus miniatus** Meek, 1902

Field Mus. Nat. Hist. Zool. Ser., 3(6): 94.

Holotype: FMNH 3680. Mexico: Lago de Chalco near Chalco.

Paratypes: 4. Same locality.

GOODEIDAE

**Zygonectes pulvereus** Evermann, 1892

Bull. U.S. Fish Comm. 1891, 11: 85.

Paratype. Texas: Corpus Christi.

CYPRINODONTIDAE

## List of Taxa Represented by Types

### Arranged Alphabetically by Family

#### ACANTHOCLINIDAE

*Acanthoplesiops hiatti* Schultz, 1953

#### ACANTHURIDAE

*Acanthurus randalli* Briggs and Caldwell, 1957

*Acanthurus triostegus marquesensis* Schultz and Woods, 1948

*Teuthis atrimentatus* Jordan and Evermann, 1903

#### ACIPENSERIDAE

*Acipenser oxyrhynchus desotoi* Vladikov, 1955

#### AGENEIOSIDAE

*Ageneiosus guianensis* Eigenmann, 1912

*Ageneiosus madeirensis* Fisher, 1917

*Ageneiosus marmoratus* Eigenmann, 1912

*Tympanopleura piperata* Eigenmann, 1912

#### AGONIDAE

*Ithuriella kasawae* Jordan and Hubbs, 1925

#### ALBULIDAE

*Albula virgata* Jordan and Jordan, 1922

#### ALEPOCEPHALIDAE

*Alepocephalus owstoni* Tanaka, 1908

*Alepocephalus umbriceps* Jordan and Thompson, 1914

*Grimatrocetes bullisi* Grey, 1958

#### ANABANTIDAE

*Ctenopoma multispinis* Peters, 1844

#### ANOSTOMIDAE

*Anostomus plicatus* Eigenmann, 1912

*Leporinodus retropinnis* Eigenmann, 1922

*Leporinus alternus* Eigenmann, 1912

*Leporinus arcus* Eigenmann, 1912

*Leporinus brunneus* Myers, 1950

*Leporinus ecuadoriensis* Eigenmann and Henn, 1916

*Leporinus granti* Eigenmann, 1912

*Leporinus steyermarki* Inger, 1956

*Pseudanos irinae* Winterbottom, 1980

*Schizodontopsis laticeps* Eigenmann, 1912

#### ANTENNARIIDAE

*Antennarius bermudensis* Schultz, 1957

*Antennarius pauciradiatus* Schultz, 1957

*Antennarius verrucosus* Bean, 1906

*Phrynelox atra* Schultz, 1957

*Phrynelox zebrinus* Schultz, 1957

#### APLOCHEILIDAE

*Aplocheilus chobensis* Fowler, 1935

*Aplocheilus luzonensis* Herre and Ablan, 1934

*Cynodonichthys tenuis* Meek, 1904

*Pterolebias hoignei* Thomerson, 1974

*Pterolebias maculipinnis* Weibezahn, 1964

*Rachovia pyropunctata* Taphorn and Thomerson, 1978

*Rivulus breviceps* Eigenmann, 1909

*Rivulus brunneus* Meek and Hildebrand, 1913

*Rivulus compressus* Henn, 1916

*Rivulus frenatus* Eigenmann, 1912

*Rivulus holmiae* Eigenmann, 1909

*Rivulus lanceolatus* Eigenmann, 1909

*Rivulus magdalenae* Eigenmann and Henn, 1916

*Rivulus montium* Hildebrand, 1938

*Rivulus stagnatus* Eigenmann, 1909

*Rivulus stellifer* Thomerson and Turner, 1973

*Rivulus volcanus* Hildebrand, 1938

*Rivulus waimacui* Eigenmann, 1909

#### APOGONIDAE

*Amia sialis* Jordan and Thompson, 1914

*Apogonichthys melampodus* Blosser, 1909

*Apogon abo* Herre, 1935

*Apogon beauforti* Weber, 1908

*Apogon leptocaulus* Gilbert, 1972

*Apogon nigrofasciatus* Lachner, 1953

*Apogon snyderi* Jordan and Evermann, 1903



*Apogon wichmanni* Weber, 1908  
*Gymnapogon gracilicauda* Lachner, 1953  
*Gymnapogon urospilotus* Lachner, 1953  
*Mionorus pacificus* Herre, 1935  
*Scepterias fragilis* Jordan and Jordan, 1922  
*Siphamia cuprea* Lachner, 1953  
*Siphamia fuscolineata* Lachner, 1953

#### APTERONOTIDAE

*Adontosternarchus clarkae* Mago-Leccia, Lundberg, and Baskin, 1985  
*Adontosternarchus devenanzii* Mago-Leccia, Lundberg, and Baskin, 1985  
*Porotergus gimbeli* Ellis, 1912  
*Porotergus gymnotus* Ellis, 1912  
*Sternarchus hasemani* Ellis, 1913  
*Sternarchus leptorhynchus* Ellis, 1912  
*Sternarchus mariae* Eigenmann and Fisher, 1914  
*Sternarchus rostratus* Meek and Hildebrand, 1913

#### ARGENTINIDAE

*Argentina alicae* Cohen and Atsaiades, 1969  
*Argentina brucei* Cohen and Atsaiades, 1969  
*Argentina georgei* Cohen and Atsaiades, 1969  
*Argentina stewarti* Cohen and Atsaiades, 1969  
*Glossanodon pygmaeus* Cohen, 1958

#### ARIIDAE

*Arius kanganamanensis* Herre, 1935  
*Arius nox* Herre, 1935  
*Arius solidus* Herre, 1935  
*Arius tuyra* Meek and Hildebrand, 1923  
*Felichthys amblops* Meek and Hildebrand, 1913  
*Galeichthys aguadulce* Meek, 1904  
*Hemipimelodus papillifer* Herre, 1915  
*Hemipimelodus velutinus* Weber, 1908  
*Netuma osakae* Jordan and Kasawa, 1925  
*Netuma planifrons* Higuchi, Reis, and Araújo, 1982

#### ASPREDINIDAE

*Agmus lyriformis* Eigenmann, 1912  
*Bunocephalus amaurus* Eigenmann, 1912  
*Bunocephalus chamaizelus* Eigenmann, 1912  
*Bunocephalus colombianus* Eigenmann, 1912  
*Bunocephalus depressus* Haseman, 1911  
*Ernstichthys intonsus* Stewart, 1985  
*Hoplomyzon papillatus* Stewart, 1985  
*Xiliphius magdalenae* Eigenmann, 1912

#### ASTROBLEPIDAE

*Astroblepus cyclopus santanderensis* Eigenmann, 1917

*Astroblepus frenatus* Eigenmann, 1917  
*Astroblepus grixalvii micrescens* Eigenmann, 1917  
*Astroblepus latidens* Eigenmann, 1917  
*Astroblepus pholeter* Collette, 1962  
*Astroblepus rosei* Eigenmann, 1922  
*Cyclopium chapmani* Eigenmann, 1912  
*Cyclopium pirrense* Meek and Hildebrand, 1913  
*Cyclopium trifasciatum* Eigenmann, 1912  
*Cyclopium unifasciatum* Eigenmann, 1912  
*Cyclopium vanceae* Eigenmann, 1913  
*Cyclopium ventrale* Eigenmann, 1912

#### ASTRONESTHIDAE

*Astronesthes ijimai* Tanaka, 1908

#### ATHERINIDAE

*Allanetta crenolepis* Schultz, 1953  
*Atherina evermanni* Eigenmann, 1903  
*Atherina insularum* Jordan and Evermann, 1903  
*Atherina ovalaua* Herre, 1935  
*Atherinichthys nouhuysi* Weber, 1910  
*Atherinops affinis littoralis* Hubbs, 1918  
*Atherinops insularum cedroscensis* Hubbs, 1918  
*Atherinops insularum guadalupae* Hubbs, 1918  
*Atherion elymus aphrozoicus* Schultz, 1953  
*Atherion elymus freyi* Schultz, 1953  
*Chirostoma attenuatum* Meek, 1902  
*Chirostoma consocium* Jordan and Hubbs, 1919  
*Chirostoma jordani* Woolman, 1894  
*Chirostoma labarcaei* Meek, 1902  
*Chirostoma mezquital* Meek, 1904  
*Chirostoma patzcuaro* Meek, 1902  
*Chirostoma regani* Jordan and Hubbs, 1919  
*Chirostoma zirahuen* Meek, 1902  
*Hypoatherina barnesi* Schultz, 1953  
*Ischnomembras gabunensis* Fowler, 1903  
*Melaniris balsanus* Meek, 1902  
*Melaniris sardina* Meek, 1907  
*Menidia chagresi* Meek and Hildebrand, 1914  
*Menidia lisa* Meek, 1904  
*Menidia peninsulae atrimentis* Kendall, 1902  
*Menidia starksi* Meek and Hildebrand, 1923  
*Mugilops cyanellus* Meek and Hildebrand, 1923  
*Poblana ferdebueni* Solórzano and López, 1965  
*Thyrina colombiensis* Hubbs, 1920  
*Thyrina guija* Hildebrand, 1925  
*Thyrina meeki* Miller, 1907

#### AUCHENIPTERIDAE

*Auchenipterus brevior* Eigenmann, 1912  
*Auchenipterus demerarae* Eigenmann, 1912  
*Entomocorus benjamini* Eigenmann, 1917

*Epapterus blohmi* Vari, Jewett, Taphorn, and Gilbert, 1984  
*Gelanoglanis stroudi* Böhlke, 1980  
*Glanidium ribeiroi* Haseman, 1911  
*Pseudauchenipterus nigrolineatus* Fowler, 1911  
*Pseudotatia parva* Mees, 1974  
*Trachelyichthys exilis* Greenfield and Glodek, 1977  
*Trachycorystes fisheri* Eigenmann, 1916

#### BAGRIDAE

*Bathylagus tetranema* Bailey and Stewart, 1984  
*Leiocassis robustus* Inger and Chin, 1959  
*Leiogaster melanopus* Weber, 1913  
*Liobagrus anguillicauda* Nichols, 1926  
*Liobagrus nantoenis* Oshima, 1919  
*Lophiobagrus aquilus* Bailey and Stewart, 1984  
*Lophiobagrus brevispinis* Bailey and Stewart, 1984  
*Mystus sabanus* Inger and Chin, 1959  
*Pseudobagrus adiposalis* Oshima, 1919  
*Pseudobagrus taiwanensis* Oshima, 1919

#### BALISTIDAE

*Monacanthus tuckeri* Bean, 1906

#### BATHYCLUPEIDAE

*Bathyclupea schroederi* Dick, 1962

#### BATHYLAGIDAE

*Bathylagus greyae* Cohen, 1958  
*Nansenia ardesiaca* Jordan and Thompson, 1914

#### BATRACHOIDIDAE

*Batrachoides waltersi* Collette and Russo, 1981  
*Porichthys bathoiketes* Gilbert, 1968  
*Porichthys kymosemeum* Gilbert, 1968  
*Porichthys myriaster* Hubbs and Schultz, 1939  
*Thalassophryne quadrizonatus* Eigenmann, 1922  
*Triathalassothia gloverensis* Greenfield and Greenfield, 1973

#### BELONIDAE

*Belone syetovidovi* Knudsen and Wolff, 1970  
*Rhaphiobelone robusta* Schultz, 1953

#### BELONTIIDAE

*Macropodus filamentosus* Oshima, 1919

#### BLENNIDAE

*Alticus novemmaculosus* Snyder, 1908  
*Blenniulus proteus* Krejsa, 1960  
*Blennius nicholsi* Tavalga, 1954

*Chasmodes bosquianus longimaxilla* Williams, 1983  
*Entomacrodus randalli* Springer, 1967  
*Hypsoblennius lignus* Meek and Hildebrand, 1928  
*Petroscirtes kulambangrae* Herre, 1935  
*Plagiotremus phenax* Smith-Vaniz, 1976  
*Salarias martini* Herre, 1942  
*Salarias walensis* Herre, 1935  
*Spinoblennius spiniger* Herre, 1935  
*Tagusa delicata* Herre, 1935

#### BOTHIDAE

*Ancylosetta antillarum* Gutherz, 1966  
*Ancylosetta cycloidea* Tyler, 1959  
*Ancylosetta kumperae* Tyler, 1959  
*Ancylosetta microctenus* Gutherz, 1966  
*Engyprosopon xenandrus* Gilbert, 1905  
*Monolene megalepis* Woods, 1961  
*Platophrys inermis* Gilbert, 1905  
*Trichopsetta caribbaea* Anderson and Gutherz, 1967

#### BRAMIDAE

*Eumegistus illustris* Jordan and Jordan, 1922

#### BRANCHIOSTOMATIDAE

*Branchiostoma bermudae* Hubbs, 1922  
*Branchiostoma floridae* Hubbs, 1922

#### CALLICHTHYIDAE

*Aspidoras maculosus* Nijssen and Isbrücker, 1976  
*Cascadura maculocephala* Ellis, 1913  
*Chaenothorax eigenmanni* Ellis, 1913  
*Corydoras australe* Eigenmann and Ward, 1907  
*Corydoras blochi* Nijssen, 1971  
*Corydoras bondi coppenamensis* Nijssen, 1970  
*Corydoras ellisae* Gosline, 1940  
*Corydoras heteromorphus* Nijssen, 1970  
*Corydoras schwartzi surinamensis* Nijssen, 1970  
*Corydoras weitzmani* Nijssen, 1971  
*Hoplosternum magdalenae* Eigenmann, 1913  
*Hoplosternum punctatum* Meek and Hildebrand, 1916

#### CALLIONYMIDAE

*Callionymus distethommatus* Fowler, 1941  
*Callionymus hudsoni* Fowler, 1941  
*Callionymus tenuis* Fricke, 1981  
*Callionymus wilburi* Herre, 1935  
*Calliurichthys astrinius* Jordan and Jordan, 1922  
*Calliurichthys zanectes* Jordan and Jordan, 1922

*Calymmichthys xenicus* Jordan and Thompson, 1914

*Synchiropus ijimae* Jordan and Thompson, 1914

*Synchiropus minutulus* Fricke, 1981

#### CAPROIDAE

*Antigonia combatia* Berry and Rathjen, 1958

*Antigonia malayana* Weber, 1913

#### CARANGIDAE

*Alectis breviventralis* Wakiya, 1924

*Alectis temmincki* Wakiya, 1924

*Caranx formosanus* Jordan and Snyder, 1909

*Caranx ishikawai* Wakiya, 1924

*Caranx miyakamii* Wakiya, 1924

*Caranx oshimai* Wakiya, 1924

*Caranx rastrosus* Jordan and Snyder, 1909

*Caranx tanakai* Wakiya, 1924

*Caranx uii* Wakiya, 1924

*Decapterus dayi* Wakiya, 1924

*Decapterus tabl* Berry, 1968

*Hemicaranx rhomboides* Meek and Hildebrand, 1925

*Scomberoides formosanus* Wakiya, 1924

*Trachinotus jordani* Wakiya, 1924

*Trachurus argenteus* Wakiya, 1924

*Ulva richardsoni* Jordan and Snyder, 1909

#### CARAPIDAE

*Disparichthys fluviatilis* Herre, 1935

*Echiodon dawsoni* Williams and Shipp, 1982

#### CARCHARHINIDAE

*Rhizoprionodon oligolinx* Springer, 1964

#### CATOSTOMIDAE

*Carpiodes elongatus* Meek, 1904

*Carpiodes labiosus* Meek, 1904

*Carpiodes microstomus* Meek, 1904

*Catostomus conchos* Meek, 1902

*Catostomus sonorensis* Meek, 1902

*Moxostoma rubriques* Hubbs, 1930

*Pantosteus jordani* Evermann, 1893

*Pantosteus santaanae* Snyder, 1908

#### CENTRARCHIDAE

*Lepomis occidentalis* Meek, 1902

*Micropterus pseudaplites* Hubbs, 1927

#### CENTRISCIDAE

*Centriscus capito* Oshima, 1922

#### CENTROPOMIDAE

*Ambassis lafa* Jordan and Seale, 1906

*Ambassis vaivasensis* Jordan and Seale, 1906

#### CETOPSIDAE

*Hemicetopsis amphioxus* Eigenmann, 1914

*Hemicetopsis macilentus* Eigenmann, 1912

*Hemicetopsis minutus* Eigenmann, 1912

*Hemicetopsis othonops* Eigenmann, 1912

#### CHAENOPSIDAE

*Acanthemblemaria greenfieldi* Smith-Vaniz and Palacio, 1974

*Chaenopsis alepidota californiensis* Böhlke, 1957

*Chaenopsis roseola* Hastings and Shipp, 1980

*Ekemblemaria myersi* Stephens, 1963

*Emblemaria angelica* Böhlke and Mead, 1957

*Emblemaria hyltoni* Johnson and Greenfield, 1976

*Emblemaria nigra* Meek and Hildebrand, 1928

*Emblemariopsis pricei* Greenfield, 1975

#### CHARACIDAE

*Acanthocharax microlepis* Eigenmann, 1912

*Acestrorhynchus altus* Menezes, 1969

*Acestrorhynchus guianensis* Menezes, 1969

*Acestrorhynchus nasutus* Eigenmann, 1912

*Agoniates anchovia* Eigenmann, 1914

*Alestes langi* Fowler, 1935

*Alestes thamalakanensis* Fowler, 1935

*Ammocryptocharax elegans* Weitzman and Kanazawa, 1976

*Aphyocharax anisitsi* Eigenmann and Kennedy, 1903

*Aphyocharax erythrurus* Eigenmann, 1912

*Aphyocharax melanotus* Eigenmann, 1912

*Aphyocharax paraguayensis* Eigenmann, 1915

*Aphyocheiroidon hemigrammus* Eigenmann, 1915

*Aphyodite grammica* Eigenmann, 1912

*Asiphonichthys hemigrammus* Eigenmann, 1912

*Astyanax abramoides* Eigenmann, 1909

*Astyanax aeneus costaricensis* Meek, 1914

*Astyanax albeolus* Eigenmann, 1908

*Astyanax aurocaudatus* Eigenmann, 1913

*Astyanax bimaculatus novae* Eigenmann, 1911

*Astyanax bimaculatus paraguayensis* Eigenmann, 1921

*Astyanax daguae* Eigenmann, 1913

*Astyanax essequibensis* Eigenmann, 1909

*Astyanax grandis* Meek and Hildebrand, 1912

*Astyanax guaporensis* Eigenmann, 1911

*Astyanax guianensis* Eigenmann, 1909

*Astyanax gymnogenys* Eigenmann, 1911

*Astyanax hasemani* Eigenmann, 1914

*Astyanax heterurus* Eigenmann and Wilson, 1914

*Astyanax kompi* Hildebrand, 1938

- Astyanax magdalenae* Eigenmann and Henn, 1916
- Astyanax marionae* Eigenmann, 1911
- Astyanax metae* Eigenmann, 1914
- Astyanax microlepis* Eigenmann, 1913
- Astyanax mucronatus* Eigenmann, 1909
- Astyanax mutator* Eigenmann, 1909
- Astyanax nasutus* Meek, 1907
- Astyanax notemigonoides* Fowler, 1911
- Astyanax paranahybae* Eigenmann, 1911
- Astyanax potaroensis* Eigenmann, 1909
- Astyanax regani* Meek, 1909
- Astyanax ribeirae* Eigenmann, 1911
- Astyanax robustus* Meek, 1912
- Astyanax ruberrimus* Eigenmann, 1913
- Astyanax scabripinnis paranae* Eigenmann, 1914
- Astyanax scierus* Fowler, 1911
- Bleptonema amazonae* Eigenmann, 1914
- Bleptonema paraguayensis* Eigenmann, 1914
- Brachyhalcinus guianensis* Boeseman, 1952
- Bramocharax elongatus* Meek, 1907
- Bryconamericus alfredae* Eigenmann, 1927
- Bryconamericus alpha* Eigenmann, 1914
- Bryconamericus beta* Eigenmann, 1914
- Bryconamericus cascajalensis* Meek and Hildebrand, 1916
- Bryconamericus caucanus* Eigenmann, 1913
- Bryconamericus chocoensis* Eigenmann, 1913
- Bryconamericus cismontanus* Eigenmann, 1914
- Bryconamericus conventus* Eigenmann, 1913
- Bryconamericus decurrens* Eigenmann, 1913
- Bryconamericus dentatus* Eigenmann, 1913
- Bryconamericus deuterodonoides* Eigenmann, 1914
- Bryconamericus deuterodonoides caudovittatus* Inger, 1956
- Bryconamericus diquensis* Eigenmann, 1913
- Bryconamericus grosvenori* Eigenmann, 1927
- Bryconamericus henni* Eigenmann, 1914
- Bryconamericus hyphessus* Eigenmann, 1909
- Bryconamericus magdalenensis* Eigenmann, 1913
- Bryconamericus novae* Eigenmann and Henn, 1914
- Bryconamericus ortholepis* Eigenmann, 1913
- Bryconamericus osgoodi* Eigenmann and Allen, 1942
- Bryconamericus pachacuti* Eigenmann, 1927
- Bryconamericus peruanus ricae* Eigenmann, 1908
- Bryconamericus scopiferus* Eigenmann, 1913
- Bryconamericus scopiferus guaytarae* Eigenmann and Henn, 1914
- Bryconamericus tolimae* Eigenmann, 1913
- Bryconamericus zeteki* Hildebrand, 1938
- Brycon argenteus* Meek and Hildebrand, 1913
- Brycon behreae* Hildebrand, 1938
- Brycon henni* Eigenmann, 1913
- Brycon obscurus* Hildebrand, 1938
- Brycon petrosus* Meek and Hildebrand, 1913
- Brycon siebenthalae* Eigenmann, 1912
- Ceratobranchia binghami* Eigenmann, 1927
- Characidium blennioides* Eigenmann, 1909
- Characidium catenatum* Eigenmann, 1909
- Characidium caucanus* Eigenmann, 1912
- Characidium laterale* Eigenmann, 1909
- Characidium pellucidum* Eigenmann, 1909
- Characidium phoxocephalum* Eigenmann, 1912
- Characidium pteroides* Eigenmann, 1909
- Characidium vintoni* Eigenmann, 1909
- Characidium zebra* Eigenmann, 1909
- Charax metae* Eigenmann, 1922
- Charax rupununi* Eigenmann, 1912
- Cheirodon annae* McAtee, 1903
- Cheirodon dialepturus* Fink and Weitzman, 1974
- Cheirodon eigenmanni* Meek, 1912
- Cheirodon galusdae* Eigenmann, 1927
- Cheirodon interruptus ibicuhiensis* Eigenmann, 1915
- Cheirodon madeirae* Eigenmann, 1915
- Cheirodon microdon* Eigenmann, 1915
- Cheirodon notomelas* Eigenmann, 1915
- Cheirodon parahybae* Eigenmann, 1915
- Cheirodon stenodon* Eigenmann, 1915
- Compsura heterura* Eigenmann, 1915
- Creagrutus beni* Eigenmann, 1911
- Creagrutus brevipinnis* Eigenmann, 1913
- Creagrutus caucanus* Eigenmann, 1913
- Creagrutus magdalenae* Eigenmann, 1913
- Creagrutus melanzonus* Eigenmann, 1909
- Creagrutus notropoides* Meek and Hildebrand, 1912
- Creagrutus simus* Meek and Hildebrand, 1913
- Cynopotamus essequebensis* Eigenmann, 1912
- Dermatocheir catablepta* Durbin, 1909
- Deuterodon acanthogaster* Eigenmann, 1911
- Deuterodon atracaudata* Meek and Hildebrand, 1912
- Deuterodon pinnatus* Eigenmann, 1909
- Deuterodon potaroensis* Eigenmann, 1909
- Fowlerina franciscensis* Eigenmann, 1914
- Genycharax tarpon* Eigenmann, 1912
- Gephyrocharax caucanus* Eigenmann, 1912
- Gephyrocharax chocoensis* Eigenmann, 1912
- Gephyrocharax intermedius* Meek and Hildebrand, 1916
- Gephyrocharax melanocheir* Eigenmann, 1912
- Gephyrocharax whaleri* Hildebrand, 1938

- Glandulocauda inequalis* Eigenmann, 1911  
*Glandulocauda melanogenys* Eigenmann, 1911  
*Glandulocauda melanopleura* Eigenmann, 1911  
*Hasemania bilineata* Ellis, 1911  
*Hasemania maxillaris* Ellis, 1911  
*Hasemania melanura* Ellis, 1911  
*Hemibrycon boquiae* Eigenmann, 1913  
*Hemibrycon colombianus* Eigenmann, 1914  
*Hemibrycon dariensis* Meek and Hildebrand, 1916  
*Hemibrycon helleri* Eigenmann, 1927  
*Hemigrammus analis* Durbin, 1909  
*Hemigrammus brevis* Ellis, 1911  
*Hemigrammus compressus* Meek, 1904  
*Hemigrammus cylindricus* Durbin, 1909  
*Hemigrammus erythrozonus* Durbin, 1909  
*Hemigrammus iota* Durbin, 1909  
*Hemigrammus kennedyi* Eigenmann, 1903  
*Hemigrammus marginatus* Ellis, 1911  
*Hemigrammus melanopterus* Eigenmann and Kennedy, 1903  
*Hemigrammus minutus* Meek and Hildebrand, 1912  
*Hemigrammus orthus* Durbin, 1909  
*Hemigrammus riddlei* Meek, 1907  
*Hemigrammus rodwayi* Durbin, 1909  
*Heterocharax macrolepis* Eigenmann, 1912  
*Holesthes heterodon* Eigenmann, 1915  
*Holopristis riddlei* Meek, 1907  
*Hyphessobrycon bifasciatus* Ellis, 1911  
*Hyphessobrycon duragenys* Ellis, 1911  
*Hyphessobrycon ecuadoriensis* Eigenmann and Henn, 1914  
*Hyphessobrycon eos* Durbin, 1909  
*Hyphessobrycon melanopleurus* Ellis, 1911  
*Hyphessobrycon minimus* Durbin, 1909  
*Hyphessobrycon minor* Durbin, 1909  
*Hyphessobrycon panamensis daguae* Eigenmann, 1922  
*Hyphessobrycon parvulus* Ellis, 1911  
*Hyphessobrycon poecilioides* Eigenmann, 1913  
*Hyphessobrycon proteus* Eigenmann, 1913  
*Hyphessobrycon reticulatus* Ellis, 1911  
*Hyphessobrycon rosaceus* Durbin, 1909  
*Hyphessobrycon stictus* Durbin, 1909  
*Hyphessobrycon taurocephalus* Ellis, 1911  
*Hysteronotus megalostomus* Eigenmann, 1911  
*Landonia latidens* Eigenmann and Henn, 1914  
*Macropsobrycon uruguayanae* Eigenmann, 1915  
*Megalamphodus megalopterus* Eigenmann, 1915  
*Megalamphodus micropterus* Eigenmann, 1915  
*Metynnis guaporensis* Eigenmann, 1915  
*Metynnis mola* Eigenmann and Kennedy, 1903  
*Metynnis roosevelti* Eigenmann, 1915  
*Microbrycon minutus* Eigenmann and Wilson, 1914  
*Microgenys minutus* Eigenmann, 1913  
*Microschemobrycon guaporensis* Eigenmann, 1915  
*Moenkhausia browni* Eigenmann, 1909  
*Moenkhausia lepidura hasemani* Eigenmann, 1917  
*Moenkhausia metae* Eigenmann, 1922  
*Moenkhausia phaeonota* Fink, 1979  
*Moenkhausia profunda* Eigenmann, 1912  
*Moenkhausia shideleri* Eigenmann, 1909  
*Mylossoma ocellatum* Eigenmann, 1915  
*Nematobrycon amphiloxus* Eigenmann and Wilson, 1914  
*Nematobrycon lacortei* Weitzman and Fink, 1971  
*Odontostilbe hastatus* Eigenmann, 1913  
*Odontostilbe melandetus* Eigenmann, 1912  
*Odontostilbe paraguayensis* Eigenmann and Kennedy, 1903  
*Odontostilbe trementinae* Eigenmann and Kennedy, 1903  
*Oligobrycon microstomus* Eigenmann, 1915  
*Parastremma sadina* Eigenmann, 1912  
*Parecbasis cyclolepis* Eigenmann, 1914  
*Petersius barnardi* Herre, 1936  
*Petersius conserialis* Hilgendorf, 1894  
*Phenacogaster beni* Eigenmann, 1911  
*Phenacogaster franciscoensis* Eigenmann, 1911  
*Phenacogaster megalostictus* Eigenmann, 1909  
*Phenacogaster microstictus* Eigenmann, 1909  
*Phenagoniates wilsoni* Eigenmann, 1914  
*Piabina analis* Eigenmann, 1914  
*Poecilocharax bovalii* Eigenmann, 1909  
*Poecilurichthys moenkhausii* Eigenmann and Kennedy, 1903  
*Pristella aubyni* Eigenmann, 1909  
*Probolodus heterostomus* Eigenmann, 1911  
*Psalidodon gymnodontus* Eigenmann, 1911  
*Pseudocheiroidon affinis* Meek and Hildebrand, 1916  
*Pseudocorynopoma heterandria* Eigenmann, 1914  
*Pterobrycon landoni* Eigenmann, 1913  
*Pygocentrus bilineatus* Eigenmann, 1909  
*Rhoadsia altipinna* Fowler, 1911  
*Roeboides cauae* Eigenmann, 1922  
*Roeboides hildebrandi* Eigenmann, 1922  
*Roeboides macrolepis* Meek and Hildebrand, 1913  
*Roeboides magdalenae* Eigenmann, 1922  
*Roeboides meeki* Eigenmann, 1922

*Roeboides occidentalis* Meek and Hildebrand, 1916  
*Roeboides salvadoris* Hildebrand, 1925  
*Roeboides thurni* Eigenmann, 1912  
*Serrasalmo hollandi* Eigenmann, 1915  
*Serrasalmo humeralis gracilior* Eigenmann, 1915  
*Serrasalmus coccogenis* Fowler, 1911  
*Spintherobolus broccae* Myers, 1925  
*Spintherobolus papilliferus* Eigenmann, 1911  
*Stethaprion crenatus* Eigenmann, 1916  
*Stevardia aliata* Eigenmann, 1914  
*Tetragonopterus alleni* Eigenmann and McAtee, 1907  
*Vesicatrux tegatus* Eigenmann, 1911

#### CHILODONTIDAE

*Tylobronchus maculosus* Eigenmann, 1912

#### CHIMAERIDAE

*Psychichthys eidolon* Jordan and Hubbs, 1925

#### CHLOROPHTHALMIDAE

*Bathypterois bigelowi* Mead, 1958  
*Bathypterois dubius* Vaillant, 1888  
*Bathypterois pectoralis* Garman, 1899  
*Bathytrophops marionae* Mead, 1958

#### CICHLIDAE

*Aequidens awani* Haseman, 1911  
*Aequidens azurifer* Fowler, 1911  
*Aequidens chimantanus* Inger, 1956  
*Aequidens duopunctata* Haseman, 1911  
*Aequidens guaporensis* Haseman, 1911  
*Aequidens mariae* Eigenmann, 1922  
*Aequidens plagiozonatus* Kullander, 1984  
*Aequidens potaroensis* Eigenmann, 1912  
*Apistogramma caetei* Kullander, 1980  
*Apistogramma inconspicua* Kullander, 1982  
*Apistogramma macmasteri* Kullander, 1979  
*Astronotus orbiculatus* Haseman, 1911  
*Chromis mossambicus* Peters, 1852  
*Chromis sparsidens* Hilgendorf, 1905  
*Cichlasoma acutum* Miller, 1907  
*Cichlasoma alfari* Meek, 1907  
*Cichlasoma boliviense* Kullander, 1983  
*Cichlasoma calobrense* Meek and Hildebrand, 1913  
*Cichlasoma centrale* Meek, 1906  
*Cichlasoma dorsatum* Meek, 1907  
*Cichlasoma eigenmanni* Meek, 1902  
*Cichlasoma evermanni* Meek, 1904  
*Cichlasoma frontale* Meek, 1909  
*Cichlasoma grammodes* Taylor and Miller, 1980  
*Cichlasoma granadense* Meek, 1907

*Cichlasoma hartwegi* Taylor and Miller, 1980  
*Cichlasoma hedricki* Meek, 1904  
*Cichlasoma meeki* Hildebrand, 1925  
*Cichlasoma milleri* Meek, 1907  
*Cichlasoma mojarra* Meek, 1904  
*Cichlasoma nigratum* Meek, 1907  
*Cichlasoma ornatum gephyrum* Eigenmann, 1922  
*Cichlasoma paranaense* Kullander, 1983  
*Cichlasoma punctatum* Meek, 1909  
*Cichlasoma spilatum* Meek, 1912  
*Cichlasoma tenue* Meek, 1906  
*Cichlasoma tuba* Meek, 1912  
*Cichlasoma turyense* Meek and Hildebrand, 1913  
*Cichlasoma umbriferum* Meek and Hildebrand, 1913  
*Cichlasoma urophthalmus mayorum* Hubbs, 1936  
*Cichlasoma zonatum* Meek, 1905  
*Crenicara altispinosa* Haseman, 1911  
*Crenicichla alta* Eigenmann, 1912  
*Crenicichla dorsocellata* Haseman, 1911  
*Crenicichla iguassuensis* Haseman, 1911  
*Crenicichla jaguarensis* Haseman, 1911  
*Crenicichla santaremensis* Haseman, 1911  
*Crenicichla simoni* Haseman, 1911  
*Geophagus brasiliensis iporangensis* Haseman, 1911  
*Geophagus brasiliensis itapicuruensis* Haseman, 1911  
*Gobiocichla ethelwynnae* Roberts, 1982  
*Gobiocichla wonderi* Kanazawa, 1951  
*Heros beani* Jordan, 1889  
*Heterogramma corumbae* Eigenmann and Ward, 1907  
*Heterogramma ortmanni* Eigenmann, 1912  
*Heterogramma ritense* Haseman, 1911  
*Heterogramma taeniatum pertense* Haseman, 1911  
*Heterogramma trifasciatum maciliense* Haseman, 1911  
*Lamprologus obliquus* Nichols and Griscom, 1917  
*Nannacara bimaculata* Eigenmann, 1912  
*Nanochromis transvestitus* Stewart and Roberts, 1984  
*Neetroplus carpintis* Jordan and Snyder, 1900  
*Neetroplus fluviatilis* Meek, 1912  
*Neetroplus panamensis* Meek and Hildebrand, 1913  
*Thorichthys ellioti* Meek, 1904

## CITHARINIDAE

*Distichodus stigmaturus* Fowler, 1935

## CLARIIDAE

*Clarias malaris* Nichols and Griscom, 1917

*Clarias mossambicus* Peters, 1852

## CLINIDAE

*Gibbonsia elegans erroli* Hubbs, 1952

*Gibbonsia elegans montereyensis* Hubbs, 1927

*Gibbonsia elegans velifera* Hubbs, 1952

*Gibbonsia metzi* Hubbs, 1927

*Gibbonsia norae* Hubbs, 1952

## CLUPEIDAE

*Alosa ohiensis* Evermann, 1902

*Dorosoma anale* Meek, 1904

*Dorosoma chavesi* Meek, 1907

*Gonostoma ebelingi* Grey, 1960

*Ilisha argentata* Meek and Hildebrand, 1923

*Jenkinsia majua majua* Whitehead, 1963

*Jenkinsia majua woodsi* Whitehead, 1963

*Odontognathus compressus* Meek and Hildebrand, 1923

*Opisthonema berlangai* Berry and Barrett, 1963

*Opisthonema medirastre* Berry and Barrett, 1963

*Pellonula tenuis* Nichols and Griscom, 1917

*Rhinosardinia serrata* Eigenmann, 1912

*Sardinella marquesensis* Berry and Whitehead, 1968

*Signalosa atchafalayae vanhyningi* Weed, 1925

*Signalosa mexicana campi* Weed, 1925

## COBITIDAE

*Acanthophtalmus mariae* Inger and Chin, 1962  
*Acanthophtalmus sandakanensis* Inger and Chin, 1962

*Barbatula toni posteroventralis* Nichols, 1925

*Barbatula yarkandensis sellaefer* Nichols, 1925

*Botia purpurea* Nichols, 1925

*Cobitis aurata vallahica* Nalbant, 1957

*Cobitis taenia melanoleuca* Nichols, 1925

*Crossostoma fascicauda* Nichols, 1926

*Misgurnus anguillicaudatus tungting* Nichols, 1925

*Misgurnus mizolepis fukien* Nichols, 1925

*Misgurnus mizolepis grangeri* Nichols, 1925

*Misgurnus moholyi yunnan* Nichols, 1925

*Oreias furcatus* Banareescu and Nalbant, 1976

## CONGRIDAE

*Conger jordani* Kanazawa, 1958

*Leptocephalus ectenurus* Jordan and Richardson, 1909

*Leptocephalus harringtonensis* Mowbray, 1931

## CORYPHAENIDAE

*Ectenias brunneus* Jordan and Thompson, 1914

## COTTIDAE

*Clinocottus analis australis* Hubbs, 1926

*Leptocottus armatus australis* Hubbs, 1921

## CURIMATIDAE

*Curimata cerasina* Vari, 1983

*Curimata incompta* Vari, 1983

*Curimatus atratoensis* Eigenmann, 1912

*Curimatus gillii* Eigenmann and Kennedy, 1903

*Curimatus issororoensis* Eigenmann, 1912

*Curimatus metae* Eigenmann, 1922

*Curimatus morawhannae* Eigenmann, 1912

*Curimatus patiae* Eigenmann, 1914

*Curimatus peruanus* Eigenmann, 1922

*Psectrogaster curviventris* Eigenmann and Kennedy, 1903

## CYCLOPTERIDAE

*Careproctus burkei* Jordan and Thompson, 1914

*Careproctus gilberti* Jordan and Thompson, 1914

## CYNOGLOSSIDAE

*Areliscus hollandi* Jordan and Metz, 1913

*Cynoglossus heterolepis* Weber, 1910

*Dollfusichthys sinusarabici* Chabanaud, 1931

## CYPRINIDAE

*Achahara jusanensis* Jordan and Hubbs, 1925

*Acheilognathus barbatus* Nichols, 1926

*Acheilognathus gracilis* Nichols, 1926

*Acheilognathus tabira* Jordan and Thompson, 1914

*Agosia adobe* Jordan and Evermann, 1891

*Agosia yarrowi* Jordan and Evermann, 1891

*Alburnops illecebrosus* Girard, 1857

*Algansea aphanea* Barbour and Miller, 1978

*Algansea monticola* Barbour and Contreras, 1968

*Algansea rubescens* Meek, 1902

*Amblypharyngodon grandisquamis* Jordan and Starks, 1917

*Aztecula mexicana* Meek, 1902

*Barbus akeleyi* Hubbs, 1918

*Barbus anoplus* Weber, 1897

*Barbus athi* Hubbs, 1918

*Barbus bantolanensis* Day, 1914

*Barbus bifrenatus* Fowler, 1935

*Barbus carens* Boulenger, 1912

*Barbus fitzsimonsi* Fowler, 1935

- Barbus helleri* Hubbs, 1918  
*Barbus oatesii* Boulenger, 1893  
*Barbus tsotsorogensis* Fowler, 1935  
*Beirabarbuis palustris* Herre, 1936  
*Belligobio eristigma* Jordan and Hubbs, 1925  
*Chondrostoma cyri orientalis* Bianco and Ban-arescu, 1982  
*Coripareius septentrionalis* Nichols, 1925  
*Culter aokii* Oshima, 1919  
*Dangila sabana* Inger and Chin, 1962  
*Dionda diaboli* Hubbs and Brown, 1956  
*Dionda erimyzonops* Hubbs and Miller, 1974  
*Evarra tlahuacensis* Meek, 1902  
*Formosania gilberti* Oshima, 1919  
*Gila minacae* Meek, 1902  
*Gnathopogon iijimae* Oshima, 1919  
*Gnathopogon ishikawae* Jordan and Thompson, 1914  
*Gnathopogon longifilis* Jordan and Hubbs, 1925  
*Gnathopogon majimae* Jordan and Hubbs, 1925  
*Gnathopogon suwae* Jordan and Hubbs, 1925  
*Gnathopogon tsuchigae* Jordan and Hubbs, 1925  
*Gobio coriparoides* Nichols, 1925  
*Gobio kesleri banaticus* Banarescu, 1953  
*Gobio rivuloides* Nichols, 1925  
*Gobio soldatovi minus* Nichols, 1925  
*Hampala macrolepidota sabana* Inger and Chin, 1962  
*Hemiculterella engraulis* Nichols, 1925  
*Hybognathus hankinsoni* Hubbs, 1929  
*Hybopsis x-punctata trautmani* Hubbs and Crowe, 1956  
*Hybopsis x-punctata x-punctata* Hubbs and Crowe, 1956  
*Labeo fisheri* Jordan and Starks, 1917  
*Labeo jordani* Oshima, 1919  
*Lepidomeda mollispinis mollispinis* Miller and Hubbs, 1960  
*Leuciscus carletoni* Kendall, 1903  
*Leucogobio polytaenia* Nichols, 1925  
*Moroco yamamotis* Jordan and Hubbs, 1925  
*Nematabramis alestes borneensis* Inger and Chin, 1962  
*Nototropis forlonensis* Meek, 1904  
*Nototropis nazas* Meek, 1904  
*Notropis aztecus* Woolman, 1894  
*Notropis chihuahua* Woolman, 1894  
*Notropis edwardraneyi* Suttkus and Clemmer, 1968  
*Notropis greenei* Hubbs and Ortenburger, 1929  
*Notropis muskoka* Meek, 1899  
*Notropis panarcys* Hubbs and Miller, 1978  
*Notropis robustus* Meek, 1902  
*Notropis saladonis* Hubbs and Hubbs, 1958  
*Notropis santarosaliae* Meek, 1902  
*Paracrossochilus aceris* Inger and Chin, 1962  
*Parapelecus eigenmanni* Jordan and Metz, 1913  
*Phoxinus cumberlandensis* Starnes and Starnes, 1978  
*Phoxinus orcuttii* Eigenmann and Eigenmann, 1890  
*Phoxiscus kikuchii* Oshima, 1919  
*Pimephales anuli* Kendall, 1903  
*Pseudaspius atrilatus* Jordan and Thompson, 1914  
*Pseudaspius bergi* Jordan and Metz, 1913  
*Pseudaspius modestus* Jordan and Metz, 1913  
*Pseudogobio chinssuensis* Nichols, 1926  
*Pseudoperilampus hondae* Jordan and Metz, 1913  
*Pseudorasbora altipinna* Nichols, 1925  
*Pseudorasbora depressirostris* Nichols, 1925  
*Puntius joaquinae* Wood, 1968  
*Puntius snyderi* Oshima, 1919  
*Puntius umali* Wood, 1968  
*Rhodeus chosenicus* Jordan and Metz, 1913  
*Rhodeus kurumeus* Jordan and Thompson, 1914  
*Sarcocheilichthys morii* Jordan and Hubbs, 1925  
*Scaphesthes tamusuiensis* Oshima, 1919  
*Spinibarbus hollandi* Oshima, 1919  
*Varicorhinus robustus* Nichols, 1925
- ## CYPRINODONTIDAE
- Cyprinodon bifasciatus* Miller, 1968  
*Cyprinodon dearborni* Meek, 1909  
*Cyprinodon fontinalis* Smith and Miller, 1980  
*Cyprinodon meeki* Miller, 1976  
*Cyprinodon orthonotus* Peters, 1844  
*Fundulus cubensis* Eigenmann, 1903  
*Fundulus neumanni* Hilgendorf, 1905  
*Fundulus oaxacae* Meek, 1902  
*Orestias mundus* Parenti, 1984  
*Profundulus candalarius* Hubbs, 1924  
*Zygonectes pulvereus* Evermann, 1892
- ## DACTYLOSCOPIDAE
- Gillellus greyae* Kanazawa, 1952
- ## DASYATIDAE
- Dasyatis ushieii* Jordan and Hubbs, 1925
- ## DICERATIIDAE
- Diceratias pileatus* Uwate, 1979
- ## DIRETMIDAE
- Diretmus pauciradiatus* Woods, 1973



## DORADIDAE

- Anduzedoras arleoi* Fernández-Yépez, 1968  
*Anduzedoras copei* Fernández-Yépez, 1968  
*Apuredoras rivasi* Fernández-Yépez, 1950  
*Doraops zuloagai* Schultz, 1944  
*Doras lentiginosus* Eigenmann, 1917  
*Hemidoras leporhinus* Eigenmann, 1912  
*Hemidoras micropoecus* Eigenmann, 1912  
*Hemidoras microstomus* Eigenmann, 1912  
*Hemidoras notospilus* Eigenmann, 1912  
*Hildadoras bolivarensis* Fernández-Yépez, 1968  
*Leptodoras linnelli* Eigenmann, 1912  
*Rhinodoras boehlkei* Glodek, Whitmire, and Orcés, 1976  
*Rhinodoras thomersoni* Taphorn and Lilystrom, 1984  
*Rhynchodoras woodsi* Glodek, 1976

## DRACONETTIDAE

- Draconetta oregona* Briggs and Berry, 1959

## ELEOTRIDAE

- Alepideleotris tigris* Herre, 1935  
*Boroda malua* Herre, 1935  
*Bostrychus zonatus* Weber, 1908  
*Dormitator gymnocephalus* Eigenmann, 1912  
*Eleotris herwerdenii* Weber, 1910  
*Eleotris heterodon* Weber, 1908  
*Eleotris isthmensis* Meek and Hildebrand, 1916  
*Eleotris latifasciatus* Meek and Hildebrand, 1912  
*Eleotris macrolepis* Meek, 1914  
*Encaeura evides* Jordan and Hubbs, 1925  
*Euleptoeleotris clarki* Hildebrand, 1938  
*Euleptoeleotris shropshirei* Hildebrand, 1938  
*Hemieleotris levis* Eigenmann, 1917  
*Leptophilypnus fluviatilis* Meek and Hildebrand, 1916  
*Microeleotris mindii* Meek and Hildebrand, 1916  
*Microeleotris panamensis* Meek and Hildebrand, 1916  
*Pogoneleotris microps* Weber, 1908

## EMBIOTOCIDAE

- Crossochir koelzi* Hubbs, 1933

## ENGRAULIDAE

- Anchoa hepsetus colonensis* Hildebrand, 1943  
*Anchoa lamprotaenia* Hildebrand, 1943  
*Anchoa mitchilli diaphana* Hildebrand, 1943  
*Anchovia arenicola* Meek and Hildebrand, 1923  
*Anchovia brevirostra* Meek and Hildebrand, 1923  
*Anchovia eigenmannia* Meek and Hildebrand, 1923  
*Anchovia elongata* Meek and Hildebrand, 1923

- Anchovia parva* Meek and Hildebrand, 1923  
*Anchoviella belizensis* Thomerson and Greenfield, 1975  
*Anchoviella helleri* Hubbs, 1921  
*Stolephorus branchiomelas* Eigenmann, 1917  
*Stolephorus guianensis* Eigenmann, 1912  
*Stolephorus lyolepis* Evermann and Marsh, 1902

## EVERMANNELLIDAE

- Evermannella ahlstromi* Johnson and Glodek, 1975  
*Evermannella sicaria* Rofen, 1963

## EXOCOETIDAE

- Cypselurus alienus* Herre, 1935  
*Cypselurus antoncichi* Woods and Schultz, 1953

## GASTEROPELECIDAE

- Thoracocharax brevis* Eigenmann, 1912  
*Thoracocharax magdalenae* Eigenmann, 1912

## GEMPYLIDAE

- Diplogonurus maderensis* Noronha, 1926  
*Epinnula orientalis americana* Grey, 1953  
*Epinnula orientalis pacifica* Grey, 1953

## GERREIDAE

- Diapterus evermanni* Meek and Hildebrand, 1925  
*Eucিনostomus californiensis elongatus* Meek and Hildebrand, 1925

## GOBIESOCIDAE

- Cotylin nigripinnis woodsi* Schultz, 1944  
*Gobiesox costaricensis* Meek, 1912  
*Gobiesox fluviatilis* Briggs and Miller, 1960  
*Gobiesox fulvus* Meek, 1907  
*Gobiesox marijeanae* Briggs, 1960  
*Gobiesox mexicanus* Briggs and Miller, 1960  
*Gobiesox paradiseus* Herre, 1935  
*Gobiesox potamius* Briggs, 1955

## GOBIIDAE

- Amblygobius myersi* Herre, 1935  
*Aparrius aurocingulus* Herre, 1935  
*Atuona tricuspidata* Herre, 1935  
*Awaous decemlineatus* Eigenmann, 1917  
*Barbuligobius boehlkei* Lachner and McKinney, 1974  
*Brachygobius kabiliensis* Inger, 1958  
*Brachygobius sabanus* Inger, 1958  
*Brachygobius xanthomelas* Herre, 1937  
*Callogobius ocellatus* Herre, 1935  
*Chasmias misakius* Jordan and Snyder, 1901

*Coryphopterus punctipectophorus* Springer, 1960  
*Cryptocentrus geniornatus* Herre, 1935  
*Ctenogobius aterrimus* Herre, 1935  
*Ctenogobius leavelli* Herre, 1935  
*Ctenogobius malekulae* Herre, 1935  
*Ctenogobius myxodermus* Herre, 1935  
*Ctenogobius opalescens* Herre, 1936  
*Ctenogobius villadolidi* Herre, 1936  
*Ctenogobius virgatulus* Jordan and Snyder, 1901  
*Ctenogobius waigiensis* Herre, 1935  
*Euctenogobius liolepis* Meek and Hildebrand, 1928  
*Euctenogobius panamensis* Meek and Hildebrand, 1928  
*Eviota japonica* Jewett and Lachner, 1983  
*Evorthodus minutus* Meek and Hildebrand, 1928  
*Gerhardinus nudus* Meek and Hildebrand, 1928  
*Gladiogobius ensifer* Herre, 1933  
*Glossogobius abacopus* Jordan and Richardson, 1909  
*Glossogobius koragensis* Herre, 1935  
*Glossogobius parvus* Oshima, 1919  
*Glossogobius sandakanensis* Inger, 1957  
*Gnatholepis corlettei* Herre, 1935  
*Gnatholepis knighti* Jordan and Evermann, 1903  
*Gobiella birtwistlei* Herre, 1934  
*Gobionellus saepepallens* Gilbert and Randall, 1968  
*Gobiopsis aporia* Lachner and McKinney, 1978  
*Gobiopsis woodsi* Lachner and McKinney, 1978  
*Gobius claytonii* Meek, 1902  
*Gobius cliffordpopei* Nichols, 1925  
*Gobius daguae* Eigenmann, 1917  
*Gobius parvus* Meek, 1902  
*Macgregorella badia* Herre, 1935  
*Macgregorella santa* Herre, 1935  
*Macrodonogobius wilburi* Herre, 1936  
*Quisquilius eugenius* Jordan and Evermann, 1903  
*Rhinogobius formosanus* Oshima, 1919  
*Rhinogobius mowbrayi* Bean, 1906  
*Rhinogobius taiwanus* Oshima, 1919  
*Sagamia russula* Jordan and Snyder, 1901  
*Sicydium altum* Meek, 1907  
*Sicydium hildebrandi* Eigenmann, 1917  
*Sicydium montanum* Hubbs, 1920  
*Suruga fundicola* Jordan and Snyder, 1901  
*Tridentiger kuroiwae* Jordan and Tanaka, 1927  
*Vaimosa balteata* Herre, 1935  
*Vaimosa horiae* Herre, 1936  
*Vaimosa osgoodi* Herre, 1935  
*Varicus bucca* Robins and Böhlke, 1961  
*Varicus imswae* Greenfield, 1981

## GOBIOIDIDAE

*Taenioides snyderi* Jordan and Hubbs, 1925

## GOODEIDAE

*Characodon multiradiatus* Meek, 1904  
*Goodea captiva* Hubbs, 1924  
*Goodea toweri* Meek, 1904  
*Goodea withei* Meek, 1904  
*Skiffia lermiae* Meek, 1902  
*Skiffia variegata* Meek, 1902  
*Zoogoneticus diazi* Meek, 1902  
*Zoogoneticus miniatus* Meek, 1902

## GRAMMISTIDAE

*Rhegma bermudensis* Kanazawa, 1952

## HAEMULIDAE

*Anisotremus williamsi* Meek, 1905  
*Pomadasys grandis* Meek, 1907  
*Pomadasys starri* Meek, 1904  
*Pomadasys templei* Meek, 1904

## HELOGENIDAE

*Helogenes unidorsalis* Glodek and Carter, 1978

## HEMIODONTIDAE

*Bivibranchia protractila* Eigenmann, 1912  
*Hemiodus orthonops* Eigenmann and Kennedy, 1903  
*Hemiodus parnaguae* Eigenmann and Henn, 1916  
*Hemiodus quadrimaculatus* Pellegrin, 1908

## HEMIRAMPHIDAE

*Hemiramphus archipelagicus* Collette, 1978  
*Hemiramphus bermudensis* Collette, 1962  
*Hyporhamphus hildebrandi* Jordan and Evermann, 1927  
*Hyporhamphus melanopterus* Collette and Parin, 1978  
*Hyporhamphus snyderi* Meek and Hildebrand, 1923  
*Odontorhamphus cancellori* Weed, 1933  
*Zenarchopterus robertsi* Collette, 1982  
*Zenarchopterus sepikensis* Herre, 1935

## HETERENCHELYIDAE

*Heterenchelys biaggii* Böhlke, 1956

## HEXAGRAMMIDAE

*Pleurogrammus azonus* Jordan and Metz, 1913  
*Stellistius katsukii* Jordan and Tanaka, 1927

## HOLOCENTRIDAE

- Adioryx poco* Woods, 1965  
*Flammeo scythrops* Jordan and Evermann, 1903  
*Holocentrus bullisi* Woods, 1955  
*Holocentrus ensifer* Jordan and Evermann, 1903  
*Holocentrus meeki* Bean, 1906  
*Holocentrus xantherythrus* Jordan and Evermann, 1903  
*Myripristis argyromus* Jordan and Evermann, 1903  
*Myripristis berndti* Jordan and Evermann, 1903  
*Myripristis bowditchae* Woods, 1953  
*Myripristis chryseres* Jordan and Evermann, 1903  
*Myripristis mooreanus* Herre, 1935  
*Myripristis randalli* Greenfield, 1974  
*Myripristis symmetricus* Jordan and Evermann, 1903  
*Myripristis tiki* Greenfield, 1974  
*Myripristis undecimalis* Herre, 1935  
*Myripristis woodsi* Greenfield, 1974  
*Ostichthys acanthorhinus* Randall, Shimizu, and Yamakawa, 1982  
*Sargocentron macrosquamis* Golani, 1984

## HOMALOPTERIDAE

- Gastromyzon fasciatus* Inger and Chin, 1961  
*Gastromyzon lepidogaster* Roberts, 1982  
*Gastromyzon megalepis* Roberts, 1982  
*Gastromyzon pauciradiatus* Inger and Chin, 1961  
*Gastromyzon punctulatus* Inger and Chin, 1961  
*Gastromyzon ridens* Roberts, 1982  
*Homaloptera hingi* Herre, 1934  
*Protomyzon aphelocheilus* Inger and Chin, 1962

## ICTALURIDAE

- Ameiurus australis* Meek, 1904  
*Ameiurus mexicanus* Meek, 1904  
*Noturus albater* Taylor, 1969  
*Noturus flavater* Taylor, 1969  
*Noturus munitus* Suttkus and Taylor, 1965

## JENYNSHIDAE

- Fitzroyia eigenmanni* Haseman, 1911

## KNERIIDAE

- Grasseichthys gabonensis* Géry, 1964

## KUHLIIDAE

- Kuhlia bilunulata* Herre, 1935

## LABRIDAE

- Halichoeres fijiensis* Herre, 1935  
*Hemipteronotus baldwini* Jordan and Evermann, 1903

- Hemipteronotus evides* Jordan and Richardson, 1909  
*Iridio bimaculata* Wilson, 1916  
*Iridio decoratus* Bean, 1906  
*Iridio elegans* Bean, 1906  
*Iridio meyeri* Bean, 1906  
*Iridio microstomus* Bean, 1906  
*Lepidaplois strophodes* Jordan and Evermann, 1903  
*Thalassoma cranei* Herre, 1935  
*Xyrichthys niveilatus* Jordan and Evermann, 1903  
*Xyrichthys sciistius* Jordan and Thompson, 1914

## LABRISOMIDAE

- Exerpes asper earli* Hubbs, 1952  
*Labrisomus lentiginosus* Bean, 1906  
*Labrisomus multiporosus* Hubbs, 1953  
*Malacotenus afueræ multipunctatus* Springer, 1958  
*Malacotenus ebisui* Springer, 1958  
*Malacotenus gigas* Springer, 1958  
*Malacotenus hubbsi hubbsi* Springer, 1958  
*Malacotenus hubbsi polyporosus* Springer, 1958  
*Malacotenus margaritæ mexicanus* Springer, 1958  
*Malacotenus puertoricensis* Evermann and Marsh, 1900  
*Malacotenus triangulatus* Springer, 1958  
*Malacotenus zacae* Springer, 1958  
*Malacotenus zonifer sudensis* Springer, 1958  
*Mnierpes macrocephalus catherinae* Hubbs, 1952  
*Nemaclinus atelestos* Böhlke and Springer, 1975  
*Paraclinus integripinnis zacae* Hubbs, 1952  
*Starksia occidentalis* Greenfield, 1979  
*Starksia variabilis* Greenfield, 1979  
*Starksia y-lineata* Gilbert, 1965

## LEBIASINIDAE

- Archicheir minutus* Eigenmann, 1909  
*Copeina osgoodi* Eigenmann, 1922  
*Nannostomus marginatus* Eigenmann, 1909  
*Nannostomus minimus* Eigenmann, 1909  
*Nannostomus simplex* Eigenmann, 1909  
*Poecilobrycon auratus* Eigenmann, 1909  
*Poecilobrycon erythrus* Eigenmann, 1909  
*Poecilobrycon harrisoni* Eigenmann, 1909  
*Poecilobrycon ocellatus* Eigenmann, 1909  
*Pyrhulina australe* Eigenmann and Kennedy, 1903

## LEIOGNATHIDAE

- Gazza achlamys* Jordan and Starks, 1917

## LETHRINIDAE

*Gnathodentex oculomaculatus* Herre, 1935

## LOPHIIDAE

*Lophiodes heroe* Caruso, 1981

*Lophiodes reticulatus* Caruso and Suttkus, 1979

## LORICARIIDAE

*Acanthicus canensis* Meek and Hildebrand, 1913

*Acestridium discus* Haseman, 1911

*Ancistrus lithurgicus* Eigenmann, 1912

*Ancistrus melas* Eigenmann, 1916

*Ancistrus planiceps* Meek and Hildebrand, 1913

*Ancistrus spinosus* Meek and Hildebrand, 1916

*Chaetostomus leucomelas* Eigenmann, 1917

*Corymbophanes andersoni* Eigenmann, 1909

*Farlowella azygia* Eigenmann and Vance, 1917

*Farlowella hasemani* Eigenmann and Vance, 1917

*Farlowella jauruensis* Eigenmann and Vance, 1917

*Hemiancistrus daguae* Eigenmann, 1912

*Hemiancistrus mayoloi* Eigenmann, 1912

*Hemiancistrus wilsoni* Eigenmann, 1917

*Lasiancistrus brevispinis* Heitmans, Nijssen, and Isbrücker, 1983

*Lasiancistrus caucanus* Eigenmann, 1912

*Lithogenes villosus* Eigenmann, 1909

*Lithoxus lithoides* Eigenmann, 1912

*Loricaria capetensis* Meek and Hildebrand, 1913

*Loricaria filamentosa latiura* Eigenmann and Vance, 1912

*Loricaria filamentosa seminuda* Eigenmann and Vance, 1912

*Loricaria fimbriata* Eigenmann and Vance, 1912

*Loricaria griseus* Eigenmann, 1909

*Loricaria gymnogaster* Eigenmann and Vance, 1912

*Loricaria gymnogaster lagoichthys* Schultz, 1944

*Loricaria microdon* Eigenmann, 1909

*Loricaria stewarti* Eigenmann, 1909

*Loricaria submarginatus* Eigenmann, 1909

*Loricaria tuyrensis* Meek and Hildebrand, 1913

*Loricaria uracantha rupestre* Schultz, 1944

*Loricaria variegata venezuelae* Schultz, 1944

*Oxyloricaria citurensis* Meek and Hildebrand, 1913

*Oxyloricaria dariensis* Meek and Hildebrand, 1913

*Parotocinclus jimi* Garavello, 1976

*Plecostomus derbyi* Haseman, 1911

*Plecostomus hemiurus* Eigenmann, 1912

*Pseudancistrus carnegiei* Eigenmann, 1916

*Pseudancistrus nigrescens* Eigenmann, 1912

*Pseudancistrus pediculatus* Eigenmann, 1917

*Pterygoplichthys anisitsi* Eigenmann and Kennedy, 1903

*Rineloricaria formosa* Isbrücker and Nijssen, 1979

## LUTJANIDAE

*Etelis evurus* Jordan and Evermann, 1903

*Pristipomoides freemani* Anderson, 1966

## MACROURIDAE

*Coelorhynchus gilberti* Jordan and Hubbs, 1925

*Coryphaenoides bonanox* Jordan and Thompson, 1914

*Hymenocephalus aterrimus* Gilbert, 1905

*Hymenocephalus grimaldii* Weber, 1913

*Macrurus richardi* Weber, 1913

*Optonurus atherodon* Gilbert and Cramer, 1896

*Squalogadus intermedius* Grey, 1959

## MASTACEMBELIDAE

*Mastacembelus keithi* Herre, 1940

*Mastacembelus oatesii* Boulenger, 1893

## MELAMPHAIDAE

*Melamphaes pumilus* Ebeling, 1962

## MELANOTAENIIDAE

*Melanotaenia kabia* Herre, 1935

*Melanotaenia rosacea* Herre, 1935

*Rhombatractus affinis* Weber, 1908

*Rhombatractus sentaniensis* Weber, 1908

*Rhombosoma sepikensis* Herre, 1935

## MICRODESMIDAE

*Gunnellichthys viridesens* Dawson, 1968

*Microdesmus carri* Gilbert, 1966

*Microdesmus woodsi* Kanazawa, 1952

## MORIDAE

*Physiculus grinnelli* Jordan and Jordan, 1922

## MUGILIDAE

*Joturus daguae* Eigenmann, 1917

*Liza formosae* Oshima, 1922

*Liza parva* Oshima, 1922

*Liza pescadorensis* Oshima, 1922

*Mugil anpinensis* Oshima, 1922

*Myxus malayanus* Herre, 1936

## MULLIDAE

*Mulloides flammeus* Jordan and Evermann, 1903

*Pseudupeneus chrysonemus* Jordan and Evermann, 1903

*Upeneus arge* Jordan and Evermann, 1903

## MURAENIDAE

*Anarchias yoshiae* Kanazawa, 1952

*Echidna unicolor* Schultz, 1953

*Echidna vineta* Jenkins, 1903

*Echidna zonophaea* Jordan and Evermann, 1903

*Enchelycore carychroa* Böhlke and Böhlke, 1976

*Gymnothorax bayeri* Schultz, 1953

*Gymnothorax bikiniensis* Schultz, 1953

*Gymnothorax efatensis* Herre, 1935

*Gymnothorax leucostigma* Jordan and Richardson, 1909

*Rabula fuscomaculata* Schultz, 1951

*Rabula marshallensis* Schultz, 1953

*Uropterygius fuscoguttatus* Schultz, 1953

## MYCTOPHIDAE

*Dasyscopelus orientalis* Gilbert, 1913

*Diaphus gigas* Gilbert, 1913

*Diaphus glandulifer* Gilbert, 1913

*Diaphus latus* Gilbert, 1913

*Diaphus sagamiensis* Gilbert, 1913

*Lampanyctus jordani* Gilbert, 1913

*Lampanyctus micropunctatus* Chapman, 1939

*Macrostoma quercinum japonicum* Tanaka, 1908

*Myctophum hollandi* Jordan and Jordan, 1922

## MYXINIDAE

*Myxine biniplicata* Richardson and Jowett, 1951

## NEMIPTERIDAE

*Parascolopsis townsendi* Boulenger, 1901

*Scolopsis eriomma* Jordan and Richardson, 1909

## NEOSCOPELIDAE

*Scopelogys clarkei* Butler and Ahlstrom, 1976

*Solivomer arenidens* Miller, 1947

## NEOSTETHIDAE

*Neostethus borneensis* Herre, 1939

## NOMEIDAE

*Icticus ischanus* Jordan and Thompson, 1914

## NOTACANTHIDAE

*Polyacanthonotus merretti* Sulak, Crabtree, and Hureau, 1984

## OGCOCEPHALIDAE

*Ogcocephalus corniger* Bradbury, 1980

*Ogcocephalus declivirostris* Bradbury, 1980

*Ogcocephalus pantostictus* Bradbury, 1980

*Ogcocephalus pumilus* Bradbury, 1980

*Ogcocephalus rostellum* Bradbury, 1980

## OPHICHTHIDAE

*Aplatophis chauliodus* Böhlke, 1956

*Callechelys bilinearis* Kanazawa, 1952

*Cirriacaecula johnsoni* Schultz, 1953

*Ophichthus evermanni* Jordan and Richardson, 1909

*Ophichthus hyposagmatus* McCosker and Böhlke, 1984

*Sphagebranchus conklini* Eigenmann, 1916

## OPHIDIIDAE

*Calamopteryx jeb* Cohen, 1973

*Dicrolene kanazawai* Grey, 1958

*Grammonus mowbrayi* Grey, 1951

*Lepophidium aporrhox* Robins, 1961

*Lepophidium pheromystax* Robins, 1960

*Lepophidium staurophor* Robins, 1958

*Luciobrotula corethromycter* Cohen, 1954

*Monomitopus kumae* Jordan and Hubbs, 1925

*Othophidium schmidtii* Woods and Kanazawa, 1951

*Porogadus silus* Carter and Sulak, 1984

*Spectrunculus radcliffei* Jordan and Thompson, 1914

## OPISTOGNATHIDAE

*Lonchopisthus higmani* Mead, 1959

## OSMERIDAE

*Spirinchus dilatus* Schultz and Chapman, 1934

*Spirinchus verecundus* Jordan and Metz, 1913

## PANGASIIDAE

*Pangasius tubbi* Inger and Chin, 1959

## PARALEPIDAE

*Paralepis philippinus* Fowler, 1934

## PARODONTIDAE

*Apareiodon hasemani* Eigenmann, 1916

*Apareiodon itapicuruensis* Eigenmann and Henn, 1916

*Parodon bifasciatus* Eigenmann, 1912

*Parodon dariensis* Meek and Hildebrand, 1913

*Parodon ecuadoriensis* Eigenmann and Henn, 1914

*Parodon terminalis* Eigenmann and Henn, 1914

## PEMPHERIDAE

*Liopempheris sasakii* Jordan and Hubbs, 1925

*Pempheris dispar* Herre, 1935

## PERCICHTHYIDAE

*Malakichthys wakiyai* Jordan and Hubbs, 1925

## PERCIDAE

*Etheostoma obeyense* Kirsch, 1892

*Hadropterus scierus apristis* Hubbs and Hubbs, 1954

*Hololepis saludae* Hubbs and Cannon, 1935

*Stizostedion glaucum* Hubbs, 1926

*Zingel streber nerensis* Banarescu and Nalbant, 1979

## PERCOPHIDAE

*Bembrops greyi* Poll, 1959

## PETROMYZONTIDAE

*Ichthyomyzon unicuspis* Hubbs and Trautman, 1937

## PHOTICHTHYIDAE

*Vinciguerria mabahiss* Johnson and Feltes, 1984

## PIMELODIDAE

*Brachyglanis frenata* Eigenmann, 1912

*Brachyglanis melas* Eigenmann, 1912

*Brachyglanis phalacra* Eigenmann, 1912

*Cephalosilurus fowleri* Haseman, 1911

*Cetopsorhamdia boquillae* Eigenmann, 1922

*Cetopsorhamdia nasus* Eigenmann and Fisher, 1916

*Cetopsorhamdia phantasia* Stewart, 1985

*Chasmocranus brevior* Eigenmann, 1912

*Chasmocranus chimantanus* Inger, 1956

*Chasmocranus longior* Eigenmann, 1912

*Chasmocranus rosae* Eigenmann, 1922

*Cheirocerus eques* Eigenmann, 1917

*Heptapterus stewarti* Haseman, 1911

*Imparfinis hollandi* Haseman, 1911

*Imparfinis microps* Eigenmann and Fisher, 1916

*Imparfinis mirini* Haseman, 1911

*Leptoglanis essequibensis* Eigenmann, 1912

*Megalonema platycephalum* Eigenmann, 1912

*Megalonema punctatum* Meek and Hildebrand, 1913

*Megalonema robustum* Meek and Hildebrand, 1913

*Megalonema xanthum* Eigenmann, 1912

*Microglanis iheringi* Gomes, 1946

*Microglanis poecilus* Eigenmann, 1912

*Microglanis secundus* Mees, 1974

*Microglanis variegatus* Eigenmann, 1914

*Myoglanis potaroensis* Eigenmann, 1912

*Nannorhamdia nemacheir* Eigenmann and Fisher, 1916

*Pimelodella avanhandavae* Eigenmann, 1917

*Pimelodella boliviana* Eigenmann, 1917

*Pimelodella eigenmanni* Meek, 1905

*Pimelodella griffini* Eigenmann, 1917

*Pimelodella hasemani* Eigenmann, 1917

*Pimelodella itapicuruensis* Eigenmann, 1917

*Pimelodella laticeps* Eigenmann, 1917

*Pimelodella laticeps australis* Eigenmann, 1917

*Pimelodella macturki* Eigenmann, 1912

*Pimelodella megalops* Eigenmann, 1912

*Pimelodella metae* Eigenmann, 1917

*Pimelodella notomelas* Eigenmann, 1917

*Pimelodella serrata* Eigenmann, 1917

*Pimelodus clarias coprophagus* Schultz, 1944

*Pimelodus heteropleurus* Eigenmann, 1912

*Pimelodus ortmanni* Eigenmann, 1911

*Platysilurus barbatus* Haseman, 1911

*Pseudopimelodus albomarginatus* Eigenmann, 1912

*Pseudopimelodus villosus* Eigenmann, 1912

*Rhamdella montana* Eigenmann, 1913

*Rhamdia barbata* Meek, 1907

*Rhamdia branneri* Haseman, 1911

*Rhamdia branneri voulezi* Haseman, 1911

*Rhamdia cabreræ* Meek, 1906

*Rhamdia laticauda typhla* Greenfield, Greenfield, and Woods, 1982

*Rhamdia luigiana* Villa, 1977

*Rhamdia microps* Eigenmann, 1917

*Rhamdia nasuta* Meek, 1909

*Rhamdia oaxacæ* Meek, 1902

*Rhamdia regani* Meek, 1907

*Rhamdiopsis moreirai* Haseman, 1911

*Sovichthys abuelo* Schultz, 1944

## PLESIOPIDAE

*Plesiops cephalotaenia* Inger, 1955

## PLOTOSIDAE

*Copidoglanis novaeguineae* Weber, 1908

*Plotosus papuensis* Weber, 1910

## POECILIIDAE

*Acanthophaelus bifurcus* Eigenmann, 1909

*Acanthophaelus melanzonus* Eigenmann, 1909

*Alfaro acutiventralis* Meek, 1912

*Cnesterodon carnegiei* Haseman, 1911

*Diphyacantha chochoensis* Henn, 1916

*Gambusia alvarezii* Hubbs and Springer, 1957

*Gambusia bonita* Meek, 1904

*Gambusia caliensis* Eigenmann and Henn, 1916 in Henn, 1916

*Gambusia cana* Meek and Hildebrand, 1913

*Gambusia cascajalensis* Meek and Hildebrand, 1913  
*Gambusia darienensis* Meek and Hildebrand, 1913  
*Gambusia fasciata* Meek, 1904  
*Gambusia heterochir* Hubbs, 1957  
*Gambusia hurtadoi* Hubbs and Springer, 1957  
*Gambusia infans* Woolman, 1894  
*Gambusia latipunctata* Meek and Hildebrand, 1913  
*Gambusia nigroventralis* Eigenmann and Henn, 1912  
*Gambusia panuco* Hubbs, 1926  
*Gambusia parismina* Meek, 1912  
*Gambusia regani* Hubbs, 1926  
*Gambusia turrubarensis* Meek, 1912  
*Gambusia umbratilis* Meek, 1912  
*Gambusia vittata* Hubbs, 1926  
*Gambusia xanthosoma* Greenfield, 1983  
*Gambusia yucatanica australis* Greenfield, 1985  
*Girardinus garmani* Eigenmann, 1903  
*Glaridichthys falcatus* Eigenmann, 1903  
*Heterandria colombianus* Eigenmann and Henn, 1912  
*Heterandria cubensis* Eigenmann, 1903  
*Heterandria hasemani* Henn, 1916  
*Heterandria lutzi* Meek, 1902  
*Limia hollandi* Henn, 1916  
*Limia rivasi* Franz and Burgess, 1983  
*Neoheterandria elegans* Henn, 1916  
*Phalloptychus eigenmanni* Henn, 1916  
*Phallotorynus fasciolatus* Henn, 1916  
*Platypoecilus tropicus* Meek, 1907  
*Platypoecilus variatus* Meek, 1904  
*Poecilia amates* Miller, 1907  
*Poecilia caudata* Meek, 1909  
*Poecilia fria* Eigenmann and Henn, 1914  
*Poecilia latipunctata* Meek, 1904  
*Poecilia pittieri* Meek, 1912  
*Poecilia tenuis* Meek, 1907  
*Poeciliopsis balsas* Hubbs, 1926  
*Priapichthys annectens hesperis* Hubbs, 1924  
*Priapichthys fosteri* Hildebrand, 1925  
*Priapichthys letonai* Hildebrand, 1925  
*Priapichthys olomina* Meek, 1914  
*Priapichthys panamensis* Meek and Hildebrand, 1916  
*Tomeurus gracilis* Eigenmann, 1909  
*Toxus riddlei* Eigenmann, 1903  
*Xiphophorus jalapae* Meek, 1902

#### POMACANTHIDAE

*Centropyge argi* Woods and Kanazawa, 1951  
*Centropyge heraldi* Woods and Schultz, 1953

*Centropyge tutuilae* Jordan and Jordan, 1922  
*Holacanthus lunatus* Blosser, 1909

#### POMACENTRIDAE

*Abudefduf lorenzi* Hensley and Allen, 1977  
*Chromis alta* Greenfield and Woods, 1980  
*Chromis atripectoralis* Welander and Schultz, 1951  
*Chromis limbaughi* Greenfield and Woods, 1980  
*Chromis meridiana* Greenfield and Woods, 1980  
*Chromis randalli* Greenfield and Hensley, 1970  
*Chromis villadolidi* Jordan and Tanaka, 1927  
*Chromis weberi* Fowler and Bean, 1928  
*Chromis woodsi* Bruner and Arnam, 1979  
*Eupomacentrus chrysus* Bean, 1906  
*Eupomacentrus rocasensis* Emery, 1972  
*Glyphisodon troscheli* Gill, 1863  
*Microspathodon frontatus* Emery, 1970  
*Pomacentrus cranei* Herre, 1935  
*Pomacentrus jenkinsi* Jordan and Evermann, 1903  
*Pomacentrus reidi* Fowler and Bean, 1928  
*Pomacentrus stigma* Fowler and Bean, 1928  
*Stegastes baldwini* Allen and Woods, 1980

#### PROCHILODONTIDAE

*Prochilodus maripicru* Eigenmann, 1912

#### PSEUDOCROMIDAE

*Nesiotes purpurascens* De Vis, 1884  
*Pseudochromis aurea marshallensis* Schultz, 1953  
*Pseudoplesiops sargenti* Schultz, 1953

#### PSILORHYNCHIDAE

*Psilorhynchus gracilis* Rainboth, 1983

#### RAJIDAE

*Pseudoraja fischeri* Bigelow and Schroeder, 1954  
*Raja hollandi* Jordan and Richardson, 1909

#### RHAMPHICHTHYIDAE

*Gymnorhamphichthys hypostomus* Ellis, 1912

#### RHINOCHIMAERIDAE

*Neoharriotta carri* Bullis and Carpenter, 1966

#### SALMONIDAE

*Coregonus stanleyi* Kendall, 1903  
*Leucichthys alpenae* Koelz, 1924  
*Leucichthys kiyi* Koelz, 1921  
*Leucichthys reighardi* Koelz, 1924  
*Oncorhynchus adonis* Jordan and McGregor, 1925

*Oncorhynchus ishikawae* Jordan and McGregor, 1925  
*Oncorhynchus kawamurae* Jordan and McGregor, 1925  
*Oncorhynchus rhodurus* Jordan and McGregor, 1925  
*Salmo bathoecetor* Meek, 1899  
*Salmo clarki declivifrons* Meek, 1899  
*Salmo clarki jordani* Meek, 1899  
*Salmo roosevelti* Evermann, 1906  
*Salvelinus imbrius* Jordan and McGregor, 1925  
*Salvelinus timagamiensis* Henn and Rinkensbach, 1925

#### SCARIDAE

*Cryptotomus crassiceps* Bean, 1906  
*Scarus laevis* Jordan and Evermann, 1903  
*Sparisoma rhomaleum* Meek and Hildebrand, 1928

#### SCHILBIDAE

*Physalia villiersi* Boulenger, 1912

#### SCIAENIDAE

*Menticirrhus focaliger* Ginsburg, 1952  
*Ophioscion punctatissimus* Meek and Hildebrand, 1925  
*Ophioscion venezuelae* Schultz, 1945  
*Otolithes lini* Herre, 1935  
*Pseudotolithus brunneolus* Jordan and Richardson, 1909  
*Sciaena bathytatos* Chao and Miller, 1975  
*Sciaena iharae* Jordan and Metz, 1913  
*Sciaena trewavasae* Chao and Miller, 1975  
*Stellifer colonensis* Meek and Hildebrand, 1925  
*Stellifer melanocheir* Eigenmann, 1917  
*Stellifer mindii* Meek and Hildebrand, 1925  
*Umbrina tumacoensis* Wilson, 1916

#### SCOMBRIDAE

*Pneumatophorus peruanus* Jordan and Hubbs, 1925

#### SCOPELARCHIDAE

*Scopelarchus beebe* Rofen, 1963  
*Scopelarchus sagax* Rofen, 1963

#### SCORPAENIDAE

*Brachirus bellus* Jordan and Hubbs, 1925  
*Pterois ellioti* Meek, 1897  
*Scorpaenopsis catocala* Jordan and Evermann, 1903  
*Sebastapistes hadiorufus* Herre, 1935  
*Sebastodes clavilatus* Starks, 1911

*Sebastodes deani* Starks, 1911  
*Sebastodes emphaeus* Starks, 1911  
*Sebastodes ijimae* Jordan and Metz, 1913  
*Sebastodes owstoni* Jordan and Thompson, 1914  
*Sebastodes thompsoni* Jordan and Hubbs, 1925  
*Thysanichthys evides* Jordan and Thompson, 1914

#### SCYLIORHINIDAE

*Pristiurus sauteri* Jordan and Richardson, 1909

#### SERRANIDAE

*Anthias conspicuus* Heemstra, 1973  
*Anthias mooreanus* Herre, 1935  
*Anthias woodsi* Anderson and Heemstra, 1980  
*Bodianus stellatus* Blosser, 1909  
*Centropristes melanus* Ginsburg, 1952  
*Cephalopholis boninius* Jordan and Thompson, 1914  
*Diplectrum labarum* Rosenblatt and Johnson, 1974  
*Diplectrum pacificum* Meek and Hildebrand, 1925  
*Entonanthias pascualis* Jordan and Tanaka, 1927  
*Epinephelus elongatus* Schultz, 1953  
*Epinephelus emoryi* Schultz, 1953  
*Epinephelus ionthas* Jordan and Metz, 1913  
*Epinephelus melanostigma* Schultz, 1953  
*Epinephelus spilotoceps* Schultz, 1953  
*Franzia ardens* Jordan and Thompson, 1914  
*Liopropoma mowbrayi* Woods and Kanazawa, 1951  
*Pikea aurora* Jordan and Evermann, 1903  
*Pikea mexicanus atlanticus* Schultz, 1958  
*Pikea mexicanus mexicanus* Schultz, 1958  
*Plectranthias garrupellus* Robins and Starck, 1961  
*Prionodes baldwini* Evermann and Marsh, 1899  
*Serraniculus pumilio* Ginsburg, 1952  
*Serranus chionaraia* Robins and Starck, 1961  
*Serranus maytagi* Robins and Starck, 1961  
*Symphysanodon berryi* Anderson, 1970  
*Symphysanodon octoactinus* Anderson, 1970  
*Ypsigramma lineata* Schultz, 1953

#### SILURIDAE

*Kryptopterus parvanalis* Inger and Chin, 1959  
*Ompok sabanus* Inger and Chin, 1959  
*Wallago maculatus* Inger and Chin, 1959

#### SISORIDAE

*Glyptothorax housei* Herre, 1942

#### SOLEIDAE

*Achirus fluviatilis* Meek and Hildebrand, 1928  
*Soleonassus finis* Eigenmann, 1912



## SOLENOTOMIDAE

*Solenostomus paegnius* Jordan and Thompson, 1914

## SPARIDAE

*Archosargus oviceps* Ginsburg, 1952  
*Diplodus bermudensis* Caldwell, 1965

## SQUALIDAE

*Squaliolus sarmenti* Noronha, 1926

## STERNOPTYCHIDAE

*Araiophos gracilis* Grey, 1961  
*Polyipnus nuttingii* Gilbert, 1905  
*Sonoda megalophthalma* Grey, 1959

## STERNOPYGIDAE

*Sternopygus dariensis* Meek and Hildebrand, 1916  
*Sternopygus pejateron* Schultz, 1949

## STICHAEIDAE

*Anoplarchus purpureus archolepis* Hubbs, 1926  
*Lumpenus longirostris* Evermann and Goldsborough, 1907

## SYNAPHOBRANCHIDAE

*Dysomma abyssale* Kamohara, 1938  
*Synaphobranchus oregoni* Castle, 1960

## SYNGNATHIDAE

*Bombonia uxorius* Herre, 1935  
*Corythoichthys elerae* Evermann and Seale, 1907  
*Corythoichthys polynotatus* Dawson, 1977  
*Doryrhamphus melanopleura cooki* Weed and Woods, 1953  
*Hippocampus brunneus* Bean, 1906  
*Hippocampus fisheri* Jordan and Evermann, 1903  
*Micrognathus suvensis* Herre, 1935  
*Siphostoma brevicaudum* Meek, 1904  
*Siphostoma eigenmanni* Wilson, 1916  
*Siphostoma elcapitanense* Meek and Hildebrand, 1914  
*Syngnathus mackayi nesiotus* Herald, 1942  
*Syngnathus scovelli makaxi* Herald and Dawson, 1972

## SYNODONTIDAE

*Synodus binotatus* Schultz, 1953  
*Synodus ulae* Schultz, 1953

## TERAPONIDAE

*Helotes lorentzi* Weber, 1910  
*Terapon habbema* Weber, 1910

## TETRAODONTIDAE

*Chonerhinus amabilis* Roberts, 1982  
*Chonerhinus nefastus* Roberts, 1982  
*Chonerhinus remotus* Roberts, 1982  
*Chonerhinus silus* Roberts, 1982  
*Spheroides asterias* Blosser, 1909  
*Sphoeroides kendalli* Meek and Hildebrand, 1928  
*Sphoeroides parvus* Shipp and Yerger, 1969  
*Tetraodon kretamensis* Inger, 1953

## TRACHICHTHYIDAE

*Hoplostethus occidentalis* Woods, 1973  
*Paratrachichthys argyrophanus* Woods, 1961

## TRACHIPTERIDAE

*Trachipterus fukuzakii* Fitch, 1964

## TRIACANTHODIDAE

*Parahollandia schmidtii* Woods, 1959

## TRICHOMYCTERIDAE

*Ochmacanthus flabelliferus* Eigenmann, 1912  
*Pygidium alternatum* Eigenmann, 1917  
*Pygidium banneaui* Eigenmann, 1912  
*Pygidium barbouri* Eigenmann, 1911  
*Pygidium bogotense* Eigenmann, 1912  
*Pygidium caliense* Eigenmann, 1912  
*Pygidium chapmani* Eigenmann, 1912  
*Pygidium conradi* Eigenmann, 1912  
*Pygidium davisi* Haseman, 1911  
*Pygidium dorsostriatum* Eigenmann, 1917  
*Pygidium gracilior* Eigenmann, 1912  
*Pygidium guianensis* Eigenmann, 1909  
*Pygidium hasemani* Eigenmann, 1914  
*Pygidium iheringi* Eigenmann, 1917  
*Pygidium latistriatum* Eigenmann, 1917  
*Pygidium paolence* Eigenmann, 1917  
*Pygidium proops parahybae* Eigenmann, 1918  
*Pygidium punctulatum piurae* Eigenmann, 1922  
*Pygidium reinhardti* Eigenmann, 1917  
*Pygidium santaeritae* Eigenmann, 1918  
*Pygidium septentrionale* Behre, 1928  
*Pygidium stellatum* Eigenmann, 1918  
*Pygidium straminium* Eigenmann, 1917  
*Pygidium striatum* Meek and Hildebrand, 1913  
*Pygidium triguttatum* Eigenmann, 1918  
*Pygidium vermiculatum* Eigenmann, 1917  
*Pygidium zonatum* Eigenmann, 1918  
*Scleronema operculatum* Eigenmann, 1917

*Vandellia hasemani* Eigenmann, 1918  
*Vandellia sanguinea* Eigenmann, 1917

#### TRIGLIDAE

*Bellator ribeiroi* Miller, 1965  
*Chelidonichthys ischyrius* Jordan and Thompson, 1914  
*Peristedion gilberti* Jordan, 1922  
*Peristedion greyae* Miller, 1967  
*Prionotus aspersus* Meek and Hildebrand, 1928

#### TRIPTERYGIIDAE

*Enneapterygius punctulatus* Herre, 1935  
*Enneapterygius waigiensis* Herre, 1935

#### UMBRIDAE

*Novumbra hubbsi* Schultz, 1929

#### URANOSCOPIDAE

*Zalescopus satsumae* Jordan and Hubbs, 1925  
*Zalescopus tosae* Jordan and Hubbs, 1925

#### XENCONGRIDAE

*Kaupichthys atlanticus* Böhlke, 1956  
*Kaupichthys brachychirus* Schultz, 1953

#### ZOARCIDAE

*Allolepis hollandi* Jordan and Hubbs, 1925  
*Lycodes tanakae* Jordan and Thompson, 1914  
*Rhigophila dearborni* DeWitt, 1962  
*Zestichthys tanakae* Jordan and Hubbs, 1925

#### Acknowledgments

This catalogue represents part of the junior author's matching commitment to National Science Foundation Grant DEB 82-04071 that financed in part the technical help needed for curation of

Field Museum's type collection; Terry Grande and Robert Schmitz (both at FMNH) assisted in that regard. We would especially like to thank Ian Chavez (Antioch College, Yellow Springs, Ohio) and Roy Yanong (Yale University, New Haven, Connecticut) for their assistance with library research and diverse other tasks. Rosetta Arrigo (FMNH) wrote the computer program that was used to print the type list and generously assisted with various other computer-related needs. Lynn Hobbs (FMNH) drafted the illustration. We also thank Robert K. Johnson and Robert M. Timm (FMNH) for constructive criticism of the manuscript. Numerous colleagues assisted by providing advice, information on manuscript names, and help with other problems.

#### Literature Cited

- BARTHOLOMEW, J. C., ET AL., EDs. 1980. The Times Atlas of the World. Comprehensive Edition, 6th ed. Times Books, New York.
- BIOSIS. 1981. Serial Sources for the BIOSIS Data Base, 1980. BioSciences Informative Service, Philadelphia.
- COLLETTE, B. B., AND E. A. LACHNER. 1976. Fish collections in the United States and Canada. *Copeia*, 1976(3): 625-644.
- GREY, M. 1947. Catalogue of the type specimens of fishes in Chicago Natural History Museum. *Fieldiana: Zool.*, 32(3): 109-205.
- HENN, A. W. 1928. II. List of types of fishes in the collection of the Carnegie Museum on September 1, 1928. *Ann. Carnegie Mus.*, 19(4): 38-39.
- NELSON, J. S. 1984. *Fishes of the World*, 2nd ed. John Wiley and Sons, New York.
- NITECKI, M. H. 1980. Featured institution—Field Museum of Natural History. *ASC Newsletter*, 8(5): 61-70.
- ROBINS, C. R., ET AL. 1980. A List of Common and Scientific Names of Fishes from the United States and Canada, 4th ed. Special Publication No. 12, American Fisheries Society, Bethesda, Md.

## Species Index

- abacopus*, *Glossogobius* 40, 96  
*abo*, *Apogon* 10, 86  
*abramoides*, *Astyanax* 12, 89  
*abuelo*, *Sovichthys* 80, 100  
*abyssale*, *Dysomma* 33, 103  
*acanthogaster*, *Deuterodon* 31, 90  
*acanthorhinus*, *Ostichthys* 63, 97  
*acerus*, *Paracrossochilus* 63, 94  
*achlamys*, *Gazza* 38, 97  
*acutiventralis*, *Alfaro* 7, 100  
*acutum*, *Cichlasoma* 25, 92  
*adiposalis*, *Pseudobagrus* 70, 88  
*adobe*, *Agosia* 6, 93  
*adonis*, *Oncorhynchus* 62, 101  
*aeneus costaricensis*, *Astyanax* 12, 89  
*affinis littoralis*, *Atherinops* 14, 87  
*Pseudocheirodon* 70, 91  
*Rhombatractus* 75, 98  
*afuerae multipunctatus*, *Malacotenus* 54, 97  
*aguadulce*, *Galeichthys* 36, 87  
*ahlstromi*, *Evermannella* 35, 95  
*akeleyi*, *Barbus* 15, 93  
*albater*, *Noturus* 61, 97  
*albeolus*, *Astyanax* 12, 89  
*albmarginatus*, *Pseudopimelodus* 71, 100  
*alepidota californiensis*, *Chaenopsis* 21, 89  
*alestes borneensis*, *Nematabramis* 60, 94  
*alfari*, *Cichlasoma* 25, 92  
*alfredae*, *Bryconamericus* 17, 90  
*aliata*, *Stewardia* 82, 92  
*aliceae*, *Argentina* 11, 87  
*alienus*, *Cypselurus* 31, 95  
*alleni*, *Tetragonopterus* 83, 92  
*alpenae*, *Leucichthys* 52, 101  
*alpha*, *Bryconamericus* 17, 90  
*alta*, *Chromis* 24, 101  
*Crenicichla* 28, 92  
*alternatum*, *Pygidium* 72, 103  
*alternus*, *Leporinus* 51, 86  
*altipinna*, *Pseudorasbora* 71, 94  
*Rhoadsia* 75, 91  
*altispinosa*, *Crenicara* 28, 92  
*altum*, *Sicydium* 79, 96  
*altus*, *Acestrorhynchus* 5, 89  
*alvarezii*, *Gambusia* 36, 100  
*amabilis*, *Chonerhinus* 24, 103  
*amates*, *Poecilia* 67, 101  
*amaurus*, *Bunocephalus* 19, 87  
*amazonae*, *Bleptonema* 16, 90  
*amblops*, *Felichthys* 36, 87  
*amphiloxus*, *Hemicetopsis* 44, 89  
*Nematobrycon* 60, 91  
*anale*, *Dorosoma* 33, 93  
*analis australis*, *Clinocottus* 27, 93  
*Hemigrammus* 44, 91  
*Piabina* 65, 91  
*anchovia*, *Agoniatas* 6, 89  
*andersoni*, *Corymbophanes* 28, 98  
*angelica*, *Emblemaria* 34, 89  
*anguillicauda*, *Liobagrus* 52, 88  
*anguillicaudatus tungting*, *Misgurnus* 58, 93  
*anisitsi*, *Aphyocharax* 9, 89  
*Pterygoplichthys* 72, 98  
*annae*, *Cheirodon* 23, 90  
*annectens hesperis*, *Priapichthys* 69, 101  
*anoplus*, *Barbus* 15, 93  
*anpinensis*, *Mugil* 58, 98  
*antillarum*, *Ancylopsetta* 8, 88  
*antoncichi*, *Cypselurus* 31, 95  
*anuli*, *Pimephales* 66, 94  
*aokii*, *Culter* 29, 94  
*aphanea*, *Algansea* 7, 93  
*aphelocheilus*, *Protomyzon* 70, 97  
*aporia*, *Gobiopsis* 41, 96  
*aporrhox*, *Lepophidium* 51, 99  
*aquilus*, *Lophiobagrus* 53, 88  
*archipelagicus*, *Hemiramphus* 45, 96  
*arcus*, *Leporinus* 51, 86  
*ardens*, *Franzia* 36, 102  
*ardesiaca*, *Nansenia* 59, 88  
*arenicola*, *Anchovia* 8, 95  
*arenidens*, *Solivomer* 80, 99  
*argentata*, *Ilisha* 49, 93  
*argenteus*, *Brycon* 19, 90  
*Trachurus* 84, 89  
*arge*, *Upeneus* 84, 99  
*argi*, *Centropyge* 21, 101  
*argyromus*, *Myripristis* 58, 97  
*argyrophanus Paratrachichthys* 64, 103  
*arleoi*, *Anduzedoras* 8, 95  
*armatus australis*, *Leptocottus* 52, 93  
*asper earli*, *Exerpes* 36, 97  
*aspersus*, *Prionotus* 69, 104  
*asterias*, *Spheroides* 80, 103  
*astrinus*, *Calliurichthys* 20, 88  
*atcafalayae vanhyningi*, *Signalosa* 79, 93  
*ateleostus*, *Nemaclinus* 60, 97  
*aterrimus*, *Ctenogobius* 29, 96  
*Hymenocephalus* 47, 98  
*atherodon*, *Optonurus* 63, 98  
*athi*, *Barbus* 15, 93  
*atlanticus*, *Kaupichthys* 50, 104  
*atracaudata*, *Deuterodon* 31, 90  
*atratoensis*, *Curimatus* 30, 93  
*atra*, *Phrynelox* 65, 86  
*atrilatus*, *Pseudaspius* 70, 94  
*atrimentatus*, *Teuthis* 83, 86  
*atripectoralis*, *Chromis* 24, 101  
*attenuatum*, *Chiostoma* 23, 87  
*aubynei*, *Pristella* 69, 91  
*aurata vallahica*, *Cobitis* 27, 93  
*auratus*, *Poecilobrycon* 68, 97  
*aurea marshallensis*, *Pseudochromis* 71, 101  
*aurocaudatus*, *Astyanax* 12, 89  
*aurocingulus*, *Aparrius* 9, 95  
*aurora*, *Pikea* 65, 102  
*australe*, *Corydoras* 27, 88  
*Pyrrhulina* 74, 97  
*australis*, *Ameiurus* 7, 97  
*avanhandavae*, *Pimelodella* 65, 100  
*awani*, *Aequidens* 5, 92  
*azonus*, *Pleurogrammus* 67, 96  
*aztecus*, *Notropis* 60, 94  
*azurifer*, *Aequidens* 5, 92  
*azygia*, *Farlowella* 36, 98  
*badia*, *Macgregorella* 54, 96  
*badiorufus*, *Sebastapistes* 78, 102  
*baldwini*, *Hemipteronotus* 45, 97  
*Prionodes* 69, 102  
*Stegastes* 81, 101  
*balsanus*, *Melaniris* 55, 87  
*balsas*, *Poeciliopsis* 68, 101  
*balteata*, *Vaimosa* 84, 96  
*banneai*, *Pygidium* 72, 103  
*bantolanensis*, *Barbus* 15, 93  
*barbata*, *Rhamdia* 74, 100  
*barbatus*, *Acheilognathus* 5, 93  
*Platysilurus* 67, 100  
*barbouri*, *Pygidium* 72, 103  
*barnardi*, *Pygidium* 64, 91  
*barnesi*, *Hypoatherina* 48, 87  
*bathoecetor*, *Salmo* 77, 102  
*bathoiketes*, *Porichthys* 69, 88  
*bathytatos*, *Sciaena* 78, 102  
*bayeri*, *Gymnothorax* 42, 99  
*beani*, *Heros* 46, 92  
*beauforti*, *Apogon* 10, 86  
*beebe*, *Scopelarchus* 78, 102  
*behrae*, *Brycon* 19, 90  
*belizensis*, *Anchoviella* 8, 95  
*bellus*, *Brachirus* 16, 102  
*beni*, *Creagrutus* 28, 90  
*Phenacogaster* 64, 91  
*benjamini*, *Entomocorus* 34, 87  
*bergi*, *Pseudaspius* 70, 94  
*berlangai*, *Opisthonema* 62, 93  
*bermudae*, *Branchiostoma* 17, 88  
*bermudensis*, *Antennarius* 9, 86  
*Diplodus* 32, 103  
*Hemiramphus* 45, 96  
*Rhegma* 75, 96  
*berndti*, *Myripristis* 58, 97  
*beroe*, *Lophiodes* 53, 98  
*berryi*, *Symphysanodon* 82, 102  
*beta*, *Bryconamericus* 17, 90  
*biaggii*, *Heterenchelys* 46, 96  
*bifasciatus*, *Cyprinodon* 31, 94  
*Hyphessobrycon* 48, 91  
*Parodon* 64, 99  
*bifrenatus*, *Barbus* 15, 93  
*bifurcus*, *Acanthophaelus* 4, 100  
*bigelowi*, *Bathypterois* 15, 92  
*bikiniensis*, *Gymnothorax* 42, 99  
*bilinearis*, *Callechelys* 19, 99  
*bilineata*, *Hasemania* 43, 91  
*bilineatus*, *Pygocentrus* 73, 91

- bilunulata*, *Kuhlia* 50, 97  
*bimaculata*, *Iridio* 49, 97  
*Nannacara* 59, 92  
*bimaculatus novae*, *Astyanax* 12, 89  
*paraguayensis*, *Astyanax* 12, 89  
*binghami*, *Ceratobranchia* 21, 90  
*biniplicata*, *Myxine* 59, 99  
*binotatus*, *Synodus* 82, 103  
*birtwistlei*, *Gobiella* 41, 96  
*blennioides*, *Characidium* 22, 90  
*blochi*, *Corydoras* 27, 88  
*blohmi*, *Epapterus* 34, 88  
*boehlkei*, *Barbuligobius* 15, 95  
*Rhinodoras* 75, 95  
*bogotense*, *Pygidium* 72, 103  
*bolivarensis*, *Hildadoras* 46, 95  
*boliviana*, *Pimelodella* 65, 100  
*boliviense*, *Cichlasoma* 25, 92  
*bonanox*, *Coryphaenoides* 28, 98  
*bondi coppenamensis*, *Corydoras* 27, 88  
*boninius*, *Cephalopholis* 21, 102  
*bonita*, *Gambusia* 37, 100  
*boquiae*, *Hemibrycon* 43, 91  
*boquillae*, *Cetopsorhamdia* 21, 100  
*borneensis*, *Neostethus* 60, 99  
*bosquianus longimaxilla*, *Chasmodes* 23, 88  
*bovalii*, *Poecilocharax* 68, 91  
*bowditchae*, *Myripristis* 58, 97  
*brachychirus*, *Kaupichthys* 50, 104  
*branchimelas*, *Stolephorus* 82, 95  
*branneri voulezi*, *Rhamdia* 74, 100  
*Rhamdia* 74, 100  
*brasiliensis iporangensis*, *Geophagus* 38, 92  
*itapicuruensis*, *Geophagus* 38, 92  
*brevicaudum*, *Siphostoma* 79, 103  
*breviceps*, *Rivulus* 76, 86  
*brevior*, *Auchenipterus* 14, 87  
*Chasmocranus* 22, 100  
*brevipinnis*, *Creagrutus* 28, 90  
*brevirostra*, *Anchovia* 8, 95  
*brevis*, *Hemigrammus* 44, 91  
*Thoracocharax* 83, 95  
*brevispinis*, *Lasiancistrus* 51, 98  
*Lophiobagrus* 53, 88  
*breviventralis*, *Alectis* 6, 89  
*broccae*, *Spintherobolus* 80, 92  
*browni*, *Moenkhausia* 57, 91  
*brucei*, *Argentina* 11, 87  
*brunneolus*, *Pseudolithus* 71, 102  
*brunneus*, *Ectenias* 33, 93  
*Hippocampus* 46, 103  
*Leporinus* 51, 86  
*Rivulus* 76, 86  
*bucca*, *Varicus* 85, 96  
*bullisi*, *Grimatrocetes* 42, 86  
*Holocentrus* 47, 97  
*burkei*, *Careproctus* 20, 93  
*cabrerae*, *Rhamdia* 74, 100  
*caetei*, *Apistogramma* 10, 92  
*caliense*, *Pygidium* 72, 103  
*caliensis*, *Gambusia* 37, 100  
*californiensis elongatus*, *Eucinos-tomus* 35, 95  
*calobrense*, *Cichlasoma* 25, 92  
*cana*, *Gambusia* 37, 100  
*candalarius*, *Profundulus* 70, 94  
*canensis*, *Acanthicus* 4, 98  
*capetensis*, *Loricaria* 53, 98  
*capito*, *Centrus* 21, 89  
*capitva*, *Goodea* 42, 96  
*carens*, *Barbus* 15, 93  
*caribbaea*, *Trichopsetta* 84, 88  
*carletoni*, *Leuciscus* 52, 94  
*carnegiei*, *Cnesterodon* 27, 100  
*Pseudancistrus* 70, 98  
*carpintis*, *Neotroplus* 59, 92  
*carri*, *Microdesmus* 56, 98  
*Neoharriotta* 60, 101  
*carychroa*, *Enchelycore* 34, 99  
*cascajalensis*, *Bryconamericus* 17, 90  
*Gambusia* 37, 101  
*catalepta*, *Dermatocheir* 31, 90  
*catenatum*, *Characidium* 22, 90  
*catocala*, *Scorpaenopsis* 78, 102  
*cauae*, *Roeboides* 76, 91  
*caucanum*, *Characidium* 22, 90  
*caucanus*, *Bryconamericus* 17, 90  
*Creagrutus* 28, 90  
*Gephyrocharax* 38, 90  
*Lasiancistrus* 51, 98  
*caudata*, *Poecilia* 67, 101  
*centrale*, *Cichlasoma* 25, 92  
*cephalotaenia*, *Plesiops* 67, 100  
*cerasina*, *Curimata* 30, 93  
*chagresi*, *Menidia* 56, 87  
*chamaizelus*, *Bunocephalus* 19, 87  
*chancellori*, *Odontorhamphus* 61, 96  
*chapmani*, *Cyclopium* 30, 87  
*Pygidium* 72, 103  
*chauliodus*, *Aplatophis* 10, 99  
*chavesi*, *Dorosoma* 33, 93  
*chiuhuahua*, *Notropis* 60, 94  
*chimantanus*, *Aequidens* 5, 92  
*Chasmocranus* 22, 100  
*chinssuensis*, *Pseudogobio* 71, 94  
*chionaraia*, *Serranus* 79, 102  
*chobensis*, *Aplocheilus* 10, 86  
*chocoensis*, *Bryconamericus* 17, 90  
*Diphyacantha* 32, 100  
*Gephyrocharax* 39, 90  
*chosenicus*, *Rhodeus* 75, 94  
*chryseres*, *Myripristis* 58, 97  
*chrysonemus*, *Pseudupeneus* 71, 99  
*chrysus*, *Eupomacentrus* 35, 101  
*cismontanus*, *Bryconamericus* 18, 90  
*citurensis*, *Oxyloricaria* 63, 98  
*clarias coprophagus*, *Pimelodus* 66, 100  
*clarkae*, *Adontosternarchus* 5, 87  
*clarkei*, *Scopelogys* 78, 99  
*clarki declivifrons*, *Salmo* 77, 102  
*jordani*, *Salmo* 77, 102  
*Euleptocheilichthys* 35, 95  
*clavilatus*, *Sebastodes* 78, 102  
*claytonii*, *Gobius* 41, 96  
*cliffordpopei*, *Gobius* 42, 96  
*coccogenis*, *Serrasalmus* 79, 92  
*colombianus*, *Bunocephalus* 19, 87  
*Hemibrycon* 43, 91  
*Heterandria* 46, 101  
*colombiensis*, *Thyrina* 83, 87  
*colonensis*, *Stellifer* 81, 102  
*combata*, *Antigonia* 9, 89  
*compressus*, *Hemigrammus* 44, 91  
*Odontognathus* 61, 93  
*Rivulus* 76, 86  
*conchos*, *Catostomus* 21, 89  
*conklini*, *Sphagebranchus* 80, 99  
*conradi*, *Pygidium* 72, 103  
*conserialis*, *Petersius* 64, 91  
*consocium*, *Chirostoma* 24, 87  
*conspicuum*, *Anthias* 9, 102  
*conventus*, *Bryconamericus* 18, 90  
*copei*, *Anduzedoras* 9, 95  
*corethromycter*, *Luciobrotula* 54, 99  
*coriparoides*, *Gobio* 41, 94  
*corlettei*, *Gnatholepis* 40, 96  
*corniger*, *Ogcocephalus* 61, 99  
*corumbae*, *Heterogramma* 46, 92  
*costaricensis*, *Gobiosoma* 41, 95  
*cranei*, *Pomacentrus* 68, 101  
*Thalassoma* 83, 97  
*crassiceps*, *Cryptotomus* 29, 102  
*crenatus*, *Stethapron* 82, 92  
*crenolepis*, *Allanetta* 7, 87  
*cubensis*, *Fundulus* 36, 94  
*Heterandria* 46, 101  
*cumberlandensis*, *Phoxinus* 65, 94  
*cuprea*, *Siphamia* 79, 87  
*curviventris*, *Psectrogaster* 70, 93  
*cyanelus*, *Mugilops* 58, 87  
*cycloidea*, *Ancylorsetta* 8, 88  
*cyclolepis*, *Parecbasis* 64, 91  
*cyclopus santanderensis*, *Astroblepus* 11, 87  
*cylindricus*, *Hemigrammus* 44, 91  
*cyri orientalis*, *Chondrostoma* 24, 94  
*daguae*, *Astyanax* 12, 89  
*Gobius* 42, 96  
*Hemiancistrus* 43, 98  
*Joturus* 50, 98  
*dariensis*, *Gambusia* 37, 101  
*dariensis*, *Hemibrycon* 43, 91  
*Oxyloricaria* 63, 98  
*Parodon* 64, 99  
*Sternopygus* 81, 103  
*davisi*, *Pygidium* 72, 103  
*dawsoni*, *Echiodon* 33, 89  
*dayi*, *Decapterus* 31, 89  
*deani*, *Sebastodes* 78, 102  
*dearborni*, *Cyprinodon* 31, 94  
*Rhigophila* 75, 104  
*decemlineatus*, *Awaous* 14, 95  
*declivirostris*, *Ogcocephalus* 62, 99  
*decoratus*, *Iridio* 49, 97  
*decurrens*, *Bryconamericus* 18, 90  
*delicata*, *Tagusa* 82, 88  
*demerarae*, *Auchenipterus* 14, 87  
*dentatus*, *Bryconamericus* 18, 90

- depressirostris*, *Pseudorasbora* 71, 94  
*depressus*, *Bunocephalus* 19, 87  
*derbyi*, *Plecostomus* 67, 98  
*deuterodonoides caudovittatus*, *Bryconamericus* 18, 90  
*Bryconamericus* 18, 90  
*devenanzii*, *Adontosternarchus* 5, 87  
*diaboli*, *Dionda* 32, 94  
*dialepturus*, *Cheirodon* 23, 90  
*diazii*, *Zoogoneticus* 86, 96  
*dilatatus*, *Spirinchus* 81, 99  
*diquensis*, *Bryconamericus* 18, 90  
*discus*, *Acestridium* 4, 98  
*dispar*, *Pempheris* 64, 99  
*distethommatum*, *Callionymus* 19, 88  
*dorsatum*, *Cichlasoma* 25, 92  
*dorsocellata*, *Crenicichla* 28, 92  
*dorsostriatum*, *Pygidium* 72, 103  
*dubius*, *Bathypetris* 15, 92  
*duopunctata*, *Aequidens* 5, 92  
*duragenys*, *Hyphessobrycon* 48, 91  
*ebelingi*, *Gonostoma* 42, 93  
*ebisui*, *Malacotenus* 54, 97  
*ectenurus*, *Leptocephalus* 52, 93  
*ecuadoriensis*, *Hyphessobrycon* 48, 91  
*Leporinus* 51, 86  
*Parodon* 64, 99  
*edwardraneyi*, *Notropis* 60, 94  
*efatensis*, *Gymnothorax* 42, 99  
*eidolon*, *Psychichthys* 71, 92  
*eigenmannia*, *Anchovia* 8, 95  
*eigenmanni*, *Chaenothorax* 22, 88  
*Cheirodon* 23, 90  
*Cichlasoma* 25, 92  
*Fitzroyia* 36, 97  
*Parapelecus* 63, 94  
*Phallopteryx* 64, 101  
*Pimelodella* 66, 100  
*Siphostoma* 79, 103  
*elcapitanense*, *Siphostoma* 80, 103  
*elegans erolli*, *Gibbonsia* 39, 93  
*montereyensis*, *Gibbonsia* 39, 93  
*velifera*, *Gibbonsia* 39, 93  
*Ammocryptocharax* 7, 89  
*Iridio* 49, 97  
*Neoheterandria* 60, 101  
*elerae*, *Corythoichthys* 28, 103  
*elliotti*, *Pterois* 71, 102  
*Thorichthys* 83, 92  
*ellisae*, *Corydoras* 27, 88  
*elongata*, *Anchovia* 8, 95  
*elongatus*, *Bramocharax* 17, 90  
*Carpiodes* 20, 89  
*Epinephelus* 34, 102  
*elymus aphrozoicus*, *Atherion* 14, 87  
*freyi*, *Atherion* 14, 87  
*emoryi*, *Epinephelus* 34, 102  
*emphaeus*, *Sebastodes* 78, 102  
*engraulis*, *Hemiculterella* 44, 94  
*ensifer*, *Gladiogobius* 39, 96  
*Holocentrus* 47, 97  
*eos*, *Hyphessobrycon* 48, 91  
*eques*, *Cheirocerus* 23, 100  
*erimyzonops*, *Dionda* 32, 94  
*erionema*, *Scolopsis* 78, 99  
*eristigma*, *Belligobio* 16, 94  
*erythrozonus*, *Hemigrammus* 44, 91  
*erythrurus*, *Aphyocharax* 9, 89  
*Poecilobrycon* 68, 97  
*essequibensis*, *Asiyanax* 12, 89  
*Cynopotamus* 31, 90  
*Leptoglanis* 52, 100  
*ethelwynnae*, *Gobiocichla* 41, 92  
*eugenius*, *Quisquilus* 74, 96  
*evermanni*, *Atherina* 14, 87  
*Cichlasoma* 25, 92  
*Diapterus* 32, 95  
*Ophichthys* 62, 99  
*evides*, *Encaetra* 34, 95  
*Hemipteronotus* 45, 97  
*Thysanichthys* 83, 102  
*evurus*, *Etelis* 35, 98  
*exilis*, *Trachelyichthys* 83, 88  
*falcatus*, *Glaridichthys* 40, 101  
*fasciata*, *Gambusia* 37, 101  
*fasciatus*, *Gastromyzon* 38, 97  
*fasciata*, *Crossostoma* 29, 93  
*fasciolatus*, *Phallotorynus* 64, 101  
*ferdebueni*, *Poblana* 67, 87  
*fijiensis*, *Halichoeres* 43, 97  
*filamentosa latiuscula*, *Loricaria* 53, 98  
*seminuda*, *Loricaria* 53, 98  
*filamentosus*, *Macropodus* 54, 88  
*fimbriata*, *Loricaria* 53, 98  
*finis*, *Soleonotus* 80, 102  
*fischeri*, *Pseudoraja* 71, 101  
*fisheri*, *Hippocampus* 46, 103  
*Labeo* 50, 94  
*Trachycorystes* 84, 88  
*fitzsimonsi*, *Barbus* 15, 93  
*flabelliferus*, *Ochmacanthus* 61, 103  
*flammeus*, *Mulloidops* 58, 98  
*flavater*, *Noturus* 61, 97  
*floridae*, *Branchiostoma* 17, 88  
*fluviatilis*, *Achirus* 5, 102  
*Disparichthys* 32, 89  
*Gobiesox* 41, 95  
*Leptophilypnus* 52, 95  
*Neotroplus* 59, 92  
*focaliger*, *Menticirrhus* 56, 102  
*fontinalis*, *Cyprinodon* 31, 94  
*forlonensis*, *Nototropis* 60, 94  
*formosa*, *Rineloricaria* 76, 98  
*formosae*, *Liza* 53, 98  
*formosanus*, *Caranx* 20, 89  
*Rhinogobius* 75, 96  
*Scomberoides* 78, 89  
*fosteri*, *Priapichthys* 69, 101  
*fowleri*, *Cephalosilurus* 21, 100  
*fragilis*, *Sceptris* 78, 87  
*franciscensis*, *Fowlerina* 36, 90  
*franciscoensis*, *Phenacogaster* 65, 91  
*freemani*, *Pristipomoides* 69, 98  
*frenata*, *Brachyglanis* 17, 100  
*frenatus*, *Astroblepus* 11, 87  
*Rivulus* 76, 86  
*fria*, *Poecilia* 67, 101  
*frontale*, *Cichlasoma* 25, 92  
*frontatus*, *Microspathodon* 57, 101  
*fukuzakii*, *Trachipterus* 83, 103  
*fulvus*, *Gobiesox* 41, 95  
*fundicola*, *Suruga* 82, 96  
*furcatus*, *Oreias* 63, 93  
*fuscoguttatus*, *Uropterygius* 84, 99  
*fuscolineata*, *Siphamia* 79, 87  
*fuscomaculata*, *Rabula* 74, 99  
*gabonensis*, *Grasseichthys* 42, 97  
*gabunensis*, *Ischnomembras* 50, 87  
*galusdae*, *Cheirodon* 23, 90  
*garmani*, *Girardinus* 39, 101  
*garrupellus*, *Plectranthias* 67, 102  
*geniornatus*, *Cryptocentrus* 29, 96  
*georgei*, *Argentina* 11, 87  
*gigas*, *Diaphus* 32, 99  
*Malacotenus* 54, 97  
*gilberti*, *Careproctus* 20, 93  
*Cololrhynchus* 27, 98  
*Formosania* 36, 94  
*Peristedion* 64, 104  
*gillii*, *Curimatus* 30, 93  
*gimbeli*, *Porotergus* 69, 87  
*glandulifer*, *Diaphus* 32, 99  
*glaucum*, *Stizostedion* 82, 100  
*gloverensis*, *Triathalassothia* 84, 88  
*gracilicauda*, *Gymnapogon* 42, 86  
*gracilior*, *Pygidium* 72, 103  
*gracilis*, *Acheilognathus* 5, 93  
*Araiphos* 11, 103  
*Psilorhynchus* 71, 101  
*Tomeurus* 83, 101  
*grammica*, *Aphyodite* 10, 89  
*grammodes*, *Cichlasoma* 25, 92  
*granadense*, *Cichlasoma* 25, 92  
*grandis*, *Asiyanax* 12, 89  
*Pomadasys* 68, 96  
*grandisquamis*, *Amblypharyngodon* 7, 93  
*granti*, *Leporinus* 51, 86  
*greenei*, *Notropis* 61, 94  
*greenfieldi*, *Acanthemblemaria* 4, 89  
*greyae*, *Bathylagus* 15, 88  
*Gillellus* 39, 94  
*Peristedion* 64, 104  
*greyi*, *Bembrops* 16, 100  
*griffini*, *Pimelodella* 66, 100  
*grimaldii*, *Hymenocephalus* 47, 98  
*grinnelli*, *Physiculus* 65, 98  
*griseus*, *Loricaria* 53, 98  
*grixiavii micrescens*, *Astroblepus* 11, 87  
*grosvenori*, *Bryconamericus* 18, 90  
*guaporensis*, *Aequidens* 5, 92  
*Asiyanax* 12, 89  
*Metynnis* 56, 91  
*Microschemobrycon* 57, 91  
*guianensis*, *Acestrorhynchus* 5, 89  
*Ageneiosus* 6, 86  
*Asiyanax* 12, 89  
*Brachyhalcinus* 17, 90  
*Pygidium* 73, 103  
*Stolephorus* 82, 95  
*guuja*, *Thyrina* 83, 87

- gymnocephalus*, *Dormitator* 33, 95  
*gymnodontus*, *Psalidodon* 70, 91  
*gymnogaster lagoichthys*, *Loricaria* 53, 98  
*Loricaria* 53, 98  
*gymnogenys*, *Asryanax* 13, 89  
*gymnotus*, *Porotergus* 69, 87  
  
*habbema*, *Terapon* 83, 103  
*hankinsoni*, *Hybognathus* 47, 94  
*harringtonensis*, *Leptocephalus* 52, 93  
*harrisoni*, *Poecilobrycon* 68, 97  
*hartwegi*, *Cichlasoma* 25, 92  
*hasemani*, *Apareiodon* 9, 99  
*Asryanax* 13, 89  
*Farlowella* 36, 98  
*Heterandria* 46, 101  
*Pimelodella* 65, 100  
*Pygidium* 73, 103  
*Sternarchus* 81, 87  
*Vandellia* 84, 104  
*hastatus*, *Odontostilbe* 61, 91  
*hedricki*, *Cichlasoma* 25, 92  
*helleri*, *Anchoiella* 8, 95  
*Barbus* 15, 94  
*Hemibrycon* 43, 91  
*hemigrammus*, *Aphyocheirodon* 10, 89  
*Asiphonichthys* 11, 89  
*hemius*, *Plecostomus* 67, 98  
*henni*, *Brycon* 19, 90  
*Bryconamericus* 18, 90  
*hepsetus colonensis*, *Anchoa* 7, 95  
*heraldi*, *Centropyge* 21, 101  
*herwerdenii*, *Eleotris* 33, 95  
*heterandria*, *Pseudocorynopoma* 71, 91  
*heterochir*, *Gambusia* 37, 101  
*heterodon*, *Eleotris* 33, 95  
*Holesthes* 47, 91  
*heterolepis*, *Cynoglossus* 30, 93  
*heteromorphus*, *Corydoras* 27, 88  
*heteropleurus*, *Pimelodus* 66, 100  
*heterostomus*, *Probolodus* 70, 91  
*heterura*, *Compsura* 27, 90  
*heterurus*, *Asryanax* 13, 89  
*hiatti*, *Acanthoplesiops* 4, 86  
*higmani*, *Lonchopisthus* 53, 99  
*hildebrandi*, *Hyporhamphus* 49, 96  
*Roeboides* 76, 91  
*Sicydium* 79, 96  
*hingi*, *Homaloptera* 47, 97  
*hoignei*, *Pterolebias* 71, 86  
*hollandi*, *Allolepis* 7, 104  
*Areliscus* 11, 93  
*Imparfinis* 49, 100  
*Limia* 52, 101  
*Myctophum* 58, 99  
*Raja* 74, 101  
*Serrasalmo* 79, 92  
*Spinibarbus* 80, 94  
*holmiae*, *Rivulus* 76, 86  
*hondae*, *Pseudoperilampus* 71, 94  
*horiae*, *Vaimosa* 84, 96  
*housei*, *Glyptothorax* 40, 102  
  
*hubbsi hubbsi*, *Malacotenus* 55, 97  
*polyporus*, *Malacotenus* 54, 97  
*Novumbra* 61, 104  
*hudsoni*, *Callionymus* 20, 88  
*humeralis gracilior*, *Serrasalmo* 79, 92  
*hurtadoi*, *Gambusia* 37, 101  
*hyltoni*, *Emblemaria* 34, 89  
*hypesson*, *Bryconamericus* 18, 90  
*hyposagmatus*, *Ophichthus* 62, 99  
*hypostomus*, *Gymnorhamphichthys* 42, 101  
  
*iguassuensis*, *Crenicichla* 29, 92  
*iharae*, *Sciaena* 78, 102  
*iheringi*, *Microglanis* 57, 100  
*Pygidium* 73, 103  
*ijimai*, *Gnathopogon* 40, 94  
*ijimae*, *Sebastodes* 79, 102  
*Synchiropus* 82, 89  
*ijimai*, *Astronesthes* 12, 87  
*illecebrosus*, *Alburnops* 6, 93  
*illustris*, *Eumegistus* 35, 88  
*imbrius*, *Salvelinus* 77, 102  
*imswe*, *Varicus* 85, 96  
*incompta*, *Curimata* 30, 93  
*inconspicua*, *Apistogramma* 10, 92  
*inequalis*, *Glandulocauda* 39, 91  
*inermis*, *Platophrys* 66, 88  
*infans*, *Gambusia* 37, 101  
*insularum cedroscensis*, *Atherinops* 14, 87  
*guadalupae*, *Atherinops* 14, 87  
*Atherina* 14, 87  
*integripinnis zacae*, *Paraclinus* 63, 97  
*intermedius*, *Gephyrocharax* 39, 90  
*Squalogadus* 81, 98  
*interruptus ibicuhiensis*, *Cheirodon* 23, 90  
*intonsus*, *Ernstichthys* 35, 87  
*ionthas*, *Epinephelus* 34, 102  
*iota*, *Hemigrammus* 44, 91  
*irinae*, *Pseudanos* 70, 86  
*ischanus*, *Icticus* 49, 99  
*ischyrus*, *Chelidonichthys* 23, 104  
*ishikawae*, *Gnathopogon* 40, 94  
*Oncorhynchus* 62, 102  
*ishikawai*, *Caranx* 20, 89  
*issororoensis*, *Curimatus* 30, 93  
*isthmensis*, *Eleotris* 33, 95  
*itapicuruensis*, *Apareiodon* 9, 99  
*Pimelodella* 65, 100  
  
*jaguarensis*, *Crenicichla* 29, 92  
*jalapae*, *Xiphophorus* 85, 101  
*japonica*, *Eviota* 36, 96  
*jauruensis*, *Farlowella* 36, 98  
*jeb*, *Calamopteryx* 19, 99  
*jenkinsi*, *Pomacentrus* 68, 101  
*jimi*, *Parotocinclus* 64, 98  
*joaquinae*, *Puntius* 72, 94  
*johnsoni*, *Cirriacacula* 26, 99  
*jordani*, *Chirostoma* 24, 87  
*Conger* 27, 93  
*Labeo* 50, 94  
  
*jordani*, *Lampanyctus* 50, 99  
*Pantosteus* 63, 89  
*Trachinotus* 83, 89  
*jusanensis*, *Achahara* 4, 93  
  
*kabia*, *Melanotaenia* 56, 98  
*kabiliensis*, *Brachygnathus* 17, 95  
*kanazawai*, *Dicrolene* 32, 99  
*kanganamanensis*, *Arius* 11, 87  
*kasawae*, *Iburiella* 49, 86  
*katsukii*, *Stellistius* 81, 96  
*kawamurae*, *Oncorhynchus* 62, 102  
*keithi*, *Mastacembelus* 55, 98  
*kendalli*, *Sphoeroides* 80, 103  
*kennedyi*, *Hemigrammus* 45, 91  
*kesleri banaticus*, *Gobio* 41, 94  
*kikuchii*, *Phoxiscus* 65, 94  
*kiyi*, *Leucichthys* 52, 101  
*knightii*, *Gnatholepis* 40, 96  
*koelzi*, *Crossochir* 29, 95  
*kompi*, *Asryanax* 13, 89  
*koragensis*, *Glossogobius* 40, 96  
*kretamensis*, *Tetraodon* 83, 103  
*kulambangrae*, *Petroscirtus* 64, 88  
*kumae*, *Monomitus* 58, 99  
*kumperae*, *Ancylorsetta* 8, 88  
*kuroiwa*, *Tridentiger* 84, 96  
*kurumeus*, *Rhodeus* 75, 94  
*kymosemeum*, *Porichthys* 69, 88  
  
*labarcae*, *Chirostoma* 24, 87  
*labarum*, *Diplectrum* 32, 102  
*labiosus*, *Carpiodes* 20, 89  
*lacortei*, *Nematobrycon* 60, 91  
*lafa*, *Ambassis* 7, 89  
*lamprotaenia*, *Anchoa* 8, 95  
*lanceolatus*, *Rivulus* 76, 86  
*landoni*, *Pterobrycon* 71, 91  
*langi*, *Alestes* 6, 89  
*laterale*, *Characidium* 22, 90  
*laticauda typhla*, *Rhamdia* 74, 100  
*laticeps australis*, *Pimelodella* 66, 100  
*Pimelodella* 66, 100  
*Schizodontopsis* 78, 86  
*latidens*, *Astroblepus* 11, 87  
*Landonia* 50, 91  
*latifasciatus*, *Eleotris* 34, 95  
*latipunctata*, *Gambusia* 37, 101  
*Poecilia* 67, 101  
*latistriatum*, *Pygidium* 73, 103  
*latus*, *Diaphus* 32, 99  
*lauia*, *Scarus* 78, 102  
*leavelli*, *Ctenogobius* 29, 96  
*lentiginosus*, *Doras* 33, 95  
*Labrisomus* 50, 97  
*lepidogaster*, *Gastromyzon* 38, 97  
*lepidura hasemani*, *Moenkhausia* 57, 91  
*leporhinus*, *Hemidoras* 44, 95  
*leptocaulus*, *Apogon* 10, 86  
*leptorhynchus*, *Sternarchus* 81, 87  
*lermae*, *Skiffia* 80, 96  
*letonai*, *Priapichthys* 69, 101  
*leucomelas*, *Chaetostomus* 22, 98  
*leucostigma*, *Gymnothorax* 43, 99

- levis*, *Hemieleotris* 44, 95  
*lignus*, *Hypsoblennius* 49, 88  
*limbaughi*, *Chromis* 24, 101  
*lineata*, *Ypsigramma* 85, 102  
*lini*, *Otolithes* 63, 102  
*linelli*, *Leptodoras* 52, 95  
*liolepis*, *Euctenogobius* 35, 96  
*lisa*, *Menidia* 56, 87  
*lithoides*, *Lithoxus* 53, 98  
*lithurgicus*, *Ancistrus* 8, 98  
*longifilis*, *Gnathopogon* 40, 94  
*longior*, *Chasmocranus* 23, 100  
*longirostris*, *Lumpenus* 54, 103  
*lorentzi*, *Helotes* 43, 103  
*lorenzi*, *Abudefduf* 4, 101  
*luigiana*, *Rhamdia* 74, 100  
*lunatus*, *Holacanthus* 46, 101  
*lutzi*, *Heterandria* 46, 101  
*luzonensis*, *Aplocheilichthys* 10, 86  
*lyolepis*, *Stolephorus* 82, 95  
*lyriformis*, *Agmus* 6, 87  
  
*mabahiss*, *Vinciguerra* 85, 100  
*macilentus*, *Hemicetopsis* 44, 89  
*mackayi nesiotis*, *Syngnathus* 82, 103  
*macmasteri*, *Apistogramma* 10, 92  
*macrocephalus catherinae*, *Mniertes* 57, 97  
*macrolepidota sabana*, *Hampala* 43, 94  
*macrolepis*, *Eleotris* 34, 95  
*Heterocharax* 46, 91  
*Roeboides* 76, 91  
*macrosquamis*, *Sargocentron* 77, 97  
*macturki*, *Pimelodella* 66, 100  
*maculatus*, *Wallago* 85, 102  
*maculipinnis*, *Pterolebias* 72, 86  
*maculocephala*, *Cascadura* 21, 88  
*maculosus*, *Aspidoras* 11, 88  
*Tylobronchus* 84, 92  
*madeirae*, *Cheirodon* 23, 90  
*madeirensis*, *Ageneiosus* 6, 86  
*maderensis*, *Diplogonurus* 32, 95  
*magdalena*, *Astyanax* 13, 90  
*Creagrutus* 28, 90  
*Hoplosternum* 47, 88  
*Rivulus* 76, 86  
*Roeboides* 77, 91  
*Thoracocharax* 83, 95  
*Xiliphius* 85, 87  
*magdalenensis*, *Bryconamericus* 18, 90  
*majimae*, *Gnathopogon* 40, 94  
*majua majua*, *Jenkinsia* 50, 93  
*woodsii*, *Jenkinsia* 50, 93  
*malaris*, *Clarias* 26, 93  
*malayana*, *Antigonia* 9, 89  
*malayanus*, *Myxus* 59, 98  
*malekulae*, *Ctenogobius* 29, 96  
*malua*, *Boroda* 16, 95  
*margaritae mexicanus*, *Malacotenus* 55, 97  
*marginatus*, *Hemigrammus* 45, 91  
*Nannostomus* 59, 97  
*mariae*, *Acanthopthalmus* 4, 93  
*mariae*, *Aequidens* 5, 92  
*Sternarchus* 81, 87  
*marijeanae*, *Gobiesox* 41, 95  
*marionae*, *Astyanax* 13, 90  
*Bathytrophus* 16, 92  
*maripicru*, *Prochilodus* 70, 101  
*marmoratus*, *Ageneiosus* 6, 86  
*marquesensis*, *Sardinella* 77, 93  
*marshallensis*, *Rabula* 74, 99  
*martini*, *Salarias* 77, 88  
*maxillaris*, *Hasemanian* 43, 91  
*mayoloi*, *Hemiancistrus* 43, 98  
*maytagi*, *Serranus* 79, 102  
*medirastre*, *Opisthonema* 62, 93  
*meeki*, *Cichlasoma* 26, 92  
*Cyprinodon* 31, 94  
*Holocentrus* 47, 97  
*Roeboides* 77, 91  
*Thyrina* 83, 87  
*megalopsis*, *Gastromyzon* 38, 97  
*Monolene* 58, 88  
*megalophtalma*, *Sonoda* 80, 103  
*megalops*, *Pimelodella* 66, 100  
*megalopter*, *Megalampodus* 55, 91  
*megalostictus*, *Phenacogaster* 65, 91  
*megalostomus*, *Hysteronotus* 49, 91  
*melampodus*, *Apogonichthys* 10, 86  
*melandetus*, *Odontostilbe* 61, 91  
*melanocheir*, *Gephyrocharax* 39, 90  
*Stellifer* 81, 102  
*melanogenys*, *Glandulocauda* 39, 91  
*melanopleura cooki*, *Doryrhamphus* 33, 103  
*Glandulocauda* 39, 91  
*melanopleurus*, *Hyphessobrycon* 45, 91  
*melanopterus*, *Hemigrammus* 48, 91  
*Hyporhamphus* 49, 96  
*melanopus*, *Leiogaster* 51, 88  
*melanostigma*, *Epinephelus* 35, 102  
*melanotus*, *Aphyocharax* 10, 89  
*melanura*, *Hasemanian* 43, 91  
*melanus*, *Centropomus* 21, 102  
*melanzonus*, *Acanthophtalmus* 4, 100  
*Creagrutus* 28, 90  
*melas*, *Ancistrus* 8, 98  
*Brachyglanis* 17, 100  
*meridiana*, *Chromis* 24, 101  
*merretti*, *Polyacanthonotus* 68, 99  
*metae*, *Astyanax* 13, 90  
*Charax* 22, 90  
*Curimatus* 30, 93  
*Moenkhausia* 57, 91  
*Pimelodella* 66, 100  
*metzi*, *Gibbonsia* 39, 93  
*mexicana campi*, *Signalosa* 79, 93  
*Aztecula* 14, 93  
*mexicanus atlanticus*, *Pikea* 65, 102  
*mexicanus*, *Pikea* 65, 102  
*Ameiurus* 7, 97  
*Gobiesox* 41, 95  
*meyeri*, *Iridio* 50, 97  
*mezquital*, *Chirostoma* 24, 87  
*microctenus*, *Ancylosetta* 8, 88  
*microdon*, *Cheirodon* 23, 90  
*Loricaria* 53, 98  
*microlepis*, *Acanthocharax* 4, 89  
*Astyanax* 13, 90  
*micropocis*, *Hemidoras* 44, 95  
*microps*, *Imparfinis* 49, 100  
*Pogonoleotris* 68, 95  
*Rhamdia* 74, 100  
*micropterus*, *Megalampodus* 55, 91  
*micropunctatus*, *Lampanyctus* 50, 99  
*microstictus*, *Phenacogaster* 65, 91  
*microstomus*, *Carpododes* 21, 89  
*Hemidoras* 44, 95  
*Iridio* 50, 97  
*Oligobrycon* 62, 91  
*milleri*, *Cichlasoma* 26, 92  
*minacae*, *Gila* 39, 94  
*mindii*, *Microeleotris* 56, 95  
*Stellifer* 81, 102  
*miniatus*, *Zoogoneticus* 86, 96  
*minimus*, *Hyphessobrycon* 48, 91  
*Nannostomus* 59, 97  
*minor*, *Hyphessobrycon* 48, 91  
*minutulus*, *Synchiropus* 82, 89  
*minutus*, *Archicheir* 11, 97  
*Evorthodus* 36, 96  
*Hemicetopsis* 44, 89  
*Hemigrammus* 45, 91  
*Microbrycon* 56, 91  
*Microgenys* 56, 91  
*mirini*, *Hyphessobrycon* 49, 100  
*misakius*, *Chasmias* 22, 95  
*mittchilli diaphana*, *Anchoa* 8, 95  
*miyakamii*, *Caranx* 20, 89  
*mizolepis fukien*, *Misgurnus* 57, 93  
*grangeri*, *Misgurnus* 57, 93  
*modestus*, *Pseudaspis* 70, 94  
*moenkhausii*, *Poecilurichthys* 68, 91  
*mohoiity yunnan*, *Misgurnus* 57, 93  
*mojarra*, *Cichlasoma* 26, 92  
*mola*, *Metynnus* 56, 91  
*mollispinis mollispinis*, *Lepidomeda* 51, 94  
*montana*, *Rhamdella* 74, 100  
*montanum*, *Sicydium* 79, 96  
*monticola*, *Algansea* 7, 93  
*montium*, *Rivulus* 76, 86  
*mooreanus*, *Anthias* 9, 102  
*Myripristis* 58, 97  
*morawhannae*, *Curimatus* 30, 93  
*moreirai*, *Rhamdiopsis* 75, 100  
*morii*, *Sarcocheilichthys* 77, 94  
*mossambicus*, *Chromis* 27, 92  
*Clarias* 26, 93  
*mowbrayi*, *Grammonus* 42, 99  
*Liopropoma* 52, 102  
*Rhinogobius* 75, 96  
*mucronatus*, *Astyanax* 13, 90  
*multiporosus*, *Labrisomus* 50, 97  
*multiradiatus*, *Characodon* 22, 96  
*multispinis*, *Ctenopoma* 29, 86  
*mundus*, *Orestias* 63, 94  
*munitus*, *Noturus* 61, 97  
*muskoka*, *Notropis* 61, 94

- mutator*, *Astyanax* 13, 90  
*myersi*, *Amblygobius* 7, 95  
*Ekemblemaria* 33, 89  
*myriaster*, *Porichthys* 69, 88  
*myxodermus*, *Ctenogobius* 29, 95
- nantoenis*, *Liobagrus* 52, 88  
*nasus*, *Cetopsorhamdia* 21, 100  
*nasuta*, *Rhamdia* 74, 100  
*nasutus*, *Acestrorhynchus* 5, 89  
*Astyanax* 13, 90  
*nazas*, *Nototropis* 60, 94  
*nefastus*, *Chonerhinus* 24, 103  
*nemacheir*, *Nannorhamdia* 59, 100  
*neumanni*, *Fundulus* 36, 94  
*nicholsi*, *Blennius* 16, 88  
*nigra*, *Emblemaria* 34, 89  
*nigrescens*, *Pseudancistrus* 70, 98  
*nigripinnis woodsi*, *Cotylis* 28, 95  
*nigratum*, *Cichlasoma* 26, 92  
*nigrofasciatus*, *Apogon* 10, 86  
*nigrolineatus*, *Pseudauchenipterus* 70, 88  
*nigroventralis*, *Gambusia* 37, 101  
*niveilatus*, *Xyrichtys* 85, 97  
*norae*, *Gibbonsia* 39, 93  
*notemigonoides*, *Astyanax* 13, 90  
*notomelas*, *Cheirodon* 23, 90  
*Pimelodella* 66, 100  
*notospilus*, *Hemiodoras* 44, 95  
*notropoides*, *Creagrutus* 28, 90  
*nouhuysi*, *Atherinichthys* 14, 87  
*novae*, *Bryconamericus* 18, 90  
*novaeaguineae*, *Copidoglanis* 27, 100  
*novemmaculosus*, *Alticus* 7, 88  
*nox*, *Arius* 11, 87  
*nudus*, *Gerhardinus* 39, 96  
*nuttingii*, *Polyipnus* 68, 103
- oatesii*, *Barbus* 15, 94  
*Mastacembelus* 55, 98  
*oaxacae*, *Fundulus* 36, 94  
*Rhamdia* 74, 100  
*obeyense*, *Etheostoma* 35, 100  
*obliquus*, *Lamprologus* 50, 92  
*obscurus*, *Brycon* 19, 90  
*occidentalis*, *Hoplostethus* 47, 103  
*Lepomis* 51, 89  
*Roeboidea* 77, 92  
*Starksia* 81, 97  
*ocellatum*, *Mylossoma* 58, 91  
*ocellatus*, *Callogobius* 20, 95  
*Poecilobrycon* 68, 97  
*octoactinus*, *Symphysanodon* 82, 102  
*oculomaculatus*, *Gnathodentex* 40, 98  
*ohiensis*, *Alosa* 7, 93  
*oligolinx*, *Rhizoprionodon* 75, 89  
*olomina*, *Priapichthys* 69, 100  
*opalescens*, *Ctenogobius* 29, 96  
*operculatum*, *Scleronema* 78, 103  
*orbiculatus*, *Astronotus* 12, 92  
*orcuttii*, *Phoxinus* 65, 94  
*oregona*, *Draconetta* 33, 95
- oregoni*, *Synaphobranchus* 82, 103  
*orientalis americana*, *Epinnula* 35, 95  
*pacifica*, *Epinnula* 35, 95  
*Dasyscopelus* 31, 99  
*ornatum gephyrum*, *Cichlasoma* 26, 92  
*ortholepis*, *Bryconamericus* 18, 90  
*orthonops*, *Hemiodus* 45, 96  
*orthonotus*, *Cyprinodon* 31, 94  
*orthus*, *Hemigrammus* 45, 91  
*ortmanni*, *Heterogramma* 46, 92  
*Pimelodus* 66, 100  
*osakae*, *Netuma* 60, 87  
*osgoodi*, *Bryconamericus* 18, 90  
*Copeina* 27, 97  
*Vaimosa* 84, 96  
*oshimai*, *Caranx* 20, 89  
*othonops*, *Hemicetopsis* 44, 89  
*ovalaua*, *Atherina* 14, 87  
*oviceps*, *Archosargus* 11, 103  
*owstoni*, *Alepocephalus* 6, 86  
*Sebastodes* 79, 102  
*oxyrhynchus desotoi*, *Acipenser* 5, 86
- pachacuti*, *Bryconamericus* 18, 90  
*pacificum*, *Diplectrum* 32, 102  
*pacificus*, *Mionorus* 57, 87  
*paegnius*, *Solenostomus* 80, 103  
*palustris*, *Beirabarbus* 16, 94  
*panamensis daguae*, *Hyphessobrycon* 48, 91  
*Eucenogobius* 35, 96  
*Microeleotris* 56, 95  
*Neotrophus* 60, 92  
*Priapichthys* 69, 101  
*panarctus*, *Notropis* 61, 94  
*panostictus*, *Ogcocephalus* 62, 99  
*panuco*, *Gambusia* 37, 101  
*paolence*, *Pygidium* 73, 103  
*papillatus*, *Hoplomyzon* 47, 87  
*papillifer*, *Hemipimelodus* 45, 87  
*papilliferus*, *Spintherobolus* 81, 92  
*papuensis*, *Plotosus* 67, 100  
*paradiseus*, *Gobiesox* 41, 95  
*paraguayensis*, *Aphyocharax* 10, 89  
*Bleptonema* 16, 90  
*Odontostilbe* 61, 91  
*parahybae*, *Cheirodon* 23, 90  
*paranaense*, *Cichlasoma* 26, 92  
*paranahybae*, *Astyanax* 13, 90  
*parismina*, *Gambusia* 37, 101  
*parnaguae*, *Hemiodus* 45, 96  
*parva*, *Anchovia* 8, 95  
*Liza* 53, 98  
*Pseudotatia* 71, 88  
*parvanalis*, *Kryptopterus* 50, 102  
*parvulus*, *Hyphessobrycon* 48, 91  
*parvus*, *Glossogobius* 40, 96  
*Gobius* 42, 96  
*Sphoeroides* 80, 103  
*pascalus*, *Entonanthias* 34, 102  
*patiae*, *Curimatus* 30, 93  
*patzcuaro*, *Chiostoma* 24, 87
- pauciradiatus*, *Antennarius* 9, 86  
*Diretmus* 32, 94  
*Gastromyzon* 38, 97  
*pectoralis*, *Bathypterois* 15, 92  
*pediculatus*, *Pseudancistrus* 70, 98  
*pejeraton*, *Sternopygus* 82, 103  
*pellucidum*, *Characidium* 22, 90  
*peninsulae atrimentis*, *Menidia* 56, 87  
*peruanus ricae*, *Bryconamericus* 18, 90  
*Curimatus* 30, 93  
*Pneumatophorus* 67, 102  
*pescadorensis*, *Liza* 53, 98  
*petrosus*, *Brycon* 19, 90  
*phaeonota*, *Moenkhausia* 57, 91  
*phalacra*, *Brachyglanis* 17, 100  
*phantasia*, *Cetopsorhamdia* 21, 100  
*phenax*, *Plagiotremus* 66, 88  
*pheromystax*, *Lepophidium* 51, 99  
*philippinus*, *Paralepis* 63, 99  
*pholeter*, *Astroblepus* 12, 87  
*phoxocephalum*, *Characidium* 22, 90
- pileatus*, *Diceratias* 32, 94  
*pinnatus*, *Deuterodon* 31, 90  
*piperata*, *Tympanopleura* 84, 86  
*pirrense*, *Cyclopius* 30, 87  
*pittieri*, *Poecilia* 67, 101  
*plagiozonatus*, *Aequidens* 6, 92  
*planiceps*, *Ancistrus* 8, 98  
*planifrons*, *Netuma* 60, 87  
*platycephalum*, *Megalonema* 55, 100  
*plicatus*, *Anostomus* 9, 86  
*poco*, *Adioryx* 5, 97  
*poeciloides*, *Hyphessobrycon* 48, 91  
*poecilus*, *Microglanis* 57, 100  
*polynotatus*, *Corythoichthys* 28, 103  
*polytaenia*, *Leucogobio* 52, 94  
*potamius*, *Gobiesox* 41, 95  
*potaroensis*, *Aequidens* 6, 92  
*Astyanax* 13, 90  
*Deuterodon* 31, 90  
*Myoglanis* 58, 100  
*pricei*, *Emblemariopsis* 34, 89  
*profunda*, *Moenkhausia* 58, 91  
*proops parahybae*, *Pygidium* 73, 103  
*proteus*, *Blenniulus* 16, 88  
*Hyphessobrycon* 48, 91  
*protractile*, *Bivibranchia* 16, 96  
*pseudaplites*, *Micropterus* 57, 89  
*pteroides*, *Characidium* 22, 90  
*puertoricensis*, *Malacotenus* 55, 97  
*pulvereus*, *Zygonectes* 86, 94  
*pumilio*, *Serraniculus* 79, 102  
*pumilus*, *Melamphaes* 55, 98  
*Ogcocephalus* 62, 99  
*punctatissimus*, *Ophioscion* 62, 102  
*punctatum*, *Cichlasoma* 26, 92  
*Hoplosternum* 47, 88  
*Megalonema* 55, 100  
*punctipectophorus*, *Coryphopterus* 28, 96  
*punctulatum piurae*, *Pygidium* 73, 103



- punctulatus*, *Enneapterygius* 34, 104  
*Gastromyzon* 38, 97  
*purpurascens*, *Nesiotes* 60, 101  
*purpurea*, *Botia* 16, 93  
*purpureus*, *archolepis*, *Anoplar-chus* 9, 103  
*pygmaeus*, *Glossanodon* 40, 87  
*pyropunctata*, *Rachovia* 74, 86
- quadrimaculatus*, *Hemiodus* 45, 96  
*quadrizonatus*, *Thalassophryne* 83, 88  
*quercinum japonicum*, *Macrostoma* 54, 99
- radcliffei*, *Spectrunculus* 80, 99  
*randalli*, *Acanthurus* 4, 86  
*Chromis* 25, 101  
*Entomacrodus* 34, 88  
*Myripristis* 59, 97  
*rastrous*, *Caranx* 20, 89  
*regani*, *Astyanax* 13, 90  
*Chirostoma* 24, 87  
*Gambusia* 37, 101  
*Rhamdia* 74, 100  
*reidi*, *Pomacentrus* 68, 101  
*reighardi*, *Leucichthys* 52, 101  
*reinhardti*, *Pygidium* 73, 103  
*remotus*, *Chonerhinus* 24, 103  
*reticulatus*, *Hyphessobrycon* 48, 91  
*Lophiodes* 53, 98  
*retropinnis*, *Leporinodus* 51, 86  
*rhodurus*, *Oncorhynchus* 62, 102  
*rhomaleum*, *Sparisoma* 80, 102  
*rhomboides*, *Hemicaranx* 44, 89  
*ribeirae*, *Astyanax* 13, 90  
*ribeiroi*, *Bellara* 16, 104  
*Glanidium* 39, 88  
*richardi*, *Macrurus* 54, 98  
*richardsoni*, *Ulua* 84, 89  
*riddlei*, *Hemigrammus* 45, 91  
*Holopristis* 47, 91  
*Toxus* 83, 101  
*ridens*, *Gastromyzon* 38, 97  
*ritense*, *Heterogramma* 46, 92  
*rivasi*, *Apuredoras* 10, 95  
*Limia* 52, 101  
*rivuloides*, *Gobio* 41, 94  
*robertsi*, *Zenarchopterus* 85, 96  
*robusta*, *Rhaphiobellone* 75, 88  
*robustum*, *Megalonema* 55, 100  
*robustus*, *Astyanax* 14, 90  
*Leiocassis* 51, 88  
*Notropis* 61, 94  
*Varicorhinus* 85, 94  
*rocasensis*, *Eupomacentrus* 35, 101  
*rodwayi*, *Hemigrammus* 45, 91  
*roosevelti*, *Metynnis* 56, 91  
*Salmo* 77, 102  
*rosacea*, *Melanotaenia* 56, 98  
*rosaceus*, *Hyphessobrycon* 48, 91  
*rosae*, *Chasmocranus* 23, 100  
*rosei*, *Astroblepus* 12, 87  
*roseola*, *Chaenopsis* 21, 89  
*rostellum*, *Ogcocephalus* 62, 99  
*rostratus*, *Sternarchus* 81, 87
- ruberrimus*, *Astyanax* 14, 90  
*rubescens*, *Algansea* 7, 93  
*rubresques*, *Moxostoma* 58, 89  
*rupununi*, *Charax* 22, 90  
*russula*, *Sagamia* 77, 96
- sabana*, *Dangila* 31, 94  
*sabanus*, *Brachyogobius* 17, 95  
*Mystus* 59, 88  
*Ompok* 62, 102  
*sadina*, *Parastremma* 63, 91  
*saepepallens*, *Gobionellus* 41, 96  
*sagamiensis*, *Diaphus* 32, 99  
*sagax*, *Scopelarchus* 78, 102  
*saladonis*, *Notropis* 61, 94  
*saludae*, *Hololepis* 47, 100  
*salvadoris*, *Roeboides* 77, 92  
*sandakanensis*, *Acanthopthalmus* 4, 93  
*Glossogobius* 40, 96  
*sanguinea*, *Vandellia* 84, 104  
*santa*, *Macgregorella* 54, 96  
*santaanae*, *Pantosteus* 63, 89  
*santaeritae*, *Pygidium* 73, 103  
*santaremensis*, *Crenicichla* 29, 92  
*santarosaliae*, *Notropis* 61, 94  
*sardina*, *Melaniris* 56, 87  
*sargentii*, *Pseudoplesiops* 71, 101  
*sarmenii*, *Squaliolus* 81, 103  
*sasakii*, *Liopempheris* 52, 99  
*satsumae*, *Zaescopus* 85, 104  
*sauteri*, *Pristiurus* 69, 102  
*scabripinnis paranae*, *Astyanax* 14, 90  
*schmidtii*, *Otophidium* 63, 99  
*Parahollandia* 63, 103  
*schroederii*, *Bathyclupea* 15, 88  
*schwartzii*, *surinamensis*, *Corydoras* 27, 88  
*scierus apristis*, *Hadropterus* 43, 100  
*Astyanax* 14, 90  
*sciistius*, *Xyrichtys* 85, 97  
*scopiferus guaytarae*, *Bryconamer-icus* 19, 90  
*Bryconamericus* 19, 90  
*scovelli makaxi*, *Syngnathus* 82, 103  
*scythrops*, *Flammeo* 36, 97  
*secundus*, *Microglanis* 57, 100  
*sentaniensis*, *Rhombattractus* 75, 98  
*sepikensis*, *Rhombosoma* 75, 98  
*Zenarchopterus* 85, 96  
*septentrionalis*, *Pygidium* 73, 103  
*septentrionalis*, *Coripareius* 27, 94  
*serrata*, *Pimelodella* 66, 103  
*Rhinosardinia* 75, 93  
*shideleri*, *Moenkhausia* 58, 91  
*shropshirei*, *Euleptoeleotris* 35, 95  
*sialis*, *Amia* 7, 86  
*sicaria*, *Evermannella* 35, 95  
*siebertthaliae*, *Brycon* 19, 90  
*silus*, *Chonerhinus* 24, 103  
*Porogadus* 69, 99  
*simoni*, *Crenicichla* 29, 92  
*simplex*, *Nannostomus* 59, 97  
*simus*, *Creagrutus* 28, 90  
*sinusarabici*, *Dollfusichthys* 32, 93
- snyderi*, *Apogon* 10, 86  
*Hyporhamphus* 49, 96  
*Puntius* 72, 94  
*Taenioides* 82, 96  
*soldatovi minulus*, *Gobio* 41, 94  
*solidus*, *Arius* 11, 87  
*sonorensis*, *Catostomus* 21, 89  
*sparsidens*, *Chromis* 25, 92  
*spilotoceps*, *Epinephelus* 35, 102  
*spilotum*, *Cichlasoma* 26, 92  
*spiniger*, *Spinoblennius* 80, 88  
*spinus*, *Ancistrus* 8, 98  
*stagnatus*, *Rivulus* 76, 86  
*stanleyi*, *Coregonus* 27, 101  
*starksi*, *Menidia* 56, 87  
*starri*, *Pomadasys* 68, 96  
*staurophor*, *Lepophidium* 51, 99  
*stellatum*, *Pygidium* 73, 103  
*stellatus*, *Bodianus* 16, 102  
*stellifer*, *Rivulus* 76, 86  
*stenodon*, *Cheirodon* 23, 90  
*stewarti*, *Argentina* 11, 87  
*Heptapterus* 46, 100  
*Loricaria* 54, 98  
*steyermarki*, *Leporinus* 52, 86  
*stictus*, *Hyphessobrycon* 48, 91  
*stigma*, *Pomacentrus* 68, 101  
*stigmaturus*, *Distichodus* 32, 93  
*straminium*, *Pygidium* 73, 103  
*streber nerensis*, *Zingel* 85, 100  
*striatum*, *Pygidium* 73, 103  
*strophodes*, *Lepidaplois* 51, 97  
*stroudi*, *Gelanoglanis* 38, 88  
*submarginatus*, *Loricaria* 54, 98  
*suensis*, *Microgathus* 57, 103  
*suwae*, *Gnathopogon* 40, 94  
*svetovidovi*, *Belone* 16, 88  
*symmetricus*, *Myripristis* 59, 97
- tabira*, *Acheilognathus* 5, 93  
*tabl*, *Decapterus* 31, 89  
*taenia melanoleuca*, *Cobitis* 27, 93  
*taeniatum pertense*, *Heterogramma* 46, 92  
*taiwanensis*, *Pseudobagrus* 70, 88  
*taiwanus*, *Rhinogobius* 75, 96  
*tamuisiensis*, *Scaphesthes* 78, 94  
*tanakae*, *Lycodes* 54, 104  
*Zesticthys* 85, 104  
*tanakai*, *Caranx* 20, 89  
*tarpon*, *Genycharax* 38, 90  
*taurocephalus*, *Hyphessobrycon* 48, 91  
*tegatus*, *Vesicatrax* 85, 92  
*temminckii*, *Alectis* 6, 89  
*templei*, *Pomadasys* 69, 96  
*tenuis*, *Cichlasoma* 26, 92  
*tenuis*, *Callionymus* 20, 88  
*Cynodonichthys* 30, 86  
*Pellonula* 64, 93  
*Poecilia* 68, 101  
*terminalis*, *Parodon* 64, 99  
*tetranema*, *Bathylabrus* 15, 88  
*thamalakanensis*, *Alestes* 6, 89  
*thomersoni*, *Rhinodoras* 75, 95  
*thompsoni*, *Sebastodes* 79, 102

*thurni*, *Roeboides* 77, 92  
*tigris*, *Alepideleotris* 6, 95  
*tiki*, *Myripristis* 59, 97  
*timagamiensis*, *Salvelinus* 77, 102  
*tlahuacensis*, *Evarra* 35, 94  
*tolimae*, *Bryconamericus* 19, 90  
*toni posteroventralis*, *Barbatula* 15, 93  
*tosae*, *Zalescopus* 85, 104  
*toweri*, *Goodea* 42, 96  
*townsendi*, *Parascloopsis* 63, 99  
*transvestitus*, *Nanochromis* 59, 92  
*tremeniinae*, *Odontostilbe* 61, 91  
*trewavasae*, *Sciaena* 78, 102  
*triangulatus*, *Malacoctenus* 55, 97  
*tricuspidata*, *Atuona* 14, 95  
*trifasciatum maciliense*, *Heterogramma* 46, 92  
*Cyclopium* 30, 87  
*triguttatum*, *Pygidium* 73, 103  
*tristegus marquesensis*, *Acanthurus* 4, 86  
*tropicus*, *Platyopocilus* 67, 101  
*troscheli*, *Glyphisodon* 40, 101  
*tsotsorogensis*, *Barbus* 15, 94  
*tsuchigae*, *Gnathopogon* 41, 94  
*tuba*, *Cichlasoma* 26, 92  
*tubbi*, *Pangasius* 63, 99  
*tuckeri*, *Monacanthus* 58, 88  
*tumacoensis*, *Umbrina* 84, 102  
*turrubarensis*, *Gambusia* 37, 101  
*tutuillae*, *Centropyge* 21, 101  
*tuyra*, *Arius* 11, 87  
*tuyrense*, *Cichlasoma* 26, 92  
*tuyrensis*, *Loricaria* 54, 98  
  
*iii*, *Caranx* 20, 89  
*ulae*, *Synodus* 82, 103  
*umalii*, *Puntius* 72, 94  
*umbratilis*, *Gambusia* 38, 101  
*umbriceps*, *Alepocephalus* 6, 86  
*umbriferum*, *Cichlasoma* 26, 92  
*undecimalis*, *Myripristis* 59, 97  
*unicolor*, *Echidna* 33, 99  
*unicuspis*, *Ichthyomyzon* 49, 100  
*undorsalis*, *Helogenes* 43, 96  
*unifasciatum*, *Cyclopium* 30, 87

*uracantha rupestre*, *Loricaria* 54, 98  
*urophthalmus mayorum*, *Cichlasoma* 26, 92  
*urosipilotus*, *Gymnapogon* 42, 87  
*uruguayanae*, *Macropsobrycon* 54, 91  
*ushiei*, *Dasyatis* 31, 94  
*uxorius*, *Bombonia* 16, 103  
  
*vaivasensis*, *Ambassis* 7, 89  
*vanceae*, *Cyclopium* 30, 87  
*variabilis*, *Starksia* 81, 97  
*variatus*, *Platyopocilus* 67, 101  
*variegata venezuelae*, *Loricaria* 54, 98  
*Skiffia* 80, 96  
*variegatus*, *Microgolanis* 57, 100  
*velutinus*, *Hemipimelodus* 45, 87  
*venezuelae*, *Ophioscion* 62, 102  
*ventrale*, *Cyclopium* 30, 87  
*verecundus*, *Spirinchus* 81, 99  
*vermiculatum*, *Pygidium* 73, 103  
*verrucosus*, *Antennarius* 9, 86  
*villadolidi*, *Ctenogobius* 29, 96  
*Chromis* 25, 101  
*villiersi*, *Physalia* 65, 102  
*villosus*, *Lithogenes* 53, 98  
*Pseudopimelodus* 71, 100  
*vineta*, *Echidna* 33, 99  
*vintoni*, *Characidium* 22, 90  
*virgata*, *Albula* 6, 86  
*virgatulus*, *Ctenogobius* 29, 96  
*viridesens*, *Gunnellichthys* 42, 98  
*vittata*, *Gambusia* 38, 101  
*volcanus*, *Rivulus* 76, 86  
  
*waigiensis*, *Ctenogobius* 29, 96  
*Enneapterygius* 34, 104  
*waimacui*, *Rivulus* 76, 86  
*wakiyae*, *Malakichthys* 55, 100  
*walensis*, *Salarias* 77, 88  
*waltersi*, *Batrachoides* 15, 88  
*weberi*, *Chromis* 25, 101  
*weitzmani*, *Corydoras* 28, 88  
*whaleri*, *Gephyrocharax* 39, 90  
*whitei*, *Goodea* 42, 96  
*wichmanni*, *Apogon* 10, 87

*wilburi*, *Callionymus* 20, 88  
*Macrodonogobius* 54, 96  
*williamsi*, *Anisotremus* 9, 96  
*wilsoni*, *Hemiancistrus* 43, 98  
*Phenagoniastes* 65, 91  
*wonderi*, *Gobiocichla* 41, 92  
*woodsi*, *Anthias* 9, 102  
*Chromis* 25, 101  
*Gobiopsis* 41, 96  
*Microdesmus* 56, 98  
*Myripristis* 59, 97  
*Rhynchodoras* 75, 95

*x-punctata trautmani*, *Hybopsis* 47, 94

*x-punctata*, *Hybopsis* 47, 94  
*xantherythrus*, *Holocentrus* 47, 97  
*xanthomelas*, *Brachygobius* 17, 95  
*xanthosoma*, *Gambusia* 38, 101  
*xanthum*, *Megalonema* 55, 100  
*xenandrus*, *Engyprosope* 34, 88  
*xenicus*, *Calymnichthys* 20, 89

*y-lineata*, *Starksia* 81, 97  
*yamamotis*, *Moroco* 58, 94  
*yarkandensis sellaefer*, *Barbatula* 15, 93  
*yarrowi*, *Agosia* 6, 93  
*yoshiae*, *Anarchias* 7, 99  
*yucatana australis*, *Gambusia* 38, 101

*zacae*, *Malacoctenus* 55, 97  
*zanectes*, *Calliurichthys* 20, 88  
*zebra*, *Characidium* 22, 90  
*zebrinus*, *Phrynelox* 65, 86  
*zeteki*, *Bryconamericus* 19, 90  
*zirahuen*, *Chirostoma* 24, 87  
*zonatum*, *Cichlasoma* 26, 92  
*Pygidium* 73, 103  
*zonatus*, *Bostrychus* 16, 95  
*zonifer sudensis*, *Malacoctenus* 55, 97  
*zonophaea*, *Echidna* 33, 99  
*zuloagai*, *Doraops* 32, 95

---

---

# FIELDIANA

---

---

## Zoology

NEW SERIES, NO. 35

### **Catalogue of Type Specimens of Recent Fishes in Field Museum of Natural History**

**Myriam Ibarra**

**Donald J. Stewart**

**September 30, 1987**

**Publication 1377**

---

**PUBLISHED BY FIELD MUSEUM OF NATURAL HISTORY**

---



Field Museum of Natural History  
Roosevelt Road at Lake Shore Drive  
Chicago, Illinois 60605-2496  
Telephone: (312) 922-9410







UNIVERSITY OF ILLINOIS URBANA

590.5FIN S

COO1

FIELDIANA ZOOLOGY \$ NEW SERIES \$CHGO

29 16 1986 87



3 0112 009378719